

# Long excursions of a random walk

Endre Csáki\*, Pál Révész\* and Zhan Shi

Alfréd Rényi Institute of Mathematics, Technische Universität Wien and Université Paris VI

**Summary.** In [1] and [9] it was proved that the length of the longest excursion among the first  $n$  excursions of a plane random walk is nearly equal to the total sum of the lengths of these excursions. In this paper several results are proved in the same spirit, for plane random walks and for random walks in higher dimensions.

**Keywords.** Random walk, excursion length.

**2000 Mathematics Subject Classification.** 60G50.

## 1. Introduction

Consider a simple symmetric random walk on the lattice  $\mathbb{Z}^2$ . This means that if the moving particle is in  $x \in \mathbb{Z}^2$  at the moment  $n$  then at the moment  $n + 1$  the particle moves with equal probabilities to any of the 4 neighbours of  $x$  independently of how the particle achieved  $x$ . Let  $S_n$  be the location of the particle after  $n$  steps and assume that  $S_0 = 0$ . Introduce the following notations:

- (i)  $\xi(x, n) := \#\{k : 0 < k \leq n, S_k = x\}$ , ( $x \in \mathbb{Z}^2$ ,  $n = 1, 2, \dots$ )
- (ii)  $\xi(x, t)$  ( $x \in \mathbb{Z}^2$ ,  $t \geq 0$ ) is the continuous process obtained by linear interpolation from  $\xi(x, n)$  with  $\xi(x, 0) = 0$ ,
- (iii)  $\rho_0 := 0$ ,  
 $\rho_k := \min\{n : n > \rho_{k-1}, S_n = 0\}$ ,  $k \geq 1$ ,
- (iv)  $r(k) := \rho_k - \rho_{k-1}$ ,

---

\*Research supported by the Hungarian National Foundation for Scientific Research, Grant No. T 029621.

(v)  $M_n(1) \geq M_n(2) \geq \dots \geq M_n(\xi(0, n) + 1)$  are the order statistics of the sequence

$$r(1), r(2), \dots, r(\xi(0, n)), n - \rho_{\xi(0, n)},$$

(vi)  $\mathcal{H}$  is the set of functions

$$h(m_1, m_2, \tau, x) = \begin{cases} m_1 & \text{if } 0 \leq x < \tau, \\ m_2 & \text{if } \tau \leq x \leq 1, \end{cases}$$

where  $0 \leq m_1 \leq m_2 \leq 1$ ,  $0 \leq \tau \leq 1$ .

In words,  $\xi(x, n)$  counts the number of visits to the site  $x$  during the first  $n$  steps and is called the local time of the random walk;  $\rho_n$ , which denotes the time of the  $n$ -th return to zero, is the “inverse” of  $\xi(0, k)$  in some sense. The sections between two consecutive returns are called excursions and so  $\{r(k), k = 1, 2, \dots\}$  denotes the sequence of the lengths of the excursions. Clearly, the sequence  $\{M_n(1), \dots\}$  denotes the ordered lengths of the excursions. In this paper we study some properties of the “long” excursions, which in spite of the recurrence of the random walk behaves quite differently from the case of dimension 1.

In [1] we proved the following Strassen type

**THEOREM A.** *With probability one, the set of limit points of the sequence*

$$\left\{ \frac{\pi \xi(0, nt)}{(\log n) \log_3 n}, 0 \leq t \leq 1 \right\}$$

is  $\mathcal{H}$ .

Here and in what follows,  $\log n := \log \max(n, e)$ , and  $\log_p n$  is the  $p$ -th iteration of  $\log$ . The proof of Theorem A is based on the following

**LEMMA A.**

$$\lim_{n \rightarrow \infty} \frac{M_n(1) + M_n(2)}{n} = 1 \quad \text{a.s.}$$

We note that by the proof of Theorem A (cf. [1]) we easily get the following

**THEOREM 1.** *With probability one, the set of limit points of the sequence*

$$\left\{ \frac{\xi(0, nt)}{\xi(0, n)}, 0 \leq t \leq 1 \right\}$$

is  $\mathcal{H}$ .

Lemma A easily implies

**PROPOSITION A.**  $\rho_{n+1} - \rho_n$  is either much smaller or much bigger than  $\rho_n$ .

In order to give a more accurate form of Proposition A, let  $\{\alpha_n\}$  and  $\{\beta_n\}$  be sequences of positive numbers with

$$(1.1) \quad \sum_{n=1}^{\infty} \frac{1}{n\alpha_n} < \infty, \quad \alpha_n \geq \log_2 n,$$

$$(1.2) \quad \sum_{n=1}^{\infty} \frac{1}{n\beta_n} = \infty, \quad \inf_{n \geq 1} \beta_n > 0.$$

Further let

$$(1.3) \quad a_n = \exp\left(\frac{n}{\alpha_n}\right), \quad b_n = \exp\left(\frac{n}{\beta_n}\right), \quad c_n = \exp\left(\frac{n}{(1+\varepsilon)\beta_n}\right) \quad (\varepsilon > 0).$$

Finally let

$$T_n = \frac{\rho_{n+1} - \rho_n}{\rho_n}.$$

Then we have

**THEOREM 2.**

$$(i) \quad T_n \notin (a_n^{-1}, a_n) \quad \text{a.s.}$$

for all but finitely many  $n$ ,

$$(ii) \quad T_n \in (b_n^{-1}, c_n^{-1}) \quad \text{i.o. a.s.}$$

$$(iii) \quad T_n \in (c_n, b_n) \quad \text{i.o. a.s.}$$

**EXAMPLE 1.**

$$\alpha_n = (\log n)^{1+\delta} \quad (\delta > 0), \quad \beta_n = \log n$$

i.e.,

$$a_n = \exp\left(\frac{n}{(\log n)^{1+\delta}}\right), \quad b_n = \exp\left(\frac{n}{\log n}\right), \quad c_n = \exp\left(\frac{n}{(1+\varepsilon)\log n}\right)$$

satisfy the above conditions.

Theorem 2 clearly implies:

**COROLLARY 1.** *Let*

$$T_n^* = \frac{\rho_{n+1} - \rho_n}{\rho_{n+1}} \quad (n = 0, 1, 2, \dots).$$

Then

$$T_n^* \notin (a_n^{-1}, 1 - a_n^{-1}) \quad \text{a.s.},$$

for all but finitely many  $n$ ,

$$T_n^* \in (0, c_n^{-1}), \quad \text{i.o. a.s.}$$

$$T_n^* \in (1 - c_n^{-1}, 1) \quad \text{i.o. a.s.}$$

Lemma A also implies that  $\rho_{n+k} - \rho_n$  for any  $k = 1, 2, \dots$ , is either much bigger or much smaller than  $\rho_n$ . A refined form of this statement is given in the next corollary.

**COROLLARY 2.** *Let  $\{a_n\}$  be defined as above and assume that  $\{a_n\}$  is a non-decreasing sequence. Further let*

$$\begin{aligned} a_n^* &= \exp\left(\sum_{j=n}^{\infty} a_j^{-1}\right) - 1, \\ T_n^{(k)} &= \frac{\rho_{n+k} - \rho_n}{\rho_n}, \quad (k = 1, 2, \dots) \\ B_n &= \bigcup_{k=1}^{\infty} \{T_n^{(k)} \in (a_n^*, a_n)\}. \end{aligned}$$

Then among the events  $B_n$  only finitely many might occur with probability 1.

Note that if

$$a_n = \exp\left(\frac{n}{(\log n)^{1+\delta}}\right) \quad (\delta > 0)$$

then

$$a_n^* \leq \exp\left(-\frac{n}{(\log n)^{1+2\delta}}\right).$$

The above results also imply that the waiting time

$$\nu(n) = \min\{k : k \geq n, \rho_{k+1} - \rho_k \geq \rho_n\}$$

is finite with probability 1 for any finite  $n$ . In fact we prove

**THEOREM 3.** *Let  $\{f_n\}$  be a positive sequence such that  $n \mapsto f_n/n$  is non-decreasing, then*

$$\limsup_{n \rightarrow \infty} \frac{\nu(n)}{f_n} = \begin{cases} 0 \\ \infty \end{cases}, \text{ a.s.} \iff \sum_n \frac{1}{f_n} \begin{cases} < \infty \\ = \infty \end{cases}.$$

Corollary 1 claims that  $T_n^*$  is either nearly 0 or nearly 1. At the same time Theorem 3 suggests that between  $n$  and  $2n$  there exists about 1 integer  $k$  for which  $T_k^*$  is nearly 1. This idea is formulated in our following

**THEOREM 4.**

- (i)  $\lim_{n \rightarrow \infty} \frac{1}{\log n} \sum_{k=0}^n T_k^* = 1 \quad \text{a.s.},$
- (ii)  $\lim_{n \rightarrow \infty} \mathbf{P} \left\{ \frac{\sum_{k=0}^n T_k^* - \log n}{\sqrt{\log n}} \leq x \right\} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-u^2/2} du,$
- (iii)  $\limsup_{n \rightarrow \infty} \frac{\sum_{k=0}^n T_k^* - \log n}{\sqrt{2(\log n)(\log_3 n)}} = 1 \quad \text{a.s.}$

Now consider a simple symmetric random walk on the lattice  $\mathbb{Z}^d$ , which means that at each step, the moving particle visits each of its  $(2d)$  neighbours (on  $\mathbb{Z}^d$ ) with equal probability  $1/(2d)$ . Let  $S_n$  be the location of the particle after  $n$  steps and assume that  $S_0 = 0$ . Since the random walk in  $\mathbb{Z}^d$ ,  $d \geq 3$ , is transient, the above results cannot be true. However if we consider the longest excursion away from some  $x \in \mathbb{Z}^d$  completed by the time  $n$ , then it can be long. For  $i \geq 0$ , define the random variable  $\kappa(i)$  by

$$S_{i+j} \neq S_i, \quad j = 1, 2, \dots, \kappa(i) - 1, \quad S_{i+\kappa(i)} = S_i.$$

(If such  $\kappa(i)$  does not exist, we set  $\kappa(i) := \infty$ ). Let

$$R(n) := \max\{\kappa(i) : i + \kappa(i) \leq n\},$$

which in words denotes the length of the longest completed excursion (away from any point) at time  $n$ .

**THEOREM 5.** *Let  $d \geq 3$ . With probability one,*

$$\lim_{n \rightarrow \infty} \frac{\log R(n)}{\log n} = \begin{cases} 1 & \text{if } d = 3, 4, \\ \frac{2}{d-2} & \text{if } d \geq 5. \end{cases}$$

## 2. Preliminary results

**THEOREM B.** ([2]) *For any  $\delta > 0$  we have*

$$\exp\left((1 - \delta)\frac{\pi n}{\log_2 n}\right) \leq \rho_n \leq \exp(n(\log n)^{1+\delta}) \quad \text{a.s.},$$

*for all but finitely many  $n$ . When  $n$  goes to infinity,*

$$\mathbf{P}\{\rho_1 > n\} = \mathbf{P}\{\xi(0, n) = 0\} = \pi(\log n)^{-1} + O((\log n)^{-2}),$$

*and*

$$\mathbf{P}\{\xi(0, n) \geq x \log n\} = e^{-\pi x}(1 + O((\log n)^{-1/4}))$$

*uniformly for  $x \in (0, (\log n)^{3/4})$ .*

Since  $\{\rho_n \leq k\} = \{\xi(0, k) \geq n\}$ , Theorem B implies

**PROPOSITION B.** *For any  $2 \leq m < n$  we have*

$$\left| \mathbf{P}\{m < \rho_1 < n\} - \frac{\pi \log(n/m)}{(\log n) \log m} \right| \leq C \left( \frac{1}{(\log m)^2} + \frac{1}{(\log n)^2} \right)$$

where  $C > 0$  is an absolute constant. Furthermore,

$$\mathbf{P}\{\rho_n \leq e^{n/z}\} = e^{-\pi z}(1 + O(n^{-1/7}))$$

uniformly for  $z \in (0, n^{3/7})$  and

$$\mathbf{P}\{\rho_n > e^{n/z}\} \leq C(1 - e^{-\pi z}) \leq C\pi z, \quad z \in [n^{-1/7}, 1].$$

Similarly for any  $0 \leq z_1 < z_2 < n^{3/7}$  and  $z_2 - z_1 \leq 1$  we have

$$\left| \mathbf{P} \left\{ \exp\left(\frac{n}{z_2}\right) \leq \rho_n < \exp\left(\frac{n}{z_1}\right) \right\} - \pi e^{-\pi z_1}(z_2 - z_1) \right| \leq \frac{C}{n^{1/7}}.$$

In the proof of Theorem 4 we use results of Rényi [6], [7] concerning extreme elements. Let  $\{\xi_k\}_{k=1}^\infty$  be a sequence of i.i.d. random variables having continuous distribution.  $\xi_k$  is called an extreme element if  $\xi_k > \max_{1 \leq j \leq k-1} \xi_j$ . By convention,  $\xi_1$  is counted as an extreme element. Let  $\mu_n$  be the number of extreme elements among  $\xi_1, \dots, \xi_n$ .

**THEOREM C.** ([6], [7])

- (i)  $\lim_{n \rightarrow \infty} \frac{\mu_n}{\log n} = 1 \quad \text{a.s.},$
- (ii)  $\lim_{n \rightarrow \infty} \mathbf{P} \left\{ \frac{\mu_n - \log n}{\sqrt{\log n}} \leq x \right\} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-u^2/2} du,$
- (iii)  $\limsup_{n \rightarrow \infty} \frac{\mu_n - \log n}{\sqrt{2(\log n)(\log_3 n)}} = 1 \quad \text{a.s.}$

In the proof of Theorem 5, we need the following two results concerning the simple symmetric random walk on  $\mathbb{Z}^d$  ( $d \geq 3$ ).

**THEOREM D.** ([4]) *Let  $d \geq 3$ . Then as  $n \rightarrow \infty$*

$$\mathbf{P}\{\rho_1 = 2n\} \sim (1 - \gamma(d))^2 \mathbf{P}\{S_{2n} = 0\},$$

where  $\gamma(d) = \mathbf{P}\{\rho_1 < \infty\}$ .

**THEOREM E.** ([3]) *Let  $f(n) \uparrow \infty$  be a positive integer valued function and let  $E_n$  be the event that the paths*

$$\{S_0, S_1, \dots, S_n\} \quad \text{and} \quad \{S_{n+f(n)+1}, S_{n+f(n)+2}, \dots\}$$

*have points in common. Then*

(i) *for  $d = 3$ , if  $f(n) = n(\varphi(n))^2$  and  $\varphi(n)$  is non-decreasing, then*

$$(2.1) \quad \mathbf{P}\{E_n \text{ i.o.}\} = 0 \quad \text{or} \quad 1$$

depending on whether  $\sum_{k=1}^{\infty}(\varphi(2^k))^{-1}$  converges or diverges,

(ii) for  $d = 4$ , if  $f(n) = n\psi(n)$  and  $\psi(n)$  is non-decreasing, then we have (2.1) depending on whether  $\sum_{k=1}^{\infty}(k\psi(2^k))^{-1}$  converges or diverges,

(iii) for  $d \geq 5$ , if

$$\sup_{m \geq n} \frac{f(m)}{m} \geq C \frac{f(n)}{n}$$

(for some  $C > 0$ ) then we have (2.1) depending on whether  $\sum_{k=1}^{\infty}(f(k))^{(2-d)/d}$  converges or diverges.

### 3. Proof of Theorem 2

Let  $\{\alpha_n\}$  and  $\{\beta_n\}$  satisfy (1.1) and (1.2), respectively, and let  $\{a_n\}$ ,  $\{b_n\}$  and  $\{c_n\}$  be as in (1.3). Fix a constant  $\delta \in (0, 1/3)$ . Introduce the following notations:

$$\begin{aligned} d_n &= \exp\left(\left(1 - \delta\right)\frac{\pi n}{\log_2 n}\right), \\ e_n &= \exp(n(\log n)^{1+\delta}), \\ A_n^{(1)} &= \{T_n \in (a_n^{-1}, a_n)\}, \\ A_n^{(2)} &= \{T_n \in (b_n^{-1}, c_n^{-1})\}, \\ A_n^{(3)} &= \{T_n \in (c_n, b_n)\}. \end{aligned}$$

Clearly we have

$$A_n^{(1)} = \{A_n^{(1)} \cap \mathcal{B}_n\} \cup \{A_n^{(1)} \cap \mathcal{B}_n^c\},$$

where  $\mathcal{B}_n := \{\rho_n \in (d_n, e_n)\}$  and  $\mathcal{B}_n^c$  denotes its complementary. By Theorem B among the events

$$A_n^{(i)} \cap \mathcal{B}_n^c,$$

only finitely many might occur with probability 1.

Another application of Theorem B yields that

$$\begin{aligned} \mathbf{P}\{A_n^{(1)} \cap \mathcal{B}_n\} &= \mathbf{E}\left[\mathbf{1}_{\mathcal{B}_n} \mathbf{P}\{A_n^{(1)} \mid \rho_n\}\right] = \\ &= \mathbf{E}\left[\mathbf{1}_{\mathcal{B}_n} \left(\frac{2\pi \log a_n}{\log(a_n^{-1}\rho_n) \cdot \log(a_n\rho_n)} + O\left(\frac{1}{\log^2(a_n^{-1}\rho_n)}\right)\right)\right]. \end{aligned}$$

On  $\mathcal{B}_n$ , we have (recalling that  $\alpha_n \geq \log_2 n$ )

$$(3.1) \quad \log(a_n^{-1}\rho_n) = \log \rho_n - \frac{n}{\alpha_n} \geq (1 - \delta)\frac{\pi n}{\log_2 n} - \frac{n}{\log_2 n} \geq \frac{n}{\log_2 n},$$

so that

$$\mathbf{E} \left[ \mathbf{1}_{\mathcal{B}_n} \cdot O \left( \frac{1}{\log^2(a_n^{-1} \rho_n)} \right) \right] = O \left( \frac{(\log_2 n)^2}{n^2} \right).$$

On the other hand, since (3.1) also tells that  $\log(a_n^{-1} \rho_n) \geq (\log \rho_n)/2$  on  $\mathcal{B}_n$ , we have

$$\mathbf{E} \left[ \mathbf{1}_{\mathcal{B}_n} \cdot \frac{\log a_n}{\log(a_n^{-1} \rho_n) \cdot \log(a_n \rho_n)} \right] \leq 2(\log a_n) \mathbf{E} \left[ \frac{\mathbf{1}_{\mathcal{B}_n}}{(\log \rho_n)^2} \right],$$

which, by writing  $N := \lfloor (\log_2 n) / ((1 - \delta)\pi) \rfloor$  and using Proposition B, is

$$\begin{aligned} &\leq 2(\log a_n) \sum_{i=0}^N \mathbf{E} \left[ \frac{\mathbf{1}_{\{\exp(n/(i+1)) \leq \rho_n < \exp(n/i)\}}}{(n/(i+1))^2} \right] = \\ &= 2(\log a_n) \sum_{i=0}^N \frac{\pi \exp(-\pi i) + O(n^{-1/7})}{(n/(i+1))^2} = \\ &= O \left( \frac{\log a_n}{n^2} \right) = \\ &= O \left( \frac{1}{n\alpha_n} \right). \end{aligned}$$

Therefore,

$$\mathbf{P} \left\{ A_n^{(1)} \cap \mathcal{B}_n \right\} = O \left( \frac{(\log_2 n)^2}{n^2} \right) + O \left( \frac{1}{n\alpha_n} \right),$$

which is summable for  $n$ . We obtain (i) of Theorem 2 by an application of the Borel–Cantelli lemma.

Now, we turn to the proof of (ii) and (iii). We first assume that for some  $n_0$ ,

$$\beta_n \leq (\log n)^2, \quad n \geq n_0.$$

We fix a constant  $\theta > 1/(\inf_{n \geq 1} \beta_n)$ . Introduce the following notations:

$$\begin{aligned} \mathcal{E}_n &= \{\rho_n \in (e^{\theta n}, e^{2\theta n})\}, \\ p_n^{(2)}(x) &= \mathbf{P} \left\{ \frac{\rho_1}{x} \in (b_n^{-1}, c_n^{-1}) \right\}, \\ p_n^{(3)}(x) &= \mathbf{P} \left\{ \frac{\rho_1}{x} \in (c_n, b_n) \right\}. \end{aligned}$$

We have, for  $i = 2$  or  $3$ ,

$$\mathbf{P} \left\{ \mathcal{E}_n \cap A_n^{(i)} \right\} = \mathbf{E} \left[ \mathbf{1}_{\mathcal{E}_n} \mathbf{P} \left\{ A_n^{(i)} \mid \rho_n \right\} \right] = \mathbf{E} \left[ \mathbf{1}_{\mathcal{E}_n} \cdot p_n^{(i)}(\rho_n) \right].$$

By Proposition B, there exist two positive constants  $C_1$  and  $C_2$  such that for all  $n$ ,

$$(3.2) \quad \frac{C_1}{n\beta_n} \leq p_n^{(i)}(e^{nz}) \leq \frac{C_2}{n\beta_n}, \quad i = 2 \text{ or } 3,$$



uniformly for  $z \in (\theta, 2\theta)$ . Therefore,

$$\frac{C_1}{n\beta_n} \mathbf{P}\{\mathcal{E}_n\} \leq \mathbf{P}\{\mathcal{E}_n \cap A_n^{(i)}\} \leq \frac{C_2}{n\beta_n} \mathbf{P}\{\mathcal{E}_n\}.$$

Since  $C_3 \leq \mathbf{P}\{\mathcal{E}_n\} \leq C_4$  for some positive constants  $C_3$  and  $C_4$ , we obtain:

$$(3.3) \quad \frac{C_5}{n\beta_n} \leq \mathbf{P}\{\mathcal{E}_n \cap A_n^{(i)}\} \leq \frac{C_6}{n\beta_n}.$$

As a consequence,  $\sum_n \mathbf{P}\{\mathcal{E}_n \cap A_n^{(i)}\} = \infty$ , for  $i = 2$  or  $3$ .

In order to complete the proof of (ii) and (iii) of Theorem 2 (in case  $\beta_n \leq (\log n)^2$ ), we prove that the events  $A_n^{(i)}$  are nearly pairwise independent. Let  $0 < n < N$ . Then we have

$$\mathbf{P}\{A_n^{(i)} \mathcal{E}_n A_N^{(i)} \mathcal{E}_N\} = \mathbf{E} \left[ \mathbf{1}_{A_n^{(i)} \mathcal{E}_n \mathcal{E}_N} \cdot p_N^{(i)}(\rho_N) \right],$$

which, according to (3.2), is

$$\leq C_2 \mathbf{E} \left[ \frac{\mathbf{1}_{A_n^{(i)} \mathcal{E}_n \mathcal{E}_N}}{N\beta_N} \right] \leq \frac{C_2}{N\beta_N} \mathbf{P}\{A_n^{(i)} \mathcal{E}_n\}.$$

This, jointly considered with (3.3) and the fact that  $\sum_n \mathbf{P}\{\mathcal{E}_n \cap A_n^{(i)}\} = \infty$ , yields

$$\liminf_{m \rightarrow \infty} \frac{\sum_{n=1}^m \sum_{N=1}^m \mathbf{P}\{A_n^{(i)} \mathcal{E}_n A_N^{(i)} \mathcal{E}_N\}}{\left( \sum_{n=1}^m \mathbf{P}\{A_n^{(i)} \mathcal{E}_n\} \right)^2} < \infty.$$

According to the Kochen and Stone's version ([5]) of the Borel–Cantelli lemma, we have, for  $i = 2$  or  $3$ ,  $\mathbf{P}\{A_n^{(i)} \mathcal{E}_n \text{ i.o.}\} > 0$ . Consequently,

$$\mathbf{P}\{A_n^{(i)} \text{ i.o.}\} > 0.$$

Since the event  $\{A_n^{(i)} \text{ i.o.}\}$  is invariant under any finite permutation between the i.i.d.r.v.'s  $\{\rho_{n+1} - \rho_n\}_{n=0}^\infty$ , an immediate application of the Hewitt–Savage zero-one law yields that  $\mathbf{P}\{A_n^{(i)} \text{ i.o.}\} = 1$  for  $i = 2$  or  $3$ . This completes the proof of Theorem 2 (ii) and (iii) in case  $\beta_n \leq (\log n)^2$ .

For general  $\{\beta_n\}$ , we define

$$\tilde{\beta}_n := \min(\beta_n, (\log n)^2),$$

which satisfies (1.2), such that  $\tilde{\beta}_n \leq (\log n)^2$ . According to what we have just proved, with probability one,

$$(3.4) \quad T_n \in \left( \exp\left(\frac{n}{(1+\varepsilon)\tilde{\beta}_n}\right), \exp\left(\frac{n}{\tilde{\beta}_n}\right) \right),$$

along a (random) subsequence  $\{n(i)\}_{i \geq 1}$ . If there existed a sub-subsequence of  $\{n(i)\}_{i \geq 1}$ , say  $\{n(i(k))\}_{k \geq 1}$ , such that  $\beta_{n(i(k))} > \log^2(n(i(k)))$  for  $k \geq 1$ , then  $\tilde{\beta}_{n(i(k))} = \log^2(n(i(k)))$ , and according to (i) of Theorem 2,

$$T_{n(i(k))} \notin \left( \exp \left( -\frac{n(i(k))}{\tilde{\beta}_{n(i(k))}} \right), \exp \left( \frac{n(i(k))}{\tilde{\beta}_{n(i(k))}} \right) \right),$$

for all large  $k$ . This would contradict (3.4). As a consequence,  $\beta_{n(i)} \leq \log^2(n(i))$  for all large  $i$ , say  $i \geq i_0$ . Thus  $\tilde{\beta}_{n(i)} = \beta_{n(i)}$  for  $i \geq i_0$ , which in view of (3.4) implies

$$T_{n(i)} \in \left( \exp \left( \frac{n(i)}{(1+\varepsilon)\tilde{\beta}_{n(i)}} \right), \exp \left( \frac{n(i)}{\tilde{\beta}_{n(i)}} \right) \right), \quad i \geq i_0.$$

This proves Theorem 2 (iii). A similar argument yields Theorem 2 (ii) for general  $\{\beta_n\}$ .  $\square$

Corollary 2 follows easily from Theorem 2 and the next lemma.

**LEMMA 1.** *Let  $\{g_n\}$  be a positive non-decreasing sequence such that  $\sum_n (1/g_n) < \infty$ . If  $\{\rho_n\}_{n=1}^\infty$  is an increasing sequence with  $\rho_0 := 0$ , satisfying*

$$\frac{\rho_{n+1} - \rho_n}{\rho_n} \notin (g_n^{-1}, g_n), \quad n \geq n_0,$$

then

$$\frac{\rho_{n+k} - \rho_n}{\rho_n} \notin (g_n^*, g_n), \quad n \geq n_0, \quad k \geq 1,$$

where  $g_n^* := \exp \left( \sum_{j=n}^\infty g_j^{-1} \right) - 1$ .

## 4. Proof of Theorem 3

At first we prove two lemmas.

**LEMMA 2.** *For all integers  $n \geq 1$  and  $a \geq 2n$ ,*

$$\mathbf{P}\{\nu(n) > a\} \leq \frac{Cn}{a} + \frac{C}{n^{1/7}},$$

where  $C$  is an absolute, positive constant.

**Proof.** Clearly we have

$$\mathbf{P}\{\nu(n) > a\} = \mathbf{P}\{r(n+1) < \rho_n, \dots, r(a) < \rho_n\}.$$

Since  $r(n+1), \dots, r(a), \rho_n$  are independent, we get

$$\mathbf{P}\{\nu(n) > a\} = \mathbf{E}(F^{a-n}(\rho_n)),$$

where  $F(x) = \mathbf{P}\{\rho_1 < x\}$  is the distribution function of  $\rho_1$ . By Theorem B,  $F(x) \leq 1 - C_7/\log x$  for all  $x \geq 2$  if  $C_7 > 0$  is small enough. Hence

$$\begin{aligned} \mathbf{P}\{\nu(n) > a\} &\leq \mathbf{E} \left( 1 - \frac{C_7}{\log \rho_n} \right)^{a-n} \leq \mathbf{E} \exp \left( -\frac{C_7(a-n)}{\log \rho_n} \right) \leq \\ &\leq \exp \left( -\frac{C_7(a-n)}{n} \right) + \mathbf{P}\{\log \rho_n > n^{8/7}\} + \\ &\quad + \mathbf{E} \left( \exp \left( -\frac{C_7(a-n)}{\log \rho_n} \right) \mathbf{1}_{\{n \leq \log \rho_n \leq n^{8/7}\}} \right). \end{aligned}$$

Proposition B with  $z = n^{-1/7}$  gives

$$\mathbf{P}\{\log \rho_n > n^{8/7}\} \leq \frac{C_8}{n^{1/7}}.$$

Integration by parts yields

$$\begin{aligned} &\mathbf{E} \left( \exp \left( -\frac{C_7(a-n)}{\log \rho_n} \right) \mathbf{1}_{\{n \leq \log \rho_n \leq n^{8/7}\}} \right) = \\ &= - \int_{[n, n^{8/7}]} \exp \left( -\frac{C_7(a-n)}{x} \right) d_x \mathbf{P}\{\log \rho_n > x\} = \\ &= \exp \left( -\frac{C_7(a-n)}{n} \right) \mathbf{P}\{\log \rho_n > n\} - \\ &\quad - \exp \left( -\frac{C_7(a-n)}{n^{8/7}} \right) \mathbf{P}\{\log \rho_n > n^{8/7}\} + \\ &\quad + \int_{[n, n^{8/7}]} \frac{C_7(a-n)}{x^2} \exp \left( -\frac{C_7(a-n)}{x} \right) \mathbf{P}\{\log \rho_n > x\} dx. \end{aligned}$$

Hence we obtain

$$\begin{aligned} \mathbf{P}\{\nu(n) > a\} &\leq 2 \exp \left( -\frac{C_7(a-n)}{n} \right) + \frac{C_8}{n^{1/7}} + \\ &\quad + \frac{C_7(a-n)}{n} \int_{n^{-1/7}}^1 \exp \left( -\frac{C_7(a-n)z}{n} \right) \mathbf{P}\{\rho_n > e^{n/z}\} dz, \end{aligned}$$

which, by Proposition B, yields

$$\mathbf{P}\{\nu(n) > a\} \leq 2 \exp \left( -\frac{C_7(a-n)}{n} \right) + \frac{C_8}{n^{1/7}} +$$

$$\begin{aligned}
& + \frac{C_9(a-n)}{n} \int_0^\infty \exp\left(-\frac{C_7(a-n)z}{n}\right) (1 - e^{-\pi z}) dz = \\
& = 2 \exp\left(-\frac{C_7(a-n)}{n}\right) + \frac{C_8}{n^{1/7}} + \frac{(C_9/C_7)\pi n}{C_7(a-n) + \pi n}.
\end{aligned}$$

Since  $a \geq 2n$  and  $\sup_{x>0}(xe^{-x}) < \infty$ , this yields the lemma.  $\square$

**LEMMA 3.** *If  $n \geq n_0$  is even, and if  $a$  is an integer with  $n < a \leq n^{8/7}$ , then*

$$\mathbf{P}\{e^a < \rho_n - \rho_{n/2} < e^{2a}, \rho_{n/2} < e^n, r(i) < e^a \text{ for } n+1 \leq i \leq a\} \geq \frac{C_{10} n}{a},$$

for some absolute constant  $C_{10} > 0$ .

**Proof.** By the strong Markov property,

$$\begin{aligned}
& \mathbf{P}\{e^a < \rho_n - \rho_{n/2} < e^{2a}, \rho_{n/2} < e^n, r(i) < e^a \text{ for } n+1 \leq i \leq a\} = \\
& = \mathbf{P}\{e^a < \rho_{n/2} < e^{2a}\} \mathbf{P}\{\rho_{n/2} < e^n\} (\mathbf{P}\{\rho_1 < e^a\})^{a-n}.
\end{aligned}$$

By Proposition B and Theorem B,

$$\begin{aligned}
\mathbf{P}\{e^a < \rho_{n/2} < e^{2a}\} & \geq \frac{C_{11} n}{a}, \\
\mathbf{P}\{\rho_{n/2} < e^n\} & \geq C_{12}, \\
(\mathbf{P}\{\rho_1 < e^a\})^{a-n} & \geq (\mathbf{P}\{\rho_1 < e^a\})^a \geq \left(1 - \frac{C_{13}}{a}\right)^a \geq C_{14}.
\end{aligned}$$

The last inequality holds in the case  $a > C_{13}$  but is trivially true if  $a \leq C_{13}$  (possibly with a smaller value of  $C_{14}$ ) since  $\min_{1 \leq a \leq C_{13}} (\mathbf{P}\{\rho_1 < e^a\})^a > 0$ . Assembling these pieces completes the proof of Lemma 3.  $\square$

Now, we turn to the proof of Theorem 3.

Let  $\{f_n\}$  be a positive sequence such that  $n \mapsto f_n/n$  is non-decreasing.

First, assume  $\sum_n (1/f_n) < \infty$ . Thus  $(f_n/n) \rightarrow \infty$ . Let  $n_k = 2^k$  for all  $k \geq 1$ , which implies  $\sum_k (n_k/f_{n_k}) < \infty$ . By Lemma 2, for all large  $k$ ,

$$\mathbf{P}\{\nu(n_{k+1}) > f_{n_k}\} \leq \frac{C_{15} n_{k+1}}{f_{n_k}} + \frac{C_{15}}{n_{k+1}^{1/7}}.$$

Therefore  $\sum_k \mathbf{P}\{\nu(n_{k+1}) > f_{n_k}\} < \infty$ . By the Borel–Cantelli lemma and the monotonicity of  $n \mapsto \nu(n)$ , we have

$$\limsup_{n \rightarrow \infty} \frac{\nu(n)}{f_n} \leq 1 \quad \text{a.s.}$$

Since replacing  $f_n$  by a constant multiple of  $f_n$  does not change the nature of the test, we conclude that the “lim sup” expression vanishes almost surely. This yields the first part of Theorem 3.

To prove the other part, we assume  $\sum_n(1/f_n) = \infty$ . Let again  $n_k = 2^k$ , which implies  $\sum_k(n_k/f_{n_k}) = \infty$ . It is well-known for the proof of this kind of integral test that we can assume without loss of generality that

$$n < f_n \leq n^{8/7},$$

when  $n$  is sufficiently large. (For a rigorous justification, we may use a similar argument as the one at the end of Section 3). Define for large  $k$  (say  $k \geq k_0$ ) the events

$$A_k = \{e^{f_{n_k}} < \rho_{n_k} - \rho_{n_k/2} < e^{2f_{n_k}}, \rho_{n_k/2} < e^{n_k}, \\ r(i) < e^{f_{n_k}} \text{ for all } n_k + 1 \leq i \leq f_{n_k}\}.$$

By Lemma 3, for all  $k \geq k_0$ ,

$$\mathbf{P}\{A_k\} \geq \frac{C_{10} n_k}{f_{n_k}},$$

which implies  $\sum_{k \geq k_0} \mathbf{P}\{A_k\} = \infty$ .

We now estimate  $\mathbf{P}\{A_k \cap A_\ell\}$  for  $\ell \geq k + 2$  (and  $k \geq k_0$ ). Observe that on  $A_k$ , we have

$$\rho_{f_{n_k}} < e^{2f_{n_k}} + e^{n_k} + (f_{n_k} - n_k)e^{f_{n_k}} < e^{f_{n_\ell}},$$

using the fact that  $4f_{n_k} \leq f_{n_\ell}$  (which is a consequence of the monotonicity of  $n \mapsto f_n/n$ ). On the other hand,  $\rho_{n_\ell} > e^{f_{n_\ell}}$  on  $A_\ell$ . This means that  $f_{n_k} < n_\ell$  if  $A_k \cap A_\ell \neq \emptyset$ . Write

$$\ell_0 = \ell_0(k) := \inf\{\ell \geq k : n_\ell > f_{n_k}\},$$

and we have just proved that  $A_k \cap A_\ell = \emptyset$  for  $\ell < \ell_0$ .

When  $\ell \geq \ell_0 + 1$ , we have  $n_\ell/2 > f_{n_k}$ , so that the event  $A_k$  is independent of the variable  $\rho_{n_\ell} - \rho_{n_\ell/2}$ . Accordingly, for  $\ell \geq \ell_0 + 1$ ,

$$\begin{aligned} \mathbf{P}\{A_k \cap A_\ell\} &\leq \mathbf{P}\{A_k\} \mathbf{P}\{\rho_{n_\ell} - \rho_{n_\ell/2} > e^{f_{n_\ell}}\} \leq \\ &\leq \mathbf{P}\{\rho_{n_k} - \rho_{n_k/2} > e^{f_{n_k}}\} \mathbf{P}\{\rho_{n_\ell} - \rho_{n_\ell/2} > e^{f_{n_\ell}}\} = \\ &= \mathbf{P}\{\rho_{n_k/2} > e^{f_{n_k}}\} \mathbf{P}\{\rho_{n_\ell/2} > e^{f_{n_\ell}}\}, \end{aligned}$$

which yields that for  $\ell \geq \ell_0 + 1$ ,

$$\mathbf{P}\{A_k \cap A_\ell\} \leq C_{16} \frac{n_k}{f_{n_k}} \frac{n_\ell}{f_{n_\ell}} \leq C_{17} \mathbf{P}\{A_k\} \mathbf{P}\{A_\ell\}.$$

Therefore,

$$\begin{aligned}
\sum_{k_0 \leq k \leq \ell \leq n} \mathbf{P}\{A_k \cap A_\ell\} &= \sum_{k=k_0}^n (\mathbf{P}\{A_k\} + \mathbf{P}\{A_k \cap A_{k+1}\} + \mathbf{P}\{A_k \cap A_{\ell_0}\}) + \\
&\quad + \sum_{k=k_0}^n \sum_{\ell=\ell_0+1}^n \mathbf{P}\{A_k \cap A_\ell\} \leq \\
&\leq 3 \sum_{k=k_0}^n \mathbf{P}\{A_k\} + C_{17} \sum_{k=k_0}^n \sum_{\ell=k}^n \mathbf{P}\{A_k\} \mathbf{P}\{A_\ell\}.
\end{aligned}$$

Since  $\sum_{k \geq k_0} \mathbf{P}\{A_k\} = \infty$ , this yields

$$\liminf_{n \rightarrow \infty} \frac{\sum_{k=k_0}^n \sum_{\ell=k_0}^n \mathbf{P}\{A_k \cap A_\ell\}}{\left(\sum_{k=k_0}^n \mathbf{P}\{A_k\}\right)^2} < \infty.$$

According to Kochen and Stone's version of the Borel–Cantelli lemma ([5]),  $\mathbf{P}\{A_k \text{ i.o.}\} > 0$ . Since  $A_k \subset \{\nu(n_k) > f_{n_k}\}$ , we obtain:

$$\mathbf{P}\left\{\limsup_{n \rightarrow \infty} \frac{\nu(n)}{f_n} \geq 1\right\} > 0.$$

Clearly,  $\{\limsup_{n \rightarrow \infty} (\nu(n)/f_n) \geq 1\}$  is invariant under any finite permutation between the i.i.d.r.v.'s  $\{\rho_{n+1} - \rho_n\}_{n=0}^\infty$ , the Hewitt–Savage zero-one law yields that

$$\limsup_{n \rightarrow \infty} \frac{\nu(n)}{f_n} \geq 1 \quad \text{a.s.}$$

Replacing  $f_n$  by an arbitrary multiple of  $f_n$  implies that the “lim sup” expression is infinity almost surely. This completes the proof of Theorem 3.  $\square$

## 5. Proof of Theorem 4

First we define continuous random variables which are close to the discrete variables  $r_k = \rho_k - \rho_{k-1}$ . Let  $\{\zeta_k\}_{k=1}^\infty$  be an i.i.d. sequence of random variables, uniformly distributed on  $(0, 1)$  and also independent of the sequence  $\{r_k\}_{k=1}^\infty$ . Define

$$\hat{r}_k = r_k + \zeta_k, \quad \hat{\rho}_k = \sum_{i=1}^k \hat{r}_i, \quad \hat{T}_k = \frac{\hat{r}_{k+1}}{\hat{\rho}_{k+1}}, \quad k = 1, 2, \dots$$

Then  $\{\hat{r}_k\}_{k=1}^\infty$  is a sequence of i.i.d. random variables with continuous distribution. Now let  $\hat{Z}_1 = 1$  and for  $k > 1$  put  $\hat{Z}_k = 1$  if  $\hat{r}_{k+1}$  is an extreme element, i.e.  $\hat{r}_{k+1} > \max_{1 \leq i \leq k} \hat{r}_i$  and let  $\hat{Z}_k = 0$  otherwise .

**LEMMA 4.** *As  $k \rightarrow \infty$ , we have*

$$|\widehat{Z}_k - T_k^*| = O\left(\exp\left(-\frac{k}{\log^2 k}\right)\right) \quad \text{a.s.}$$

**Proof.** We have

$$|\widehat{Z}_k - T_k^*| \leq |\widehat{Z}_k - \widehat{T}_k| + |\widehat{T}_k - T_k^*|.$$

Simple algebra shows, using  $0 \leq \zeta_k \leq 1$ , that

$$|\widehat{T}_k - T_k^*| = \left| \frac{\widehat{r}_{k+1}}{\widehat{\rho}_{k+1}} - \frac{r_{k+1}}{\rho_{k+1}} \right| \leq \frac{k+1}{\rho_{k+1}} = O\left(\exp\left(-\frac{k}{\log^2 k}\right)\right) \quad \text{a.s.},$$

the last identity following from Theorem B. This combined with Corollary 1 yields

$$\widehat{T}_k \notin \left( \exp\left(-\frac{k}{\log^2 k}\right), 1 - \exp\left(-\frac{k}{\log^2 k}\right) \right) \quad \text{a.s.},$$

for all but finitely many  $k$ . This means that for  $k$  large enough,  $\widehat{Z}_k = 1$  implies  $\widehat{T}_k \in \left(1 - \exp\left(-\frac{k}{\log^2 k}\right), 1\right)$ , while  $\widehat{Z}_k = 0$  implies  $\widehat{T}_k \in \left(0, \exp\left(-\frac{k}{\log^2 k}\right)\right)$ , so we have also

$$|\widehat{Z}_k - \widehat{T}_k| = O\left(\exp\left(-\frac{k}{\log^2 k}\right)\right),$$

proving Lemma 4. □

As a consequence, we have

$$\left| \sum_{k=1}^n \widehat{Z}_k - \sum_{k=1}^n T_k^* \right| = O(1) \quad \text{a.s.},$$

as  $k \rightarrow \infty$ . Since  $\sum_{k=1}^n \widehat{Z}_k$  is the number of extreme elements up to  $n$  from a continuous distribution, Theorem 4 follows from Theorem C. □

## 6. Proof of Theorem 5

In this section,  $\{S_n\}_{n=0}^\infty$  denotes a simple symmetric random walk on  $\mathbb{Z}^d$  ( $d \geq 3$ ), starting from 0. Let  $E_n$  be the event as in Theorem E. Then  $\mathbf{P}\{E_n \text{ i.o.}\} = 0$  if

$$(6.1) \quad f(n) \geq f^*(n) := \begin{cases} n(\log n)^{1+\varepsilon} & \text{if } d = 3 \\ n(\log_2 n)^{1+\varepsilon} & \text{if } d = 4 \\ n^{2/(d-2)}(\log n)^{2(1+\varepsilon)/(d-2)} & \text{if } d \geq 5. \end{cases}$$

Observe that if there exists an excursion of length  $g(n)$  away from  $S_n$ , i.e. if

$$S_{n+j} \neq S_n, \quad j = 1, 2, \dots, g(n), \quad S_{n+g(n)+1} = S_n,$$

then  $E_n$  does not hold with  $f(n) = g(n)$ , i.e.  $g(n)$  must be less than  $f^*(n)$  of (6.1). Let  $N := n + f^*(n) = h(n)$ , i.e.  $n = h^{-1}(N)$ . Hence

$$R(N) \leq f^*(n) = f^*(h^{-1}(N)),$$

which, in turn, easily implies that

$$\limsup_{n \rightarrow \infty} \frac{\log R(n)}{\log n} \leq \begin{cases} 1 & \text{if } d = 3, 4, \\ \frac{2}{d-2} & \text{if } d \geq 5, \end{cases}$$

almost surely.

Now we turn to the proof of the lower bound in Theorem 5. Fix  $0 < a < \min(\frac{2}{d-2}, 1)$ . Note that  $a < 1$ . It suffices to show that

$$(6.2) \quad \liminf_{n \rightarrow \infty} \frac{\log R(n)}{\log n} \geq a \quad \text{a.s.}$$

To this end, recall (see for example [8] p. 183) that  $\mathbf{P}\{S_{2n} = 0\} \sim 2(d/4\pi)^{d/2} n^{-d/2}$  ( $n \rightarrow \infty$ ). Therefore, an application of Theorem D reveals that for all  $n \geq 1$ ,

$$(6.3) \quad \frac{C_{18}}{n^{(d-2)/2}} \leq \mathbf{P}\{n \leq \rho_1 < \infty\} \leq \frac{C_{19}}{n^{(d-2)/2}},$$

where  $C_{18}$  and  $C_{19}$  are two positive constants depending only on  $d$ . Define for  $n > j \geq 0$ ,

$$\begin{aligned} \tau_j &:= \inf\{k > j : S_k = S_j\}, \\ A_j(n) &:= \{j + n^a < \tau_j \leq j + \Theta n^a\}, \end{aligned}$$

where the constant  $\Theta = \Theta(d) > 1$  is chosen to satisfy  $\Theta > (C_{19}/C_{18})^{2/(d-2)}$ .

For all  $j \leq n - \Theta n^a$ , we have

$$\begin{aligned} \mathbf{P}\{A_j(n)\} &= \mathbf{P}\{n^a < \rho_1 \leq \Theta n^a\} = \\ &= \mathbf{P}\{n^a < \rho_1 < \infty\} - \mathbf{P}\{\Theta n^a < \rho_1 < \infty\}, \end{aligned}$$

which, in view of (6.3), yields

$$\mathbf{P}\{A_j(n)\} \geq \frac{C_{18}}{n^{a(d-2)/2}} - \frac{C_{19}}{(\Theta n^a)^{(d-2)/2}} = \frac{C_{20}}{n^{a(d-2)/2}},$$

with  $C_{20} := C_{18} - C_{19}/\Theta^{(d-2)/2} \in (0, \infty)$ . On the other hand, for all  $j \leq n - \Theta n^a$ ,

$$\mathbf{P}\{A_j(n)\} \leq \mathbf{P}\{n^a < \rho_1 < \infty\} \leq \frac{C_{19}}{n^{a(d-2)/2}},$$

the last inequality following from (6.3). As a consequence,

$$(6.4) \quad C_{21} n^{1-a(d-2)/2} \leq \mathbf{E} \left\{ \sum_{j=1}^{n-\Theta n^a} \mathbf{1}_{A_j(n)} \right\} \leq C_{19} n^{1-a(d-2)/2}.$$



Note that  $1 - a(d - 2)/2 > 0$ .

Assume for the moment that we can show that

$$(6.5) \quad \text{Var} \left( \sum_{j=1}^{n-\Theta n^a} \mathbf{1}_{A_j(n)} \right) = O \left( n^{2-a(d-2)-\delta} \right),$$

for some  $\delta > 0$ . Then by Chebyshev's inequality and (6.4),

$$\begin{aligned} \mathbf{P} \left\{ \left| \sum_{j=1}^{n-\Theta n^a} \mathbf{1}_{A_j(n)} - \mathbf{E} \left( \sum_{j=1}^{n-\Theta n^a} \mathbf{1}_{A_j(n)} \right) \right| > \frac{1}{2} \mathbf{E} \left( \sum_{j=1}^{n-\Theta n^a} \mathbf{1}_{A_j(n)} \right) \right\} &\leq \\ &\leq \frac{4 \text{Var} \left( \sum_{j=1}^{n-\Theta n^a} \mathbf{1}_{A_j(n)} \right)}{\left( \mathbf{E} \left( \sum_{j=1}^{n-\Theta n^a} \mathbf{1}_{A_j(n)} \right) \right)^2} \leq \frac{C_{22}}{n^\delta}, \end{aligned}$$

which, by the Borel–Cantelli lemma, implies that almost surely for all large  $k$ ,

$$\sum_{j=1}^{n_k - \Theta n_k^a} \mathbf{1}_{A_j(n_k)} \geq \frac{1}{2} \mathbf{E} \left( \sum_{j=1}^{n_k - \Theta n_k^a} \mathbf{1}_{A_j(n_k)} \right),$$

where  $n_k := \lfloor k^{2/\delta} \rfloor$ . By (6.4),  $\mathbf{E} \left( \sum_{j=1}^{n_k - \Theta n_k^a} \mathbf{1}_{A_j(n_k)} \right) \geq 2$ , so that for large  $k$ , there exists  $j \in [1, n_k - \Theta n_k^a]$  such that  $\mathbf{1}_{A_j(n_k)} = 1$ . This yields  $R(n_k) \geq n_k^a$ , and will complete the proof of (6.2), using the monotonicity of  $n \mapsto R(n)$ .

It remains to check (6.5). We have

$$\begin{aligned} \mathbf{E} \left[ \left( \sum_{j=1}^{n-\Theta n^a} \mathbf{1}_{A_j(n)} \right)^2 \right] &= \mathbf{E} \left( \sum_{j=1}^{n-\Theta n^a} \mathbf{1}_{A_j(n)} \right) + 2 \sum_{1 \leq i < j \leq n-\Theta n^a} \mathbf{P}\{A_i(n)A_j(n)\} = \\ &= \mathbf{E} \left( \sum_{j=1}^{n-\Theta n^a} \mathbf{1}_{A_j(n)} \right) + \\ &\quad + 2 \sum_{1 \leq i < j \leq n-\Theta n^a} \mathbf{P}\{A_i(n), A_j(n), \tau_i \leq j\} + \\ &\quad + 2 \sum_{1 \leq i < j \leq n-\Theta n^a} \mathbf{P}\{A_i(n), A_j(n), \tau_i > j\}. \end{aligned}$$

By the Markov property,

$$\mathbf{P}\{A_i(n), A_j(n), \tau_i \leq j\} \leq \mathbf{P}\{A_i(n)\} \mathbf{P}\{A_j(n)\}.$$

On the other hand, if  $A_i(n) \cap \{\tau_i > j\} \neq \emptyset$ , then  $j < \tau_i \leq i + \Theta n^a$ , which implies  $j - i \leq \Theta n^a$ . The proof of (6.5) is thus reduced to showing the following: for some  $\delta > 0$ ,

$$(6.6) \quad \sum_{(i,j) \in \Omega(n)} \mathbf{P}\{A_i(n), A_j(n), \tau_i > j\} = O \left( \frac{n^{2-a(d-2)}}{n^\delta} \right),$$

where  $\Omega(n) := \{(i, j) : 1 \leq i < j \leq n - \Theta n^a, j - i \leq \Theta n^a\}$ .

The proof of (6.6) will be carried out in a few steps, namely, we will show that

$$(6.7) \quad \sum_{(i,j) \in \Omega(n)} \mathbf{P} \left\{ A_i(n), A_j(n), \tau_j > \tau_i > j + \frac{n^a}{2} \right\} = O \left( n^{1-a(d-2)/2} b_n \right),$$

$$(6.8) \quad \sum_{(i,j) \in \Omega(n)} \mathbf{P} \left\{ A_i(n), A_j(n), j < \tau_i \leq j + \frac{n^a}{2} \right\} = O \left( \frac{n^{2-a(d-2)/2}}{n^{1-a}} \right),$$

$$(6.9) \quad \sum_{(i,j) \in \Omega(n)} \mathbf{P} \{ A_i(n), A_j(n), \tau_j \leq \tau_i \} = O \left( n^{1-a(d-2)/2} b_n \right),$$

where  $b_n := \sum_{k=1}^{\Theta n^a} k^{-(d-2)/2}$ . Since  $b_n = O(n^{1-a(d-2)/2-\delta})$  for some  $\delta > 0$  (recalling that  $a < \min(\frac{2}{d-2}, 1)$ ), the estimates (6.7)–(6.9) together suffice to yield (6.6), hence the lower bound in Theorem 5.

To check (6.7), observe that

$$\begin{aligned} I_1 &:= \mathbf{P} \left\{ A_i(n), A_j(n), \tau_j > \tau_i > j + \frac{n^a}{2} \right\} \\ &\leq \mathbf{P} \{ \tau_i \in (j + n^a/2, i + \Theta n^a], \tau_j > \tau_i \}. \end{aligned}$$

Applying the strong Markov property at  $\tau_i$  gives that

$$I_1 \leq \mathbf{E} \left( \mathbf{1}_{\{\tau_i \in (j+n^a/2, i+\Theta n^a)\}} p_1(S_i, S_j) \right),$$

where  $p_1(y, x) := \mathbf{P}^y \{ \rho_1(x) < \infty \}$  for  $x \in \mathbb{Z}^d$  and  $y \in \mathbb{Z}^d$ , and  $\mathbf{P}^y$  is the probability under which the random walk starts from  $y$ . According to Erdős and Taylor [2], there exists a constant  $C_{23} = C_{23}(d)$  such that for all  $(x, y) \in \mathbb{Z}^d \times \mathbb{Z}^d$ ,

$$(6.10) \quad \mathbf{P}^y \{ \rho_1(x) < \infty \} \leq \frac{C_{23}}{(\|x - y\| + 1)^{d-2}}.$$

Therefore

$$I_1 \leq C_{23} \mathbf{E} \left[ \frac{\mathbf{1}_{\{\tau_i \in (j+n^a/2, i+\Theta n^a)\}}}{(\|S_j - S_i\| + 1)^{d-2}} \right] = C_{23} \mathbf{E} \left[ \frac{p_2(S_j, S_i)}{(\|S_j - S_i\| + 1)^{d-2}} \right],$$

where  $p_2(y, x) := \mathbf{P}^y \{ n^a/2 < \rho_1(x) \leq i + \Theta n^a - j \}$  (and we have applied the Markov property at  $j$ ).

We now estimate  $p_2(y, x)$ . According to [8] p. 184,  $\sup_{x \in \mathbb{Z}^d} \mathbf{P} \{ S_n = x \} \leq C_{24}/n^{d/2}$ , so that for  $m < n$ ,

$$\mathbf{P}^y \{ m \leq \rho_1(x) < n \} \leq \sum_{k=m}^{n-1} \mathbf{P}^y \{ S_k = x \} \leq \sum_{k=m}^{n-1} \frac{C_{24}}{k^{d/2}} \leq C_{24} \frac{n-m}{m^{d/2}}.$$

Therefore, for all  $m < n$ ,

$$(6.11) \quad \sup_{x \in \mathbb{Z}^d} \sup_{y \in \mathbb{Z}^d} \mathbf{P}^y \{m \leq \rho_1(x) < n\} \leq C_{24} \frac{n}{m^{d/2}}.$$

In particular,  $p_2(y, x) \leq C_{24} \Theta n^a / (n^a/2)^{d/2}$  for all  $x$  and  $y$ , so that

$$I_1 \leq \frac{C_{25}}{n^{a(d-2)/2}} \mathbf{E} \left[ \frac{1}{(\|S_j - S_i\| + 1)^{d-2}} \right].$$

Recall Lemma 16.5 of [8] p. 184:  $\mathbf{P}\{S_n = x\} \leq C_{26} n^{-d/2} \exp(-\|x\|^2/2n)$  for all  $x \in \mathbb{Z}^d$  and  $n \geq 1$ , from which it follows that

$$(6.12) \quad \mathbf{E} \left( (\|S_n\| + 1)^{-(d-2)} \right) \leq C_{27} n^{-(d-2)/2}, \quad n \geq 1.$$

Accordingly,

$$I_1 \leq \frac{C_{25}}{n^{a(d-2)/2}} \frac{C_{27}}{(j-i)^{(d-2)/2}}.$$

This implies (6.7).

Now let us check (6.8). Clearly,

$$\begin{aligned} I_2 &:= \mathbf{P} \left\{ A_i(n), A_j(n), j < \tau_i \leq j + \frac{n^a}{2} \right\} \leq \\ &\leq \mathbf{P} \left\{ \tau_i \in (i + n^a, j + n^a/2], j < \tau_i, \tau_j \in (j + n^a, j + \Theta n^a) \right\}. \end{aligned}$$

Applying the strong Markov property at  $\tau_i$  gives

$$I_2 \leq \mathbf{E} \left( \mathbf{1}_{\{\tau_i \in (i+n^a, j+n^a/2], j < \tau_i\}} p_3(S_i, S_j, \tau_i) \right),$$

where  $p_3(y, x, \ell) := \mathbf{P}^y \{ \rho_1(x) \in (j + n^a - \ell, j + \Theta n^a - \ell] \}$ . By (6.11), on the event  $\{ \tau_i \in (i + n^a, j + n^a/2] \}$ , we have  $p_3(S_i, S_j, \tau_i) \leq C_{24} \Theta n^a / (n^a/2)^{d/2}$ , so that

$$\begin{aligned} I_2 &\leq \frac{C_{28}}{n^{a(d-2)/2}} \mathbf{P} \{ \tau_i \in (i + n^a, j + n^a/2] \} = \\ &= \frac{C_{28}}{n^{a(d-2)/2}} \mathbf{P} \{ n^a < \rho_1 \leq j + n^a/2 - i \} \leq \\ &\leq \frac{C_{28}}{n^{a(d-2)/2}} \frac{C_{29}}{n^{a(d-2)/2}}, \end{aligned}$$

the last inequality following from (6.11) (recalling that  $j - i \leq \Theta n^a$ ). Therefore,

$$\sum_{(i,j) \in \Omega(n)} I_2 \leq n \Theta n^a \frac{C_{28} C_{29}}{n^{a(d-2)}} = \Theta C_{28} C_{29} \frac{n^{2-a(d-2)}}{n^{1-a}},$$

proving (6.8).

Finally, to verify (6.9), we observe that

$$\begin{aligned} I_3 &:= \mathbf{P} \{A_i(n), A_j(n), \tau_j \leq \tau_i\} \leq \\ &\leq \mathbf{P} \{\tau_j \in (j + n^a, j + \Theta n^a], \tau_i \geq \tau_j\} = \\ &= \mathbf{E} \left( \mathbf{1}_{\{\tau_j \in (j+n^a, j+\Theta n^a]\}} p_1(S_j, S_i) \right), \end{aligned}$$

where  $p_1(y, x) = \mathbf{P}^y \{\rho_1(x) < \infty\}$  as before, and we have applied in the last equality the strong Markov property at  $\tau_j$ . By (6.10),  $p_1(S_j, S_i) \leq C_{23}/(\|S_j - S_i\| + 1)^{d-2}$ , so that

$$I_3 \leq C_{23} \mathbf{E} \left[ \frac{\mathbf{1}_{\{\tau_j \in (j+n^a, j+\Theta n^a]\}}}{(\|S_j - S_i\| + 1)^{d-2}} \right].$$

Using the Markov property at  $j$ , we get

$$I_3 \leq C_{23} \mathbf{E} \left[ \frac{p_4(S_j, S_j)}{(\|S_j - S_i\| + 1)^{d-2}} \right],$$

where  $p_4(y, y) := \mathbf{P}^y \{\rho_1(y) \in (n^a, \Theta n^a]\} = \mathbf{P} \{n^a < \rho_1 \leq \Theta n^a\} \leq C_{24} \Theta n^{-a(d-2)/2}$  by means of (6.11). Combined with (6.12), it follows that

$$I_3 \leq \frac{C_{23} C_{24} \Theta C_{27}}{(j-i)^{(d-2)/2} n^{a(d-2)/2}}.$$

This yields (6.9), and completes the proof of the lower bound in Theorem 5.  $\square$

## Acknowledgements

Cooperation between the authors was supported by the joint French–Hungarian Inter-governmental Grant “Balaton” (grant no. F25/97).

## References

- [1] CSÁKI, E. - RÉVÉSZ, P. - ROSEN, J. (1998) Functional laws of the iterated logarithm for local times of recurrent random walks on  $\mathbb{Z}^2$ . *Ann. Inst. Henri Poincaré, Probabilités et Statistiques* **34**, 545-563.
- [2] ERDŐS, P. - TAYLOR, S.J. (1960) Some problems concerning the structure of random walks paths. *Acta Math. Acad. Sci. Hung.* **11**, 137-162.
- [3] ERDŐS, P. - TAYLOR, S.J. (1960) Some intersection properties of random walk paths. *Acta Math. Acad. Sci. Hung.* **11**, 231-248.
- [4] GRIFFIN, P. (1990) Accelerating beyond the third dimension: returning to the origin in simple random walk. *Math. Sci.* **15**, 24-35.
- [5] KOCHEN, S.B. - STONE, C.J. (1964) A note on the Borel-Cantelli lemma. *Illinois J. Math.* **8**, 248-251.
- [6] RÉNYI, A. (1962) On the extreme elements of observations. *MTA III. Oszt. Közl.* **12**, 105-121 (in Hungarian). English translation: *Selected Papers of Alfréd Rényi* Vol. 3, pp. 50-65. Publishing House of the Hungarian Academy of Sciences, Budapest, 1976.
- [7] RÉNYI, A. (1962) Théorie des éléments saillants d'une suite d'observations. *Annales Fac. Sci. Univ. Clermont-Ferrand* **2**, No. 8, 7-12.
- [8] RÉVÉSZ, P. (1990) *Random Walk in Random and Non-Random Environments*, World Scientific, Singapore, 1990.
- [9] RÉVÉSZ, P. - WILLEKENS, E. (1988) On the maximal distance between two renewal epochs. *Stochastic Process. Appl.* **27**, 21-41.

Endre Csáki  
Alfréd Rényi Institute of Mathematics  
Hungarian Academy of Sciences  
P.O.B. 127  
H-1364 Budapest  
Hungary  
`csaki@math-inst.hu`

Pál Révész  
Institut für Statistik und Wahrscheinlichkeitstheorie  
Technische Universität Wien  
Wiedner Hauptstrasse 8-10/107  
A-1040 Vienna  
Austria  
`revesz@ci.tuwien.ac.at`

Zhan Shi  
Laboratoire de Probabilités UMR 7599  
Université Paris VI  
4 Place Jussieu  
F-75252 Paris Cedex 05  
France  
`zhan@proba.jussieu.fr`