

## Some remarks on set theory. VI.

By P. ERDŐS in Budapest and G. FODOR in Szeged,

Let  $E$  be a given non countable set of power  $\aleph_0$  and suppose that there exists a relation  $R$  between the elements of  $E$ . For any  $x \in E$ , let  $R(x)$  denote the set of the elements  $y \in E$  for which  $xRy$  holds. Two distinct elements of  $E$ ,  $x$  and  $y$ , are called *independent*, if  $x \notin R(y)$  and  $y \notin R(x)$ . A subset  $F$  of  $E$  is called free if  $F$  has only one element or if  $F$  has more elements and any two of them being independent. Let  $B$  be a system of subsets of  $E$ ; then a non empty system  $I \subset B$  is called a p-additive ideal, if  $\bigcup I$  is again a set of  $I$ , and if  $X \in I$ ,  $Y \in B$ ,  $Y \subset X$  imply  $Y \in I$ .

We assume that  $\{x\} \in B$  and  $\{x\} \in I$  for every  $x \in E$ , and one of the following conditions holds for the sets  $R(x)$ :

- (A) There is a cardinal number  $n < \aleph_0$  such that, for every  $x \in E$ ,  $|R(x)| < n$ ,
- (B)  $E$  is a metric space and  $d(x, R(x)) > 0$ , where  $d(x, R(x))$  denotes the distance of the point  $x$  from the set  $R(x)$ .

We deal in this paper first with the following question :

(i) *If  $A$  is a system of sets of  $B-I$ , does there exist a free subset  $E'$  of  $E$  such that for every  $X \in A$ ,  $X \cap E' \in B-I$ ?*

This question has been studied previously in the following special cases :

a)  $\aleph_0$  is regular, condition (A) holds,  $B$  is the set of all subsets of  $E$ ,  $I$  is the set of all subsets of  $E$ , of power less than  $\aleph_0$ , and  $A = 1$  {then  $\mathfrak{p} = \aleph_0$ }.

(See [1].)

b)  $E = [0,1]$  with the ordinary metric, condition (B) holds,  $B$  is the set of all subsets of  $E$ ,  $I$  is the set of all subsets of measure zero in the Lebesgue sense, and  $\bar{A} = 1$ .

(The answer to this question is affirmative, see [2].)

c) The same hypotheses as in b), with the only difference that  $B$  is the set of all subsets of  $[0,1]$  measurable in the Lebesgue sense.

(The answer to this question is generally in the negative. The answer is affirmative if  $g(x) = d(x, R(x))$  is a measurable function in the Lebesgue sense, see [3], [4].)

d)  $E = [0,1]$  with the ordinary metric  $d$   $B$  is a Boolean  $\sigma$ -algebra of subsets of  $[0,1]$  containing all subintervals of  $[0,1]$ , and  $I$  is the set of the sets  $X$  of  $B$  such that  $\mu(X) = 0$ , where  $\mu$  is a measure on  $B$ .<sup>1)</sup>

(If  $\mu$  is not identically zero and if there exists a function  $f$  measurable with respect to  $B$  and such that  $0 < f(x) \leq g(x) = d(x, R(x))$  for all  $x \in [0,1]$ , then there exists a free set  $F$  in  $B$  such that  $\mu(F) > 0$  (i.e.  $F \notin I$ ). This theorem is due to P. HALMOS.<sup>2)</sup>)

In section 1 first we prove making use of a method of ULAM [6] the following theorem (Theorem 1): If  $E$  is a set of power  $\aleph_\beta$  with  $\aleph_\beta$  greater than  $\aleph_0$  and less than the first aleph inaccessible in the weak sense,  $I$  is a proper  $\aleph_{\beta+1}$ -additive ideal of subsets of  $E$  such that  $\{x\} \in I$  for every  $x \in E$  and  $F \notin I$ , then  $F$  may be decomposed into the sum of a sequence of the type  $\omega_{\beta+1}$  of mutually disjoint subsets  $F_\xi$  of  $E$ , such that  $F_\xi \notin I$ .

We use this theorem in the proof of theorem 3.

In sections I and II a number of results is given with respect to question (i). For instance we shall prove that the answer to the problem is affirmative in the following cases:

1) If  $m > \aleph_0$  is less than the first aleph inaccessible in the weak sense,  $B$  is the set of all subsets of  $E$ ,  $I$  is a  $\aleph_{\gamma+1}$  additive ideal ( $\aleph_{\gamma+1} \leq m$ ),  $A = \aleph_0$  and  $f?(x) < \aleph_0$  for every  $x \in E$ .

2) If  $E$  is a metric space which contains a dense subset, the power of which is less than the first aleph inaccessible in the weak sense,  $B$  is the set of all Borel sets of  $E$ ,  $I$  is the  $\sigma$ -ideal of all sets of  $\mu$ -measure zero of  $B$ , where  $\mu$  is a measure on  $B$ ,  $A = 1$ , the condition (B) is satisfied, and also the following condition (C) holds:

(C) there is a real number  $i > 0$  such that the set  $\{x : g(x) \geq i\}$  contains in  $B$  a subset of positive measure, where  $g(x) = d(x, R(x))$ .

If, for every  $x \in E$ , the set  $Z?(x)$  is the complement of a sphere of  $E$  whose center is at  $x$ , then the condition (C) is not only sufficient, but also necessary for the existence of a free subset of  $E$  in  $B$ .

Finally, in the section III, we deal with the following question :

(ii) *Lef  $K$  be a class of subsets of  $E$ . When does there exist a relation*

<sup>1)</sup> We use the terminology of P. R. HALMOS [11].

<sup>2)</sup> See his review of the paper [3] in *Math. Reviews*, 12 (1951), p. 398.

$R$  for which the condition (A) holds and there is no free subset  $X \in K$  with respect to  $R$ ?

For instance we shall prove that if  $\bar{K} = \aleph_0$  and every element of  $K$  is of power  $\aleph_0$ , then there exists a relation  $R$ , with  $R(x) \leq 1$  for every  $x \in E$ , for which there is no free set in  $K$ .

This result shows that the answer to the problem (i) is always negative if  $\bar{\mathbf{B}} = \aleph_0$  and every element of  $\mathbf{B}$  is of power  $\aleph_0$ .

**Notation and definitions.** Throughout this paper, the symbols  $\bar{F}$  and  $\bar{\beta}$  denote the cardinal number of the set  $F$  and of the ordinal number  $\beta$  respectively. For any  $x \in E$ , let  $R^{-1}(x) = \{y : x \in R(y)\}$ . For any subset  $F$  of  $E$  let

$$R[F] = \bigcup_{x \in F} R(x) \quad \text{and} \quad R^{-1}[F] = \bigcup_{x \in F} R^{-1}(x).$$

For any cardinal number  $\kappa$  we denote by  $\varphi_\kappa$  the initial number of  $\kappa$ , by  $\kappa^+$  the smallest cardinal number for which  $\kappa$  is the sum of  $\kappa^+$  cardinal numbers each of which is smaller than  $\kappa$ , by  $\kappa^+$  the cardinal number immediately following  $\kappa$ . We say that  $\kappa$  is regular if  $\kappa^+ = \kappa$  and singular if  $\kappa^+ < \kappa$ ;  $\kappa \Rightarrow \aleph_\kappa \geq \aleph_0$  is called inaccessible in the weak sense, if  $\gamma$  is a limit number and  $\kappa$  is regular,

## I.

We assume in this section that the sets  $R(x)$  satisfy condition (A) and  $\mathbf{B}$  is the set of all subsets of  $E$ . We shall use the following

**L e m m a.** Let  $T$  be a set of power  $\aleph_{\alpha+1}$  (where  $\alpha$  is a given ordinal number  $\geq 0$ ). There exists a system  $\{A_\eta^\xi\}_{\xi < \omega_{\alpha+1}}$  of subsets of  $T$  such that

- 1)  $T = \bigcup_{\eta < \omega_{\alpha+1}} A_\eta^\xi$  for every  $\xi < \omega_{\alpha+1}$
- 2)  $A_\eta^\xi \cap A_\zeta^\xi = \emptyset$  for  $\xi < \omega_\alpha$  and  $\eta < \zeta < \omega_{\alpha+1}$ ,
- 3) the power of the set  $T - \bigcup_{\xi < \omega_\alpha} A_\eta^\xi$  is  $\leq \aleph_\alpha$  for every  $\eta < \omega_{\alpha+1}$ . (See S. ULAM [6] p. 143.)

We prove now the following

**T h e o r e m 1.** Let  $E$  be a set of power  $\aleph_\lambda$  with  $\aleph_\lambda$  greater than  $\aleph_0$  and less than the first aleph inaccessible in the weak sense, and let  $\mathbf{I}$  be a proper  $\aleph_{\lambda+1}$ -additive ideal of subsets of  $E$  such that  $\{x\} \in \mathbf{I}$  for every  $x \in E$ . If  $B \subseteq E$  and  $B \notin \mathbf{I}$ , then there exists a sequence  $\{B_\xi\}_{\xi < \omega_{\lambda+1}}$  of type  $\omega_{\lambda+1}$ , of subsets of  $E$ , such that

- (i)  $B_\xi \notin \mathbf{I}$  for every  $\xi < \omega_{\lambda+1}$ ,
- (ii)  $B_\xi \cap B_\eta = \emptyset$  for  $\xi < \eta < \omega_{\lambda+1}$ ,
- (iii)  $|B| = \bigcup_{\xi < \omega_{\lambda+1}} B_\xi$ .

P r o of"). We use transfinite induction. First we prove that our theorem is true for  $\gamma = \lambda + 1$ . Let  $\bar{E} = \aleph_{\lambda+1}$  and  $B \notin I$ . It is obvious that  $\bar{B} = \aleph_{\lambda+1}$ . By the lemma ( $a = \lambda$  and  $T = B$ ) there is a system  $\{A_{\eta}^{\xi} | \xi < \omega_{\lambda}\}_{\eta < \omega_{\lambda+1}}$  of subsets of  $B$  for which 1), 2) and 3) hold. Since  $B \notin I$  and, by 3)  $B - \bigcup_{\xi < \omega_{\lambda}} A_{\eta}^{\xi} \in I$  for every  $\eta < \omega_{\lambda+1}$ , there exists for every  $\eta < \omega_{\lambda+1}$  an ordinal number  $\xi(\eta) < \omega_{\lambda}$  such that  $A_{\eta}^{\xi(\eta)} \notin I$ . It follows that there is an ordinal number  $\xi_0 < \omega_{\lambda}$  and a sequence  $\{\eta_v\}_{v < \omega_{\lambda+1}}$  of type  $\omega_{\lambda+1}$ , of the ordinal numbers  $q < \omega_{\lambda+1}$ , such that  $\xi(\eta_v) = \xi_0$  and  $A_{\eta_v}^{\xi_0} \notin I$  for every  $v < \omega_{\lambda+1}$ . Let  $A = \{\eta : \eta < \omega_{\lambda+1} \text{ and } \eta \neq \eta_v \text{ if } v < \omega_{\lambda+1}\}$  and

$$B_v = \begin{cases} A_{\eta_0}^{\xi_0} \cup (\bigcup_{\eta \in A} A_{\eta}^{\xi_0}) & \text{for } v = 0, \\ A_{\eta_v}^{\xi_0} & \text{for } 0 < v < \omega_{\lambda+1}. \end{cases}$$

Obviously the set  $\{B_v\}_{v < \omega_{\lambda+1}}$  satisfies the conditions (i), (ii) and (iii).

Let now  $\beta$  be a given ordinal number,  $\beta > \lambda + 1$ , such that  $\aleph_{\beta}$  is less than the first aleph inaccessible in the weak sense, and suppose that the theorem is true for every  $\alpha < \beta$ . Let  $\bar{E} = \aleph_{\beta}$  and  $B \notin I$  (BEE).

If  $B < \aleph_{\beta}$ , then the theorem is true by the induction hypothesis. (Let  $I \in I$ , if and only if  $I_1 = B \cap I$ , where  $I \in I$ . Obviously  $I$  is an  $\aleph_{\lambda+1}$ -additive ideal in  $B$ .)

If  $\bar{B} = \aleph_{\beta}$ , then there are two possibilities :

- a)  $\beta$  is an ordinal number of the first kind, i. e.  $\beta = \alpha + 1$ ,
- b)  $\beta$  is an ordinal number of the second kind.

*Case a).* By the lemma ( $\beta = \alpha + 1$  and  $T = B$ ) there is a system  $\{A_{\eta}^{\xi} | \xi < \omega_{\alpha}, \eta < \omega_{\alpha+1}\}$  of subsets of  $B$  for which 1), 2) and 3) hold.

We have two subcases :

a,) if  $B = \bigcup_{\zeta < \omega_{\alpha}} C_{\zeta}$  is an arbitrary decomposition of  $B$  into the sum of  $\aleph_{\alpha}$  subsets, then there is an ordinal number  $\zeta_0 < \omega_{\alpha}$  such that  $C_{\zeta_0} \notin I$

a,)  $B$  has a decomposition  $B = \bigcup_{\zeta < \omega_{\alpha}} C_{\zeta}$  into the sum of  $\aleph_{\alpha}$  subsets such that, for every  $\zeta < \omega_{\alpha}$ ,  $C_{\zeta} \in I$ .

*Subcase a.).* For every  $\eta < \omega_{\alpha+1}$  there is an ordinal number  $\xi(\eta) < \omega_{\alpha}$  such that  $A_{\eta}^{\xi(\eta)} \notin I$ . It follows that there is an ordinal number  $\xi_0 < \omega_{\alpha}$  and a sequence  $\{\eta_v\}_{v < \omega_{\alpha+1}}$  of type  $\omega_{\alpha+1}$ , of ordinal numbers  $q < \omega_{\alpha+1}$ , such that  $\xi(\eta_v) = \xi_0$  and  $A_{\eta_v}^{\xi_0} \notin I$  for every  $v < \omega_{\alpha+1}$ . Let  $A = \{\eta : \eta < \omega_{\alpha+1} \text{ and } \eta \neq \eta_v \text{ if } v < \omega_{\alpha+1}\}$

---

<sup>3)</sup> We make use of a method of ULAM [6].

if  $\eta \triangleleft \omega_{\lambda+1}$ , and

$$B_\nu = \begin{cases} A_{\eta_0}^{\xi_0} \cup (\bigcup_{\eta \in A} A_{\eta}^{\xi_0}) & \text{for } \nu = 0, \\ A_{\eta_\nu}^{\xi_0} & \text{for } 0 < \nu < \omega_{\lambda+1}. \end{cases}$$

*Subcase a).* Let  $B = \bigcup_{\zeta \triangleleft \omega_\alpha} C_\zeta$  be a decomposition of  $B$  into the sum of  $\aleph_\alpha$  subsets such that  $C_{\zeta_1} \cap C_{\zeta_2} = 0$  for  $\zeta_1 \triangleleft \zeta_2 \triangleleft \omega_\alpha$  and  $C_\zeta \notin I$  for every  $\zeta \triangleleft \omega_\alpha$ . Consider the set  $D = \{C_\zeta\}_{\zeta \triangleleft \omega_\alpha}$ . We define an  $\aleph_{\lambda+1}$ -additive ideal  $I$  in  $D$  as follows: Let  $F \in I'$  if and only if  $F \subset D$  and  $\bigcup_{C \in F} C \notin I$ . Since  $\bar{D} = \aleph_\alpha \triangleleft \aleph_\beta$  and  $D \notin I'$ , there is, by the induction hypothesis, a decomposition

$$D = \bigcup_{\eta \triangleleft \omega_{\lambda+1}} F_\eta$$

of  $D$  into the sum of  $\aleph_{\lambda+1}$  subsets such that  $F_{\eta_1} \cap F_{\eta_2} = 0$  if  $\eta_1 \neq \eta_2$  and  $F_\eta \notin I$  for every  $\eta \triangleleft \omega_{\lambda+1}$ . Let

$$B_\eta = \bigcup_{C \in F_\eta} C$$

Obviously  $B_{\eta_1} \cap B_{\eta_2} = 0$  if  $\eta_1 \neq \eta_2$ ,  $B_\eta \notin I$  for every  $\eta \triangleleft \omega_{\lambda+1}$ , and

$$B = \bigcup_{\eta \triangleleft \omega_{\lambda+1}} B_\eta.$$

*Case b).* Since  $\aleph_\beta$  is less than the first aleph inaccessible in the weak sense,  $B$  has a decomposition  $B = \bigcup_{\zeta \triangleleft \omega_\beta} C_\zeta$  into the sum of  $\aleph_\beta \triangleleft \aleph_\beta$  subsets such that  $\aleph_\beta \triangleleft \bar{C}_\zeta \triangleleft \aleph_\beta$  and  $C_{\zeta_1} \cap C_{\zeta_2} = 0$  if  $\zeta_1 \neq \zeta_2$ .

If there is an ordinal number  $\xi_0 < \omega_\beta$  for which  $C_{\xi_0} \notin I$ , then there is, by the induction hypothesis, a decomposition

$$C_{\xi_0} = \bigcup_{\eta \triangleleft \omega_{\lambda+1}} D_\eta$$

of  $C_{\xi_0}$  such that  $D_{\xi_1} \cap D_{\xi_2} = 0$  for  $\xi_1 \neq \xi_2$  and  $D_\xi \notin I$  for every  $\xi \triangleleft \omega_{\lambda+1}$ . Let

$$B_\zeta = \begin{cases} D_\zeta \cup (\bigcup_{\xi \triangleleft \xi_0} C_\xi) & \text{for } \zeta = 0, \\ D_\zeta & \text{for } 0 < \zeta < \omega_{\lambda+1}. \end{cases}$$

Obviously the set  $\{B_\zeta\}_{\zeta \triangleleft \omega_{\lambda+1}}$  satisfies the conditions (i), (ii), and (iii).

The proof of the case, when  $C_\zeta \notin I$  for every  $\zeta < 10$ , is similar to that of case a.). Theorem 1 is proved.

*Corollary 1.* If  $I$  is a measure  $\aleph_\alpha$  is less than the first aleph inaccessible in the weak sense, then every finite measure  $\mu$ ,<sup>4)</sup> defined for all subsets of  $E$  and vanishing for all one-point sets, vanishes identically. (See S. ULAM [6].)

<sup>4)</sup> We call a measure every extended real valued, non negative, countably additive set function  $p(X)$  defined in a ring of subsets of  $E$ . A ring of sets is a non empty class  $R$  of sets such that if  $E \in R$  and  $F \in R$ , then  $E \cup F \in R$  and  $E - F \in R$ .

**P r o of.** The set of all subsets  $F$  of  $E$  for which  $\mu(F) = 0$  is an &additive ideal  $I$  containing all one-point subsets of  $E$ . If  $\mu$  is not identically zero, then there exists a subset  $F$  of  $E$  such that  $\mu(F) \neq 0$ ; i. e.  $I$  is-a proper ideal. By Theorem 1 there exists a sequence  $\{F_\xi\}_{\xi < \omega_1}$  of type  $\omega_1$ , of subsets of  $E$ , satisfying the conditions (i), (ii), (iii). Let  $H_n$  be the set of the ordinal numbers  $\xi < \omega_1$  for which  $\mu(F_\xi) > \frac{1}{n}$  ( $n = 1, 2, \dots$ ). It follows that there is a natural number  $n_0$  such that  $H_{n_0} = \aleph_0$ . Let  $\{i_n\}_{n < \omega}$  be an enumeration of  $H_{n_0}$ . By the  $\sigma$ -additivity of  $\mu$  we have

$$\mu(\bigcup_{n=1}^{\infty} F_{i_n}) = \sum_{n=1}^{\infty} \mu(F_{i_n}) \geq \frac{1}{n_0} + \frac{1}{n_0} + \dots + \frac{1}{n_0} + \dots = \infty,$$

which is impossible since  $\mu$  is finite.

**C o r o l l a r y 2.** *If  $2^\omega$  is less than the first aleph inaccessible in the weak sense, then for every subset  $F$  of the second category of the set of real numbers  $E$  there is a sequence  $\{F_\xi\}_{\xi < \omega_1}$  of type  $\omega_1$ , of mutually disjoint subsets of  $E$  of the second category, such that*

$$F = \bigcup_{\xi < \omega_1} F_\xi$$

**P roof.** The set  $I$  of all subsets of the first category of  $E$  is a n-ideal (i. e. an &additive ideal). (See W. SIERPIŃSKI [8] p. 176.)

**C o r o l l a r y 3.** *If  $2^\omega$  is less than the first aleph inaccessible in the weak sense and  $\mu^*(F)$  is an outer measure") not identically zero on the set of all subsets of the set  $E$  of real numbers such that  $\mu^*(\{x\}) = 0$  for every  $x \notin E$ , then for every subset  $F$  of  $E$  for which  $\mu^*(F) \neq 0$ , there is a sequence  $\{F_\xi\}_{\xi < \omega_1}$  of the type  $\omega_1$ , of mutually disjoint subsets  $F_\xi$  of  $E$  such that  $\mu^*(F_\xi) \neq 0$  and*

$$F = \bigcup_{\xi < \omega_1} F_\xi$$

**P roof.** The set  $I$  of all subsets  $F$  of  $E$  for which  $\mu^*(F) = 0$  is a n-ideal. (See W. SIERPIŃSKI [8] p. 109, Proposition C<sub>34</sub>.)

**T h e o r e m 2.** *Let  $\bar{E} = \aleph_0 > \aleph_0$  and suppose that there exists a relation  $R$  between the elements of  $E$ , such that for any  $x \in E$ , the power of the set  $R(x) = \{y: xRy\}$  is smaller than  $\aleph_0$ . Let furthermore  $I$  be an  $\aleph_0$ -additive proper ideal of  $E$ , such that  $\{x\} \notin I$  for any  $x \in E$ . Then there exists a free subset  $E'$  of  $E$ , such that  $E' \notin I$ ,*

---

<sup>5)</sup> An outer measure is an extended real valued, non negative, monotone and countably subadditive set function  $\tau^*$  on the -class of all subsets of  $E$ , such that  $\tau^*(\emptyset) = 0$ .

Proof. By Theorem 1 of [5]  $E$  may be decomposed into the sum of  $n$  or fewer free subsets  $E_\xi (\xi < \varphi_n)$ :

$$E = \bigcup_{\xi < \varphi_n} E_\xi.$$

Since  $I$  is an  $\aleph$ -additive proper ideal it follows the statement of Theorem 2.

**Theorem 3.** *Let  $E$  be a set of power  $\aleph_1$  with  $\aleph_1$  greater than  $\aleph_0$  and less than the first aleph inaccessible in the weak sense, and let  $R$  be a relation between the elements of  $E$  such that for any  $x \in E$  the power of the set  $R(x)$  is smaller than  $\aleph_0$ . Let furthermore  $I$  be an  $\aleph$ -additive proper ideal of  $\kappa$ -subsets of  $E$ , such that  $\{x\} \in I$  for any  $x \in E$ . If  $\{E_\xi\}_{\xi < \omega}$  is a sequence of type  $\omega$ , of subsets of  $E$ , such that  $E_\xi \notin I$  for  $\xi < \omega$ , then there exists a free subset  $E'$  of  $E$  for which  $E' \cap E_\xi \notin I$  for every  $\xi < \omega$ .*

Proof. First we define by finite induction a sequence  $\{F_\xi\}_{\xi < \eta}$  of subsets of  $E$  such that  $F_\xi \notin I$  for  $\xi < \eta$ ,  $F_\xi \cap F_\eta = 0$  if  $\xi \neq \eta$  and for every  $\xi < \omega$  there is a  $r(\xi) < \eta$  such that  $F_{r(\xi)} \sqsubset E_\xi$ . Let  $E_0 = \bigcup_{r < \omega} E_{r(\xi)}$  be a decomposition of  $E_0$  satisfying Theorem 1. Since  $E_0 \cap E_{r(\xi)} = 0$  for  $r \neq \mu$ , for every  $\xi < \omega$  there is at most one  $r = r(\xi) < \omega$  such that  $E_\xi - E_{r(\xi)} \in I$ . It follows that there is an ordinal number  $\nu < \omega_1$  for which  $E_\xi - E_{r(\xi)} \notin I$ , for every  $\xi < \omega$ . Put  $F_0 = E_{r(\xi)}$ . Let  $\beta < \omega$  be a given ordinal number  $\beta > 0$ , and suppose that all sets  $F_\xi$ , where  $0 \leq \xi < \beta$  have been already defined such that  $F_\xi \notin I$  for  $\xi < \beta$  and  $F_\xi \cap F_\eta = 0$ . Put  $E_\xi - \bigcup_{\zeta < \xi} F_\zeta = N_\xi$  ( $\xi \geq \beta$ ). Let  $U = \{\xi; \beta \leq \xi < \omega\}$ . If  $U = 0$ , then we do not define  $F_\beta$ . In this case we put  $\mu = \beta$ . If  $U = 1$ , i.e.  $U = \{k\}$ , then let  $F_\beta = N_k$  and  $\eta = \beta + 1$ . If  $U > 1$ , then we denote by  $q$  the first element of  $U$ . Let  $N_q = \bigcup_{r < \omega_1} N_{qr}$  be a decomposition of  $N_q$  satisfying Theorem 1. Since  $N_{qr} \cap N_{qv} = 0$  for  $v \neq u$ , there is a  $n < \omega$  such that  $N_\xi - N_{qr} \notin I$  for every  $\xi \in U$ . Put  $F_\beta = N_{qr}$ .

It follows from Theorem 2 that  $F_\xi$  has for every  $\xi < \eta$  a free subset  $G_\xi$  such that  $G_\xi \notin I$ . We shall now prove that there is a sequence  $\{H_\xi\}_{\xi < \eta}$  of subsets of  $E$  such that  $H_\xi \subset G_\xi$ ,  $H_\xi \notin I$  ( $\xi < \eta$ ) and  $H_\xi \cap (R[H_\xi] \cup R^{-1}[H_\xi]) = 0$  for  $\xi \neq \zeta$ . The set  $E' = \bigcup_{\xi < \eta} H_\xi$  obviously satisfies Theorem 2.

We define  $H_\xi$  as follows. Let  $G_0 = \bigcup_{\alpha < \omega_1} G_{0\alpha}$  be a decomposition of  $G_0$  satisfying Theorem 1. There is an ordinal number  $\alpha' < \omega_1$  such that  $G_\xi - R^{-1}(G_{0\alpha'}) \notin I$ . In the opposite case there would exist for every  $\alpha$  a natural number  $\xi = \xi(\alpha)$  such that  $G_{\xi(\alpha)} - R^{-1}(G_{0\alpha}) \in I$ . This would imply the existence of a natural number  $\xi$  and a sequence  $\{\alpha_k\}_{k < \omega}$  such that  $\xi = \xi(\alpha_k)$

for every  $k < \omega$ , i. e.  $G_\xi = R^{-1}[G_{\alpha_k}] \in I$  for every  $k < \omega$ . Then there would exist an element  $z \notin G_\xi$ , for which  $z \in R^{-1}[G_{\alpha_k}]$ , i. e.  $R(z) \cap G_{\alpha_k} \neq \emptyset$  for every  $k < \omega$ , which is a contradiction, because  $\overline{R(z)} \triangleleft \aleph_0$ .

Put  $G'_\xi = G_\xi - R^{-1}[G_{\alpha_\xi}] (\xi = 1, 2, \dots)$ . Let  $G'_\xi = \bigcup_{\alpha < \alpha_\xi} G'_{\alpha}$  be a decomposition of  $G'_\xi$  satisfying Theorem 1. Further let

$$U_\alpha = \bigcup_{0 < \xi < \eta, \alpha_\xi < \alpha} G'_{\alpha_\xi}.$$

It is obvious that  $U_{\alpha_1} \cap U_{\alpha_2} = \emptyset$  for  $\alpha_1 \neq \alpha_2$ .

There is a natural number  $r \downarrow$  for which  $G_{0\omega} - R^{-1}[U_r] \notin I$ . For if  $G_{0\omega} - R^{-1}[U_r] \in I$  for every  $n < \omega$ , then there would exist an element  $z \notin G_{0\omega}$  such that  $z \in R^{-1}[U_r] (r = 0, 1, 2, \dots)$  i. e.  $R(z) \cap U_r \neq \emptyset (r = 0, 1, 2, \dots)$ , which is impossible, because  $R(z) \triangleleft \aleph_0$ . Put  $H_0 = G_{0\omega} - R^{-1}[U_r]$ . It is obvious that

$$N_\xi = G'_{\xi r} - R[H_0] - R^{-1}[H_0] \notin I \quad (\xi = 1, 2, \dots)$$

We define  $H_1$  starting from  $N_1$  in the same way as  $H_0$  is defined starting from the set  $G_0$ . Obviously we can continue this process for every  $r < \eta \downarrow$ . Thus we obtain the sequence  $\{H_r\}_{r < \eta}$  satisfying our requirement. The theorem is proved.

**Corollary 4.** *If  $2^{\aleph_0}$  is less than the first aleph inaccessible in the weak sense,  $E$  is the set of the real numbers and  $R$  is a relation between the elements of  $E$  such that for any  $x \in E$  the power of the set  $R(x)$  is smaller than  $\aleph_0$ , then there exists a free subset  $E'$  of  $E$ , which is everywhere of the second category.*

**Proof.** Let  $I$  be the set of the subsets of  $E$  of the first category, and  $\{E_\xi\}_{\xi < \omega}$  a sequence of type  $\omega_1$  of all intervals of  $E$  with rational endpoints, and apply Theorem 3.

**Corollary 5.** *Under the same hypotheses as in Corollary 4 there exists a free subset  $E'$  of  $E$  such that*

$$\mu^*(E' \cap [n, b]) \neq 0$$

for every interval  $[a, b]$  of  $E$ ,  $\mu^*$  denoting Lebesgue outer measure.

**Proof.** Let  $I$  be the set of all subsets of measure zero of  $E$  and  $\{E_\xi\}_{\xi < \omega}$  a sequence of type  $\omega_1$  of all intervals of  $E$  with rational endpoints, and apply Theorem 3.

## II.

We assume in this section that  $E$  is a metric space and condition (B) holds.

First we prove the following

**Theorem 4.** *Let  $E$  be the set of all real numbers and  $R$  a relation between the elements of  $E$  such that, for any  $x \in E$ , the power of the set  $R(x)$  is smaller than  $\aleph_0$ . Then there exists a free subset  $E'$  of  $E$  such that  $E'$  is everywhere of the second category.*

Proof. Let  $(a, b)$  be an arbitrary interval of  $E$  and  $A^{(a, b)}$  the set of all subsets of  $(a, b)$  the complements of which are of the first category and  $F_\alpha$ . Let further  $\{C_\gamma\}_{\gamma < \omega_1}$  be a wellordering of the set

$$\bigcup_{(a, b) \subseteq I_\gamma} A^{(a, b)}.$$

of the type  $\varphi_c$  (where  $c = 2^{\aleph_0}$ ) and  $I_\gamma$  the interval corresponding to the set  $C_\gamma$ .

We consider the set  $H$  of all the series  $H = \{a_\xi\}_{\xi < \varphi_c}$  of elements with the properties :

- a)  $a_\xi \in C_\xi$  or  $a_\xi = 0$ ;  $\xi < \varphi_c$ ;
- b) if  $a_\xi \neq 0$ , then  $a_\nu \neq 0$  for  $\nu < \xi$ ;
- c) if  $a_\xi \neq 0$  and  $a_\nu \neq 0$ , then  $a_\xi \neq a_\nu$  for  $\xi < \nu$ ;
- d) the set of the elements of the series is a free set.

For any  $H \in H$ , let  $\tilde{H}$  denote the set of the elements of  $H$ .

We say that an element  $H \in H$  is maximal with respect to the relation  $R$  if  $\nu_d$  is the smallest ordinal number  $\xi < \varphi_c$  such that  $a_{\nu_d} = 0$  and there is no element  $k \in C_{\nu_d} - R[\tilde{H}]$  such that  $k$  and the elements  $\neq 0$  of  $H$  are independent or if  $a_\nu \neq 0$  for every  $\nu < \varphi_c$ . We define the index of  $H$  in the first case as  $\nu_d$  and in the second case as  $\varphi_c$ . Let  $H'$  be the set of the maximal elements of  $H$ .

We say that two series  $H_1$  and  $H_2$  are mutually exclusive if  $H_1 \cap H_2 = \emptyset$ .

Let  $\{H_\nu\}_{\nu < \eta}$  be a sequence of type  $\eta < \omega_1$  of mutually exclusive elements of  $H'$  with indices  $\delta_\nu < \varphi_c$ . Then by the definition of  $H'$ ,  $\tilde{H}_\nu < c$ ; consequently  $\overline{R[\tilde{H}_\nu]} < c$  for every  $\nu < \eta$ . Since  $\eta < \omega_1$  by a well-known theorem of J. KÖNIG we have

$$\overline{\bigcup_{\nu < \eta} (H_\nu \cup R[\tilde{H}_\nu])} < c,$$

i. e.

$$\overline{C_\gamma - \bigcup_{\nu < \eta} (\tilde{H}_\nu \cup R[\tilde{H}_\nu])} < c$$

for every  $\gamma < \varphi_c$ . It follows that there is an element  $H_\eta$  of  $H'$  such that  $\tilde{H}_\eta \neq 0$  and  $H_\eta \mathbf{n} \tilde{H}_\eta = 0$  for every  $\nu < \eta$ .

(1) } For every  $\delta < \varphi_c$  there is only a finite number of mutually exclusive elements of  $H'$  with the same index  $\delta$ .

Let  $\{H_n\}_{n<\omega}$  be a sequence of type  $\omega$ , of mutually exclusive elements of  $H'$ . Suppose that the series  $H_n$  ( $n = 1, 2, \dots$ ) have the same index  $\delta$ . Then the set  $C_\gamma - \bigcup_{n<\omega} \tilde{H}_n - \bigcup_{n<\omega} R[\tilde{H}_n]$  is non empty and for every element  $\alpha$  of this set  $R(\alpha) \geq \aleph_0$  hold, because  $R(\alpha) \mathbf{n} \tilde{H}_n \neq 0$  ( $n = 1, 2, \dots$ ), which is a contradiction.

Supposing that every element of  $H'$  has an index smaller than  $\varphi_c$  we can choose by (1) a sequence  $\{H_\nu\}_{\nu<\omega_1}$  of mutually exclusive elements of  $H'$  of type  $\omega_1$  such that the indices  $\beta_\nu$  of the series  $H_\nu$  are distinct. Corresponding to every interval  $I_\eta$  we choose in  $I_\eta$  a subinterval  $I'_\eta$  with rational endpoints. Since  $\{\beta_\nu\}_{\nu<\omega_1} > \aleph_0$  and  $\{I'_\gamma\}_{\gamma<\varphi_c} \leq \aleph_0$  there is an  $I'_{\gamma_0}$  and a subsequence  $\{\beta_{\nu_k}\}_{k<\omega}$  of type  $\omega$ , of  $Z = \{\beta_\nu\}_{\nu<\omega_1}$  such that  $I'_{\beta_{\nu_k}} = I'_{\gamma_0}$  for every  $k < \omega$ . Obviously the complement of the set  $L_{\gamma_0} = \bigcap_{k<\omega} C_{\beta_{\nu_k}}$  is of the first category with respect to  $I'_{\gamma_0}$ . Consequently the power of  $L_{\gamma_0}$  is  $c$ , thus

$$\overline{L_{\gamma_0} - \bigcup_{k<\omega} (\tilde{H}_{\nu_k} \cup R[\tilde{H}_{\nu_k}])} = c$$

It follows that there is an element  $z \in L_{\gamma_0} - \bigcup_{k<\omega} (\tilde{H}_{\nu_k} \cup R[\tilde{H}_{\nu_k}])$  such that  $R(z) \mathbf{n} \tilde{H}_{\nu_k} \neq 0$  ( $k = 1, 2, \dots$ ) i. e.  $R(z) \geq \aleph_0$  which is impossible, because  $R(z) < \aleph_0$ . Thus there is a free subset  $E'$  of  $E$  such that  $E' \cap C_\gamma \neq 0$  for every  $\gamma < \varphi_c$ . It is clear that  $E'$  is of the second category. The theorem is proved.

Theorem 5. Let  $E$  be the set of all real numbers and  $R$  a relation between the elements of  $E$  such that for any  $x \in E$  the power of the set  $R(x)$  is smaller than  $\aleph_0$ . Then there exists a free subset  $E'$  of  $E$  such that the Lebesgue outer measure  $\mu^*(E')$  of  $E'$  in every interval  $(a, b)$  is  $b-a$ .

Proof. Let  $(a, b)$  be an arbitrary interval of  $E$  and  $\mathbf{B}^{(a, b)}$  the set of all subsets of  $(a, b)$  of positive measure  $> \frac{1}{2}(b-a)$  and  $G_\delta$ . Let further  $\{D_\gamma\}_{\gamma<\varphi_c}$  be a wellordering of the set

$$\bigcup_{(a, b) \subseteq E} \mathbf{B}^{(a, b)}$$

of type  $\varphi_c$ , and  $I_\eta$  the interval  $(a, b)$  corresponding to  $D_\eta$ . We can prove completely analogously to the proof of the theorem 4 the existence of a free set  $E'$  such that

$$E' \cap D_\gamma \neq 0 \quad (\gamma < \varphi_c),$$

if we select in every interval  $I_\gamma = (a, b)$  an interval  $I'_\gamma = (a', b')$  with rational endpoints such that  $b' - a' > \frac{3}{4}(b - a)$ . Obviously the outer measure of  $E$  in every interval  $(a, b)$  is  $b - a$ .

It is easy to see by the method of the proofs of theorems 4 and 5 that the following theorem is valid too.

**Theorem 6.** *Let  $E$  be the set of all real numbers and  $R$  a relation between the elements of  $E$  such that for any  $x \in E$  the power of the set  $R(x)$  is smaller than  $\aleph_0$ . Then there exists a free subset  $E'$  of  $E$  such that  $E'$  is everywhere of the second category and the Lebesgue outer measure  $\mu(E')$  of  $E$  in every interval  $(a, b)$  is  $b - a$ .*

**Theorem 7.** *Let  $E$  be an interval of the set of all real numbers and suppose that there exists a relation  $R$  between the elements of  $E$ . Let further  $B$  be a  $n$ -algebra of subsets of  $E$  containing all subintervals of  $E$  and  $\mu$  a nof identically zero measure on  $B$ . If  $g(x) = d(x, R(x)) > 0$  for every  $x \in E$  and if*

*(C) there exists a real number  $i > 0$  such that the set  $\{x: g(x) \geq i\}$  contains in  $B$  a subset of positive  $\mu$ -measure,*  
*then there exists in  $B$  a free subset of  $E$  of positive  $\mu$ -measure.*

*If, for every  $x \in E$ , the set  $R(x)$  is the complement of an interval of  $E$  whose center is at  $x$ , then the condition (C) is not only sufficient, but also necessary for the existence of a free subset, of positive  $\mu$ -measure, of  $E$  in  $B$ .*

**Proof.** Let  $A$  be a subset of  $\{x: g(x) \geq i\}$  satisfying the condition (C). Let

$$x_1, x_2, \dots, x_n, \dots$$

be an enumeration of the set of rational numbers in  $E$ . For every element  $x \in E$  and  $\varepsilon > 0$  there exists an element  $x_{n_0}$  of this sequence such that  $d(x, x_{n_0}) < \varepsilon$ . For every  $n = 1, 2, \dots$  let  $U(x_n, i)$  be the open interval of length  $i$  whose center is at  $x_n$ . It is obvious that

$$\bigcup_n U(x_n, i) = E$$

Let  $A_n = A \cap U(x_n, i)$  ( $n = 1, 2, \dots$ ). Since  $U(x_n, i) \in B$  and  $A \in B$ ,  $A_n \in B$ . Let  $A^* = A_n - \bigcup_{n' \neq n} A_{n'}$  ( $n = 1, 2, \dots$ ). Since  $\mu$  is countably additive and  $\mu(A) > 0$ , there exists an index  $n'$  for which  $\mu(A_{n'}^*) > 0$ . It follows that  $\mu(A_{n'}) > 0$ . The set  $A_{n'}$  is free, because if  $x \in A_{n'}$  and  $y \in R(x)$ , then  $d(x, y) > g(x) \geq i$ .

For every  $x \in E$ , let  $U(x)$  be an interval whose center is at  $x$  and  $R(x) = E - U(x)$ . In this case condition '(C)' is also necessary for the existence of a free subset of positive  $\mu$ -measure in  $B$ , i. e. if there is in  $B$  a

free subset A of  $E$  such that  $\mu(A) > 0$ , then there exists a positive number  $i$ , for which the set  $\{x : g(x) \geq i\}$  contains in B a set of positive p-measure. Suppose the contrary. Then B contains a free subset of positive p-measure, but for every  $i > 0$  the set  $\{x : g(x) \geq i\}$  contains in B only such subsets  $F$  for which  $p(F) = 0$ . Let  $\alpha$  denote the diameter of the set A. Put

$$E_\alpha = \left\{ x : g(x) \geq \frac{\alpha}{2} \right\}.$$

By the hypothesis  $E_\alpha$  contains in B only such subsets  $F$ , for which  $p(F) = 0$ . Let  $F_1 = E_\alpha \cap A$  and  $F_2 = E_\alpha \cap (E-A)$ . Since A is free and  $R(x) = E - U(x)$  for every  $x \in E$ , we have  $g(x) \geq \frac{\alpha}{2}$  for every  $x \in A$ . Thus  $F_1 = A$ . By the definition,  $F_1 \cup F_2 = E_\alpha$ , therefore  $A = F_1 \subset E_\alpha$ . Since  $A \in B$ , it follows that  $p(A) = 0$ , which contradicts to  $\mu(A) > 0$ . The theorem is proved.

**Remark 1.** In general the condition (C) is not necessary. Consider the interval  $[0,1]$ . Let  $\mu^*$  and  $\mu_*$  denote the Lebesgue outer and inner measure, respectively. We can define the relation  $R$  such that the interval  $[0,1]$  contains a free subset of positive Lebesgue measure and

$$\mu_*(\{x : g(x) \geq i\}) = 0$$

for any  $i > 0$ , where  $g(x) = d(x, R(x))$ . We shall use the following theorem (see [7]):

The set  $E$  of the real numbers has a subset  $E'$  with the following properties :

1. for every interval  $(a, b)$  of  $E$ ,  $\mu^*(E' \cap (a, b)) = b-a$ ,
2.  $E$  can be decomposed into enumerable many sets  $E_n$  ( $n = 1, 2, \dots$ ) without common points, which are all superposable by shifting the set  $E'$ .

It follows that  $[0,1]$  can be decomposed into the sum of enumerable many sets  $S_n$  ( $n = 1, 2, \dots$ ) such that  $\mu^*(S_n) = 1$  ( $n = 1, 2, \dots$ )

For every  $x \in S_n$ , let  $K(x)$  be the open interval of length  $\frac{2}{n}$  whose center is at  $x$ . We define  $R$  as follows. Let  $N$  be the set of rational numbers and

$$R(x) = (E - K(x)) \cap N.$$

Obviously

$$g(x) = + \text{ for } x \in S_n.$$

If  $i > 1$ , then  $V_i = \{x : g(x) \geq i\} = \emptyset$ . If  $i \leq 1$ , then  $V_i \subseteq V_1 = S_1 \cup S_2 \cup \dots \cup S_{n+1}$  for some natural numbers  $n > 0$ . We have  $\mu_*(V_i) = 0$  because  $\mu_*(V_{\frac{i+1}{n+1}}) = \mu^*([0,1] - V_{\frac{i+1}{n+1}}) = 0$ .

It follows from the definition of  $R$  that the set  $U$  of the irrational numbers of  $[0,1]$  is a free set.  $U$  is measurable and  $\mu(U) = 1$ .

**R e m a r k 2.** It is easily seen that Theorem 7 remains true for a separable metric space. The following counter-example shows that for non-separable metric spaces this theorem is generally not true.

Consider the following example of ALEXANDROFF [9]. Let  $S$  be the plane with the ordinary (euclidean) metric  $d = d(x, y)$ . We define now a new distance as follows. Let  $\bar{0}$  be a given point of  $S$ ,  $x$  and  $y$  two arbitrary points of  $S$  and

$$d'(x, y) = \begin{cases} d(x, y) & \text{if } \bar{0} \text{ lies on the line } xy, \\ d(x, \bar{0}) + d(y, \bar{0}) & \text{if } \bar{0} \text{ does not lie on the line } xy. \end{cases}$$

Thus we obtain a new metric space  $S'$ , which is not separable.

Let  $\mu^*$  be the ordinary Lebesgue outer measure for the subsets of  $S$ . We define a relation  $R$  between the elements of  $S'$  as follows. If  $x = \bar{0}$ , then let  $R(x) = 0$ . If  $x \neq \bar{0}$ , then let  $r$  be a real number for which  $0 < r < d(x, \bar{0})$ ,  $E(x) = \{y : d'(x, y) < r\}$  and  $R(x) = S - E(x)$ . It follows from the definition of the distance  $d'$  that if  $x, y \in S' (x \neq y)$  and  $\bar{0}$  does not lie on the line  $xy$ , then either  $x \notin R(y)$  or  $y \notin R(x)$  i. e.  $x$  and  $y$  are not independent. Hence each free subset of  $S'$  lies on a line containing  $\bar{0}$ . But for every line  $L$ ,  $\mu^*(L) = 0$ . Thus for every free subset  $E'$ ,  $\mu^*(E') = 0$ .

For non-separable metric spaces we state the following

**Theorem 8.** Let  $E$  be a metric space. Suppose that  $E$  contains a dense subset, the power of which is less than the first aleph inaccessible in the weak sense. Let  $\mu$  be a  $\sigma$ -finite measure on the set  $\mathbf{B}$  of all Borel subsets which is not identically zero. If  $g(x) = d(x, R(x)) > 0$  for every  $x \in E$  and if condition (C) holds, then there exists in  $\mathbf{B}$  a free subset of positive  $\mu$ -measure of  $E$ .

If, for every  $x \in E$ , the set  $R(x)$  is the complement of an sphere of  $E$  whose center is at  $x$ , then the condition (C) is not only sufficient, but also necessary for the existence of a free subset of positive  $\mu$ -measure of  $E$  in  $\mathbf{B}$ .

**P r o o f.** If  $\mu$  is a  $\sigma$ -finite measure on the set of all Borel subsets of  $E$  and  $E$  contains a dense subset, the power of which is less than the first aleph inaccessible in the weak sense, then there exists a decomposition

$$E = N \cup M$$

of  $E$  into two mutually disjoint sets such that  $\mu(N) = 0$  and  $M$  is separable (where  $N$  is the sum of all open subsets of  $\{t\}$ -measure zero of  $E$ ) (see [10]). It is clear that  $\mu$  is not identically zero on  $M$ , since  $\mu(N) = 0$  and

$$\mu(N) + \mu(M) = \mu(E) \neq 0.$$

Let  $X$  be an arbitrary Borel subset of  $E$ . Since  $X \cap M = X - N$  is a Borel subset of  $E$ ,

$$\mu(X \cap M) = \mu(X) - \mu(N) = \mu(X)$$

Let  $B'$  be the set of all sets of the form  $X \cap M$ , where  $X \in B$ , and let  $r(X) = \mu(X)$  for  $X \in B$ . Hence, if the set  $\{x : g(x) \geq i\}$  contains in  $B$  a set of positive  $\mu$ -measure, then it contains in  $B'$  a set of positive  $\mu$ -measure too. Since  $B' \subseteq B$ , the converse of this statement is also true. Thus, it is sufficient to prove the theorem for  $M, B'$  and  $r$ , instead of  $E, B$  and  $\mu$ . Since  $M$  is a separable metric space and  $B'$  is a  $\sigma$ -algebra and  $\mu$  is not identically zero measure on  $B'$ , the theorem is true for  $M, B'$  and  $r$ . Thus the theorem is true for  $E, B$  and  $\mu$  too.

### III.

We deal in this section with the problem (ii).

**Theorem 9.** *Let  $E$  be a set of power  $m \geq \aleph_0$  and  $K$  a class of power  $m$ , of subsets of  $E$  of power  $m$ . There exists a relation  $R$  between the elements of  $E$  such that for every  $x \in E$  the power of the set  $R(x)$  is  $\leq 1$  and there is no free subset  $X$  in  $K$  with respect to  $R$ .*

**Proof.** Let

$$B_0, B_1, \dots, B_\omega, \dots, B_\xi, \dots \quad (\xi < \varphi_m)$$

be a wellordering of  $K$  of the type  $\varphi_m$ . Since  $B_\xi = m$  for every  $\xi < \varphi_m$ , there exist two sequences  $\{x_\xi\}_{\xi < \varphi_m}$  and  $\{y_\xi\}_{\xi < \varphi_m}$  such that

1.  $x_\xi \in B_\xi$  and  $y_\xi \in B_\xi$  for every  $\xi < \varphi_m$ ,
2.  $x_\xi \neq x_\zeta$  and  $y_\xi \neq y_\zeta$  for  $\xi < \zeta < \varphi_m$ ,
3.  $x_\xi \neq y_\xi$  for every  $\xi < \varphi_m$ .

We define  $R$  as follows : let  $R(x_\xi) = \{y_\xi\}$  for every  $\xi < \varphi_m$  and if  $x \neq x_\xi$  ( $\xi < \varphi_m$ ), then let  $R(x) = \{x_0\}$ . It is obvious that the sets  $B_\xi$  are not free.

**Corollary 6.** *Let  $E$  be the set of all real numbers. There exists a relation  $R$  between the elements of  $E$  such that for every  $x \in E$  the power of the set  $R(x)$  is  $\leq 1$  and there is no perfect free subset of  $E$ .*

**Corollary 7.** *Let  $E$  be the set of all real numbers. There exists a relation  $R$  between the elements of  $E$  such that for every  $x \in E$  the power of the set  $R(x)$  is  $\leq 1$  and there is no free Borel subset of  $E$  of power  $2^{\aleph_0}$ .*

**Theorem 10.** *Let  $E$  be a set of power  $m \geq \aleph_0$  and  $K$  a set of power  $m$ , of mutually disjoint non empty subsets of  $E$ . There exists a relation  $R$  between the elements of  $E$  such that, for every  $x \in E$  the power of the set  $R(x)$  is  $\leq 1$  and there is no such free set which has non empty intersection with every element of  $K$ .*

Proof. Let

$$B_0, B_1, \dots, B_\omega, \dots, B_\xi, \dots \quad (\xi \triangleleft \varphi_m)$$

be a wellordering of  $K$  of the type  $\varphi_m$ . Let further

$$x_0, x_1, \dots, x_\omega, \dots, x_\xi, \dots \quad (\xi \triangleleft \varphi_m)$$

be a wellordering of  $E$  of the type  $\varphi_m$ . Obviously, we may assume that  $x_\xi \notin B_\xi$ . We define  $R$  as follows: let

$$R^{-1}(x_\xi) = B_\xi.$$

Let  $F$  be a set which has non empty intersection with every element of  $K$ :

$$F \cap B_\xi \neq \emptyset \quad (\xi < \varphi_m).$$

Let  $x \in F$ . There is an ordinal number  $\eta \triangleleft \varphi_m$  such that  $x = x_\eta$ . Since  $R^{-1}(x) = B_\eta$ , we have  $b_\eta Rx$  for every  $b_\eta \in B_\eta \cap F$ . It follows that  $x$  and  $b_\eta$  ( $x \neq b_\eta$ ) are not independent, because  $x \in R(b_\eta)$ . The theorem is proved.

*Corollary 8.* If  $E$  is the set of all real numbers, then there exists a relation  $R$  between the elements of  $E$  such that, for every  $x \in E$ , the power of the set  $R(x)$  is  $\leq 1$  and there is no free subset, the complement of which is totally imperfect.

Proof. Let  $K$  be a set of power  $2^{\aleph_0}$  of non empty mutually disjoint perfect subsets of  $E$ .  $T$  a set the complement  $CT$  of which is totally imperfect, and  $K \in K$ . Since the set  $CT$  does not contain  $K$ ,  $K \cap T \neq \emptyset$ . The corollary is proved.

Finally we prove

*Theorem 11.* Let  $E$  be a set of power  $m \geq \aleph_0$  and  $\mathbf{K}$  a class of power  $g \triangleleft m$ , of mutually exclusive subsets of power  $m$  of  $E$ . If  $R$  is a relation between the elements  $x \in E$  for which the condition (A) holds, i. e.  $\overline{R(x)} \triangleleft n \triangleleft m$  for every  $x \in E$ , then there exists a free subset  $E'$  of  $E$  such that, for every  $K \in \mathbf{K}$ ,

$$\overline{KnE'} = m.$$

Proof. Let

$$K_0, K_1, \dots, K_\omega, K_{\omega+1}, \dots, K_\xi, \dots \quad (\xi \triangleleft \varphi_m)$$

be a wellordering of  $K$  of the type  $\varphi_m$ . We assume first that  $m$  is regular. We consider the set  $M$  of the matrices

$$M = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1\xi} & \dots \\ a_{21} & a_{22} & \dots & a_{2\xi} & \dots \\ \vdots & \vdots & & \vdots & \\ a_{\eta 1} & a_{\eta 2} & \dots & a_{\eta \xi} & \dots \\ I & \vdots & & \ddots & \end{bmatrix}$$

<sup>6</sup> S. MARCUS has found independently the results of our corollaries 6 and 8.

of elements with the properties:

1.  $a_{\eta\xi} \notin K_\xi$  or  $a_{\eta\xi} = 0$ ,  $\eta \triangleleft \varphi_m$  and  $\xi \triangleleft \varphi_n$
2. if  $a_{\eta\xi} \neq 0$  then  $a_{\nu\mu} \neq 0$  for  $\nu = \eta$  and  $\mu \triangleleft \xi$  or  $\nu \triangleleft \eta$  and  $\mu \triangleleft \xi$
3. if  $a_{\nu\mu} \neq 0$  and  $a_{\xi\delta} \neq 0$ , then  $a_{\nu\mu} \neq a_{\xi\delta}$  for  $\nu \neq \xi$
4. the set of the elements of the matrix is a free set.

For any  $M \in \mathbf{M}$ , let  $\tilde{M}$  denote the set of the elements of  $M$ .

We say that an element  $M \in \mathbf{M}$  is *maximal with respect to the relation R* if  $\mu_0$  and  $\nu_0$  are the smallest ordinal numbers  $\triangleleft \varphi_m$  such that  $a_{\mu_0\nu_0} = 0$  and there is no element  $k \in K_{\nu_0} - R[\tilde{M}]$  such that  $k$  and the elements  $\neq 0$  of the matrix  $M$  are independent or if  $a_{\mu\nu} \neq 0$  for every  $\mu \triangleleft \varphi_m$  and  $\nu \triangleleft \varphi_n$ . We define the *index* of  $M$  in the first case as  $\nu_0$  and in the second case as  $\varphi_n$ . Let  $M'$  be the set of the maximal elements of  $\mathbf{M}$ .

We say that two matrices  $M_1$  and  $M_2$  are mutually exclusive if  $\tilde{M}_1 \amalg \tilde{M}_2 = 0$ .

Let  $\{M_\nu\}_{\nu < \varphi_n}$  be a sequence of type  $\eta \triangleleft \varphi_m$  of mutually exclusive elements  $M_\nu$  of  $M'$  with indices  $\delta_\nu \triangleleft \varphi_n$ . Then by the definition of  $M'$ ,  $\tilde{M}_\nu < \mathfrak{m}$ , consequently  $K[\tilde{M}_\nu] < \mathfrak{m}$  for every  $\nu \triangleleft \eta$ , because  $f(x) \triangleleft \nu \triangleleft \eta$ .

Since  $\mathfrak{m}$  is regular,

$$\overline{\bigcup_{\nu < \eta} (\tilde{M}_\nu \cup R[\tilde{M}_\nu])} \triangleleft \mathfrak{m}$$

i. e.

$$\overline{K_\gamma - \bigcup_{\nu < \eta} (\tilde{M}_\nu \cup R[\tilde{M}_\nu])} < \mathfrak{m},$$

for every  $\gamma \triangleleft \varphi_n$ . It follows that there is an element  $M_\eta \notin M'$  such that  $\tilde{M}_\eta \neq 0$  and  $\tilde{M}_\eta \cap \tilde{M}_\nu = 0$  for every  $\nu \triangleleft \eta$ .

(2) { For every  $\delta \triangleleft \varphi_n$  there are less than  $\mathfrak{m}$  mutually exclusive elements of  $M'$  with the same index  $\delta$ .

Let  $\{M_\nu\}_{\nu < \varphi_n}$  be a sequence of the type  $\varphi_n$  of mutually exclusive elements  $M_\nu$  of  $M'$  with the same index  $\delta$ . Then the set

$$K_\delta = \overline{\bigcup_{\nu < \varphi_n} (\tilde{M}_\nu \cup R[\tilde{M}_\nu])}$$

is non empty and, for every element  $z$  of this set,  $R(z) \cong \mathfrak{m}$  because, by the definition of  $M'$ ,  $R(z) \amalg \tilde{M}_\nu \neq 0$  for  $\nu < \varphi_n$ , which is a contradiction. Thus (2) is proved.

Supposing that every element  $M$  of  $M'$  has an index smaller than  $\varphi_n$ , we can now define by transfinite induction a sequence  $\{M_\nu\}_{\nu < \varphi_m}$  of mutually exclusive elements of  $M'$  of the type  $\varphi_m$ . Since  $\mathfrak{g} \triangleleft \mathfrak{m}$  and  $\mathfrak{m}$  is regular, there exists a subset, of power  $\mathfrak{m}$ , of  $M'$  with the same index  $\triangleleft \varphi_n$  which contra-

dicts to (2). Thus there exists a matrix of index  $\varphi_{\mathfrak{m}^*}$ . It is obvious that the set of elements of this matrix satisfies the requirement of the theorem. Thus the theorem is true, if  $\mathfrak{m}$  is regular.

Consider now the case when  $\mathfrak{m}$  is singular'). We assume that the generalised continuum hypothesis is true. Let

$$\mathfrak{m} = \sum_{\xi < \varphi_{\mathfrak{m}^*}} \mathfrak{m}_\xi$$

be a decomposition of  $\mathfrak{m}$  such that

- 1)  $\mathfrak{m}_\xi$  is regular for every  $\xi < \varphi_{\mathfrak{m}^*}$ ,
- 2)  $\mathfrak{m}_\xi < \mathfrak{m}_\zeta$  for  $\xi < \zeta < \varphi_{\mathfrak{m}^*}$ ,
- 3)  $\mathfrak{m}_\xi \geq \max \{g_i \text{ it, } \mathfrak{m}^*\}$ ,
- 4)  $2^{\sum_{\xi < \xi} \mathfrak{m}_\xi} < \mathfrak{m}_\xi$  for every  $\xi < \varphi_{\mathfrak{m}^*}$ .

Let further

$$K_\nu = \bigcup_{\xi < \varphi_{\mathfrak{m}^*}} K_{\nu\xi} \quad (\nu < \varphi_{\mathfrak{m}})$$

be a decomposition of  $K_\nu$  into mutually exclusive subsets of  $K_\nu$  such that  $K_{\nu\xi} = \mathfrak{m}_\xi$ .

By the first part of the theorem, there exists a free subset  $L_\xi$  of  $E$  for every  $\xi < \varphi_{\mathfrak{m}^*}$  such that

$$\overline{L_\xi \cap K_{\nu\xi}} = \mathfrak{m}_\xi$$

for every  $\eta < \varphi_{\mathfrak{m}}$ . Omit for  $\xi < \eta$  all the elements of  $R[L_\xi]$  from  $L_\eta$ . Thus we get the sets

$$L'_\eta = L_\eta - \bigcup_{\xi < \eta} R[L_\xi].$$

By 1) and 3)  $\bigcup_{\xi < \eta} \overline{R[L_\xi]} < \mathfrak{m}_\eta$ , thus the power of the set  $L'_\eta$  is  $\mathfrak{m}_\eta$  and  $L'_\eta \cap \overline{K_{\nu\eta}} = \mathfrak{m}_\eta$  for 'every'  $\eta < \varphi_{\mathfrak{m}}$ . Obviously

$$R[L'_\xi] \cap \left( \bigcup_{\eta < \xi} L'_\eta \right) = 0.$$

Let

$$L'_{\nu\xi} = L'_\xi \cup K_{\nu\xi} \quad (\nu < \varphi_{\mathfrak{m}}, \xi < \varphi_{\mathfrak{m}^*}).$$

We want to construct sets  $L'_{\nu\xi}$  of power  $\mathfrak{m}_\xi$  which satisfy

$$(3) \quad R[L'_{\nu\xi}] \cap \left( \bigcup_{\kappa < \nu} \bigcup_{\eta < \xi} L''_{\kappa\eta} \right) = 0.$$

But then clearly

$$R \left[ \bigcup_{\nu < \varphi_{\mathfrak{m}}} \bigcup_{\xi < \varphi_{\mathfrak{m}^*}} L'_{\nu\xi} \right] \cap \left[ \bigcup_{\nu < \varphi_{\mathfrak{m}}} \bigcup_{\xi < \varphi_{\mathfrak{m}^*}} L''_{\nu\xi} \right] = 0,$$

i. e. the set  $\bigcup_{\nu < \varphi_{\mathfrak{m}}} \bigcup_{\xi < \varphi_{\mathfrak{m}^*}} L'_{\nu\xi}$  is free and satisfies the requirement of the theorem. Thus we only have to construct  $L'_{\nu\xi}$ . Consider the sets  $L'_{\nu\xi}$  and

---

<sup>6)</sup> The proof is due to A. HAJNAL.

$L_\xi^* = \bigcup_{r \in \varphi_\xi} \bigcup_{\zeta < \xi} L'_{r\zeta}$  ( $\xi < q_{m^*}$ ). Let  $N[L_\xi^*]$  denote the set of all subsets of  $L_\xi^*$  of the power  $\xi$ . By 3)  $\overline{N[L_\xi^*]} \subset m_\xi$ . It follows that there exists a subset  $H_{r\xi}$  of power  $m_\xi$  of  $L'_{r\xi}$  and an element  $N_{r\xi}$  of  $N[L_\xi^*]$  such that  $L_\xi^* \cap R[H_{r\xi}] = N_{r\xi}$ . Let

$$U = \bigcup_{r < \varphi_\xi} \bigcup_{\zeta < q_{m^*}} N_{r\zeta}.$$

Obviously  $\overline{U} \leq n g m^* < m_0$ . Let  $L''_{r\xi} = H_{r\xi} - U$  ( $r < \varphi_\xi$  and  $\xi < q_{m^*}$ ). These sets obviously satisfy the condition (3). The theorem is proved.

### References.

- [1] S. PICCARD, Sur un problème de M. Ruziewicz, de la théorie des relations, *Fundamenta Math.*, 29 (1937), 5-9.
- [2] G. FODOR, On a theorem in the theory of binary relations, *Compositio Math.*, 8 (1951), 250.
- [3] G. FODOR, On two problems concerning the theory of binary relations, *Publicationes Math. Debrecen*, 1 (1949-50), 199-200.
- [4] G. FODOR, On a problem concerning the theory of binary relations, *Nieuw Archief voor Wiskunde*, 23 (1951), 247-248.
- [5] G. FODOR, Proof of a conjecture of P. Erdős, *Acta Sci. Math.*, 14 (1952), 219-227.
- [6] S. ULAM, Zur Maßtheorie in der allgemeinen Mengenlehre, *Fundamenta Math.*, 16 (1930), 140-150.
- [7] W. SIERPIŃSKI, Sur un ensemble non mesurable, *Tôhoku Math. Journal*, 12 (1914), 205 - 208.
- [8] W. SIERPIŃSKI, *Hypothèse du continu* (Warszawa—Lwów, 1934).
- [9] П. С. АЛЕКСАНДРОВ, Введение в общую теорию множеств и функций (Москва, 1948).
- [10] E. MARCZEWSKI-R. SIKORSKI, Measures in non-separable metric spaces, *Colloquium Math.*, 1 (1947-1948), 133-139.
- [11] P. R. HALMOS, *Measure theory* (New York, 1950).

(Received September 5, 1957.)