

On the Lipschitz's condition for Brownian motion.

By K. L. CHUNG*, P. ERDÖS and T. SIRAO

(Received April 9, 1959)

(Revised May 25, 1959)

Let $X(t)$ ($0 \leq t < \infty$) be the Brownian motion process. Concerning the uniform continuity of $X(t)$, there exists P. Lévy's result. Before stating his result, let us define the concept of upper class and lower class with regard to the uniform continuity of $X(t)$ ($0 \leq t \leq 1$).

If there exists a positive number ε such that $|t' - t| \leq \varepsilon$ implies the relation

$$(1) \quad |f(t') - f(t)| \leq g(|t' - t|),$$

where $g(t)$ is a non-negative, continuous, non-decreasing function defined in some finite interval $(0, T)$ and vanishing with t , then we say that $f(t)$ satisfies Lipschitz's condition relative to $g(t)$. Putting $\varphi(t) = \psi\left(\frac{1}{t}\right)\sqrt{t}$, if $X(t)$ ($0 \leq t \leq 1$) satisfies Lipschitz's condition relative to $\varphi(t)$ with probability 1 we say that $\psi(t)$ belongs to the upper class. If $X(t)$ ($0 \leq t \leq 1$) does not satisfy Lipschitz's condition relative to $\varphi(t)$ with probability 1 we say that $\psi(t)$ belongs to the lower class. P. Lévy [1] proved that the function

$$\psi(t) = c(2 \log t)^{\frac{1}{2}}$$

belongs to the upper class for $c > 1$ and belongs to the lower class for $c < 1$. Following his method, T. Sirao [2] improved the result as follows: The function

$$\psi(t) = (2 \log t + c \log \log t)^{\frac{1}{2}}$$

belongs to the upper class for $c > 5$ and belongs to the lower class for $c < -1$. In this paper we shall prove the following theorems.

THEOREM 1. *A non-negative, continuous and monotone non-decreasing function $\psi(t)$ belongs to the upper or lower class according as the integral*

$$(2) \quad \int_0^\infty \psi^3(t) e^{-\frac{1}{2}\psi^2(t)} dt$$

is convergent or divergent.

* Research supported in part by the United States Air Force through the Office of Scientific Research under contract AF 49 (638)-265.

THEOREM 2. The function $\psi(t)$ defined by

$$(3) \quad \psi(t) = (2 \log t + 5 \log_{(2)} t + 2 \log_{(3)} t + \cdots + 2 \log_{(n-1)} t + c \log_{(n)} t)^{\frac{1}{2}},$$

where $\log_{(n)} t$ denotes the n -times iterated logarithm, belongs to the upper class for $c > 2$ and to the lower class for $c \leq 2$.

These theorems were quoted by P. Lévy [3] without proof. They give a definitive solution to the problem of uniform continuity of Brownian motion $X(t)$ and are comparable to A. Kolmogorov's criterion in the theory of iterated logarithm for $X(t)$ at time point ∞ .

Theorem 2 is a simple corollary of Theorem 1. Hence we prove only Theorem 1.

LEMMA 1. Without loss of generality, we may assume that

$$(4) \quad (2 \log t - 10 \log \log t)^{\frac{1}{2}} \leq \psi(t) \leq (2 \log t + 10 \log \log t)^{\frac{1}{2}}.$$

PROOF. We show that if Theorem 1 holds under the assumption (4), then it holds without (4). Let us denote the first member in (4) by $\psi_1(t)$ and the last member in (4) by $\psi_2(t)$.

Define $\hat{\psi}(t)$ as follows:

$$(5) \quad \hat{\psi}(t) = \min(\max(\psi(t), \psi_1(t)), \psi_2(t)).$$

Then the convergence of the integral (2) for $\psi(t)$ implies the same for $\hat{\psi}(t)$. In fact, let us assume the convergence of (2) for $\psi(t)$. If the set of t on which $\psi(t)$ is less than $\psi_1(t)$ is not bounded, there exists an increasing sequence $\{t_n\}$ such that $\psi(t_n) \leq \psi_1(t_n)$ and t_n tends to infinity with n . Since $\psi(t)$ is a non-negative and non-decreasing function, we have

$$\begin{aligned} \int_{t_1}^{\infty} \psi^3(t) e^{-\frac{1}{2}\psi^2(t)} dt &\geq \int_{t_1}^{t_n} \psi^3(t) e^{-\frac{1}{2}\psi^2(t)} dt \\ &\geq c \psi^3(t_n) e^{-\frac{1}{2}\psi^2(t_n)} t_n \\ &\geq c (\log t_n)^{\frac{13}{2}} \end{aligned}$$

where c is a positive constant. Since $\log t_n$ tends to infinity with n , the integral for $\psi(t)$ is divergent. This contradicts our assumption and therefore $\psi_1(t)$ must be smaller than $\psi(t)$ for large t . On the other hand the integral for $\psi_2(t)$ is convergent. These facts prove our assertion. Now we assume that the integral for $\psi(t)$ is convergent and Theorem 1 valid under the condition (4). Then the integral for $\hat{\psi}(t)$ is convergent and therefore $\hat{\psi}(t)$ belongs to the upper class. But by what has just been shown $\hat{\psi}(t) \leq \psi(t)$ for large t . So we have $\hat{\varphi}(h) \leq \varphi(h)$ for small h where $\hat{\varphi}(t)$ is defined by $\hat{\psi}(t)$ as $\varphi(t)$ is by $\psi(t)$ and therefore $\psi(t)$ belongs to the upper class. Thus Lemma 1 is proved in the convergent case.

Secondly let us assume that the integral for $\psi(t)$ is divergent. If the set of t on which $\psi(t)$ is less than $\psi_1(t)$ is bounded, then it follows that $\hat{\psi}(t)$ is less than $\psi(t)$ for large t and accordingly the integral for $\hat{\psi}(t)$ must be divergent. On the contrary, if there exists an increasing sequence $\{t_n\}$ having the property

$$(6) \quad \psi(t_n) < \psi_1(t_n), \quad t_n \rightarrow \infty \quad \text{as } n \rightarrow \infty,$$

then we have

$$(7) \quad \hat{\psi}(t_n) = \psi_1(t_n).$$

By the monotony of $\hat{\psi}(t)$, we have

$$(8) \quad \int_{t_1}^{t_n} \hat{\psi}^3(t) e^{-\frac{1}{2}\hat{\psi}^2(t)} dt \geq \psi^3(t_n) e^{-\frac{1}{2}\hat{\psi}^2(t_n)}(t_n - t_1) \\ = \psi_1^3(t_n) e^{-\frac{1}{2}\psi_1^2(t_n)}(t_n - t_1).$$

Since the last term in (8) tends to infinity with n , the integral for $\hat{\psi}(t)$ is divergent in our case. Now, by the result in [2], $\psi_2(t)$ belongs to the upper class and therefore, for almost all sample point ω , there exists ε such that

$$(9) \quad |X(t', \omega) - X(t, \omega)| < \varphi_2(|t' - t|) \quad \text{for } |t' - t| < \varepsilon,$$

where $\varphi_2(t)$ is defined by $\psi_2(t)$ in the same way as $\varphi(t)$ is by $\psi(t)$. On the other hand, since by assumption $\psi_2(t)$ belongs to the lower class, for almost all ω we can choose a sequence $\{(t_n, t_n')\}$ having the following properties

$$(10) \quad |X(t_n') - X(t_n)| > \hat{\varphi}(|t_n' - t_n|), \\ |t_n' - t_n| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

From (9) and (10), we have

$$(11) \quad \hat{\varphi}(|t_n' - t_n|) < \varphi_2(|t_n' - t_n|).$$

(11) shows that $\varphi(t)$ is at last equal to $\varphi(t)$ at $t = |t_n' - t_n|$. This fact and (10) show that $\psi(t)$ belongs to the lower class. Q. E. D.

We now proceed to prove Theorem 1.

1) Proof of the convergent case.

First of all we remark that it suffices to prove, for almost all ω , the existence of a positive ε' such that

$$X(t', \omega) - X(t, \omega) \leq \varphi(|t' - t|) \quad \text{for } |t' - t| < \varepsilon'.$$

In fact, let us assume that this assertion holds. Then it follows from the symmetry of Brownian motion that the probability of the existence of a positive ε'' satisfying the inequality

$$-\varphi(|t' - t|) \leq X(t', \omega) - X(t, \omega) \quad \text{for } |t' - t| < \varepsilon''$$

is equal to 1. Taking ε for the minimum of ε' and ε'' , we have Theorem 1. Therefore we may consider the difference $X(t') - X(t)$ instead of its absolute value.

For each triple (p, k, l) , let $E_{k,l}^p$ be the event

$$(12) \quad X\left(\frac{k+l}{2^p}\right) - X\left(\frac{k}{2^p}\right) \geq \varphi\left(\frac{l}{2^p}\right), \quad k = 0, 1, 2, \dots, 2^p, \\ l = 1, 2, \dots, p.$$

A simple computation shows that

$$P(E_{k,l}^p) \sim e^{-\frac{1}{2}\psi^2\left(\frac{2^p}{l}\right)} / (2\pi)^{\frac{1}{2}} \psi\left(\frac{2^p}{l}\right)$$

for large p . Summing up $P(E_{k,l}^p)$ for $p = 1, 2, \dots$, $k = 1, 2, \dots, 2^p$, $l = \left[\frac{p}{3}\right], \left[\frac{p}{3}\right] + 1, \dots, p$, we have

$$\sum_{p=1}^{\infty} \sum_{k=1}^{2^p} \sum_{l=\left[\frac{p}{3}\right]}^p P(E_{k,l}^p) = O(1) \sum_{p=1}^{\infty} \sum_{k=1}^{2^p} \sum_{l=\left[\frac{p}{3}\right]}^p e^{-\frac{1}{2}\psi^2\left(\frac{2^p}{l}\right)} / \psi\left(\frac{2^p}{l}\right) \\ = O(1) \sum_{p=1}^{\infty} \sum_{k=1}^{2^p} \frac{p}{\psi\left(\frac{2^p}{p}\right)} e^{-\frac{1}{2}\psi^2\left(\frac{2^p}{p}\right)}.$$

Applying Lemma 1, we obtain

$$(13) \quad \sum_{p=1}^{\infty} \sum_{k=1}^{2^p} \sum_{l=\left[\frac{p}{3}\right]}^p P(E_{k,l}^p) = O(1) \sum_{p=1}^{\infty} \frac{2^p}{p} \psi^3\left(\frac{2^p}{p}\right) e^{-\frac{1}{2}\psi^2\left(\frac{2^p}{p}\right)} \\ = O(1) \int_0^{\infty} \psi^3(t) e^{-\frac{1}{2}\psi^2(t)} dt < +\infty.$$

Next, for each triple (p, k, l) , let $F_{k,l}^p$ be the event

$$(14) \quad \max_{0 \leq t, s \leq \frac{1}{2^p}} \left\{ X\left(\frac{k+l}{2^p} + t\right) - X\left(\frac{k}{2^p} - s\right) \right\} \geq \sqrt{\frac{l}{2^p}} \psi\left(\frac{2^p}{l+2}\right), \\ k = 0, 1, 2, \dots, 2^p, \\ l = 1, 2, \dots, p.$$

For convenience' sake, we consider the $F_{k,l}^p$ only such that the time parameters t of $X(t)$ which appear in the above definition are positive and less than 1. It is well known that

$$P(\max_{0 \leq s \leq t} X(s) > a) \leq 2P(X(t) > a),$$

where a is any real number. Since the Brownian motion is an additive process, we have

$$\begin{aligned}
 P(F_{k,l}^p) &\leq P \left\{ \max_{0 \leq t \leq \frac{1}{2^p}} \left(X\left(\frac{k+l}{2^p} + t\right) - X\left(\frac{k+l}{2^p}\right) \right) + \left(X\left(\frac{k+l}{2^p}\right) - X\left(\frac{k}{2^p}\right) \right) \right. \\
 &\quad \left. + \max_{0 \leq s \leq \frac{1}{2^p}} \left(X\left(\frac{k}{2^p}\right) - X\left(\frac{k}{2^p} - s\right) \right) \geq \sqrt{\frac{l}{2^p}} \psi\left(\frac{2^p}{l+2}\right) \right\} \\
 (15) \quad &\leq 4P \left\{ X\left(\frac{k+l+1}{2^p}\right) - X\left(\frac{k+l}{2^p}\right) + X\left(\frac{k+l}{2^p}\right) - X\left(\frac{k}{2^p}\right) \right. \\
 &\quad \left. + X\left(\frac{k}{2^p}\right) - X\left(\frac{k-1}{2^p}\right) \geq \sqrt{\frac{l}{2^p}} \psi\left(\frac{2^p}{l+2}\right) \right\} \\
 &= 4P \left(X\left(\frac{k+l+1}{2^p}\right) - X\left(\frac{k-1}{2^p}\right) \geq \sqrt{\frac{l}{2^p}} \psi\left(\frac{2^p}{l+2}\right) \right)
 \end{aligned}$$

By Lemma 1 we have, for large p and l ,

$$\begin{aligned}
 P(F_{k,l}^p) &\leq \frac{4}{(2\pi)^{\frac{1}{2}} \psi\left(\frac{2^p}{l+2}\right)} e^{-\frac{l}{2(l+2)} \psi^2\left(\frac{2^p}{l+2}\right)} \\
 &\sim 4P(E_{k,l}^p) e^{\frac{1}{l+2} \psi^2\left(\frac{2^p}{l+2}\right)}.
 \end{aligned}$$

Therefore, if l is an integer existing between $\left[\frac{p}{3}\right]$ and p , there exists a positive constant c such that

$$(16) \quad P(F_{k,l}^p) \leq cP(E_{k,l}^p).$$

Combining (13) and (16), we obtain

$$(17) \quad \sum_{p=1}^{\infty} \sum_{k=1}^{2^p} \sum_{l=\left[\frac{p}{3}\right]}^p P(F_{k,l}^p) < +\infty.$$

According to Borel-Cantelli's lemma in the convergent case, (17) shows that the events $F_{k,l}^p$ appearing in (17) occur "only finitely many times" with probability 1. Or, in other words, there exists a positive ϵ with probability 1 such that if $\frac{p}{2^{p+1}}$ is smaller than ϵ , $F_{k,l}^p$ does not occur for any pair (k, l) appearing in the summation of (17).

Now, for any pair of (t, t') satisfying the condition $|t' - t| < \epsilon$, we choose p as follows:

$$(18) \quad \frac{p+1}{2^{p+1}} < |t' - t| \leq \frac{p}{2^p} < 2\epsilon.$$

If we define k and l by the following inequalities

$$(19) \quad \frac{k-1}{2^p} < \min(t, t') \leq \frac{k}{2^p} < \frac{k+l}{2^p} \leq \max(t, t') < \frac{k+l+1}{2^p},$$

it follows that $\left[\frac{p}{3} \right] < l \leq p$ and therefore we obtain

$$\begin{aligned} X(t') - X(t) &\leq \max_{0 \leq t, s \leq \frac{1}{2^p}} \left(X\left(\frac{k+l}{2^p} + t\right) - X\left(\frac{k}{2^p} - s\right) \right) \\ &\leq \left(\frac{l}{2^p}\right)^{\frac{1}{2}} \psi\left(\frac{2^p}{l+2}\right) \\ &\leq \varphi(|t' - t|) \end{aligned}$$

with probability 1.

Thus Theorem 1 is proved in the convergent case.

2) Proof of the divergent case.

Let $E_{k,t}^p$ be the event defined by (12). By the monotony of $\psi(t)$ and Lemma 1, we have

$$\begin{aligned} \sum_{p=1}^{\infty} \sum_{k=1}^{2^p} \sum_{l=\left[\frac{p}{2}\right]+1}^p P(E_{k,t}^p) &= O(1) \sum_{p=1}^{\infty} \sum_{k=1}^{2^p} \sum_{l=\left[\frac{p}{2}\right]+1}^p e^{-\frac{1}{2}\psi^2\left(\frac{2^p}{l}\right)} / \psi\left(\frac{2^p}{l}\right) \\ (20) \qquad &= O(1) \sum_{p=1}^{\infty} \frac{2^p}{p} \psi^3\left(\frac{2^{p+1}}{p}\right) e^{-\frac{1}{2}\psi^2\left(\frac{2^{p+1}}{p}\right)} \\ &= O(1) \int_0^{\infty} \psi^3(t) e^{-\frac{1}{2}\psi^2(t)} dt = +\infty. \end{aligned}$$

It is sufficient to show that $E_{k,t}^p$ occur “infinitely often” with probability 1. For this purpose, we use the following Lemma given in [4].

LEMMA 2. Let $\{E_k\}$ be a sequence of events satisfying the following conditions.

(i)
$$\sum_{k=1}^{\infty} P(E_k) = +\infty.$$

(ii) For every pair of positive integers h, n with $n \geq h$, there exist $c(h)$ and $H(n, h) > n$ such that for every $m \geq H(n, h)$ we have

$$P(E_m / E_k', \dots, E_n') > c(h)P(E_m),$$

where $P(F/E)$ denotes the conditional probability of F on the hypothesis E and E' denotes the complement of E .

(iii) There exist two absolute constants c_1 and c_2 with the following property: to each E_j there corresponds a set of events E_{j_1}, \dots, E_{j_s} belonging to $\{E_k\}$ such that

(a)
$$\sum_{i=1}^s P(E_j E_{j_i}) < c_1 P(E_{j_i})$$

and if $k > j$ but E_k is not among the E_{j_i} ($1 \leq i \leq s$) then

(b)
$$P(E_j E_k) < c_2 P(E_j) P(E_k).$$

Then the probability that the events E_k occur "infinitely often" is equal to one.

We rearrange $E_{k,t}$ and denotes it by E_m so that we may apply Lemma 2 in our case. The rule of ordering is given by the following. If $E_n = E_{k,t}$, $E_m = E_{k',t'}$, then $n < m$ if and only if one of the following three conditions holds :

- (α) $p < p'$,
- (β) $p = p'$ and $l > l'$,
- (γ) $p = p'$, $l = l'$ and $k < k'$.

Now we prove that the sequence $\{E_n\}$ satisfies the conditions of Lemma 2. (i) is a consequence of (20). For (ii), we use the characteristic property of Gaussian distribution. Let $E_m = E_{k,t}$ and put $U_m = X\left(\frac{k+l}{2^q}\right) - X\left(\frac{k}{2^p}\right)$. For every pair (h, n) with $n \geq h$, if we define U_h, U_{h+1}, \dots, U_n similarly then

$$(21) \quad \begin{aligned} E(U_i) &= 0 \quad (i = h, h+1, \dots, n), & E(U_m) &= 0, \\ E(U_i U_m) &\leq \frac{l}{2^p} \quad (i = h, h+1, \dots, n), \end{aligned}$$

where $E(U)$ denotes the expectation of U . Since $\frac{l}{2^p}$ tends to zero as p increases, (21) shows that for each i ($h \leq i \leq n$) the correlation coefficient of U_i and U_m tends to zero as m increases. In other words, U_m is asymptotically independent of the joint variable $(U_h, U_{h+1}, \dots, U_n)$. Therefore we have

$$(22) \quad \lim_{m \rightarrow \infty} \frac{P(E_m/E_h', \dots, E_n')}{P(E_m)} = \lim_{m \rightarrow \infty} \frac{P(E_h', E_{h+1}', \dots, E_n'/E_m)}{P(E_h', E_{h+1}', \dots, E_n')} = 1.$$

This shows that (ii) holds in our case. For the justification of (iii), we need some lemmas.

LEMMA 3. Let U and V be two random variables whose joint distribution is Gaussian and each of them has a standard Gaussian distribution. Let the correlation coefficient of U and V be ρ , then there exists a positive constant c_1 such that

$$(23) \quad P(U > a, V > b) \leq c_1 P(U > a)P(V > b) \quad \text{for } \rho < \frac{1}{ab}.$$

PROOF. If ρ is negative or if a or b is small, (23) holds trivially. Therefore it is sufficient to prove Lemma 3 for sufficiently large a, b and positive ρ . Without loss of generality, we may assume $a \leq b$. Then we have

$$P(U > a, V > b) = \frac{1}{2\pi(1-\rho^2)^{\frac{1}{2}}} \int_b^\infty \int_a^\infty e^{-\frac{(x^2-2\rho xy+y^2)}{2(1-\rho^2)}} dx dy$$

$$\begin{aligned}
 (24) \quad &= \frac{1}{2\pi(1-\rho^2)^{\frac{1}{2}}} \int_b^{2b} \int_a^{2b} e^{-\frac{(x-\rho y)^2}{2(1-\rho^2)}} e^{-\frac{y^2}{2}} dx dy \\
 &+ \frac{1}{2\pi(1-\rho^2)^{\frac{1}{2}}} \int_b^\infty \int_{2b}^\infty e^{-\frac{(x-\rho y)^2}{2(1-\rho^2)}} e^{-\frac{y^2}{2}} dx dy \\
 &+ \frac{1}{2\pi(1-\rho^2)^{\frac{1}{2}}} \int_{2b}^\infty \int_a^{2b} e^{-\frac{(x-\rho y)^2}{2(1-\rho^2)}} e^{-\frac{y^2}{2}} dx dy.
 \end{aligned}$$

The first term on the right side is estimated as follows:

$$\begin{aligned}
 (25) \quad &\frac{1}{2\pi(1-\rho^2)^{\frac{1}{2}}} \int_b^{2b} \int_a^{2b} e^{-\frac{(x-\rho y)^2}{2(1-\rho^2)}} e^{-\frac{y^2}{2}} dx dy \\
 &\leq \frac{1}{2\pi(1-\rho^2)^{\frac{1}{2}}} \int_b^{2b} \int_a^{2b} e^{-\frac{(x-2/a)^2}{2(1-\rho^2)}} e^{-\frac{y^2}{2}} dx dy \\
 &\leq P\{U > (a-2/a)/(1-\rho^2)^{\frac{1}{2}}\} P(V > b) \\
 &= O(1)P(U > a)P(V > b).
 \end{aligned}$$

On the other hand, for sufficiently large a , the second and third term on the right side of (24) are trivially smaller than the right side of (23) replaced c_1 by 1. These estimates assure the validity of Lemma 3. Q. E. D.

LEMMA 4. Let U and V be random variables as in Lemma 3. If the correlation coefficient of U and V is less than $1/2^{\frac{1}{2}}$ and $0 < a < b$ then there exist two positive constants c_2 and δ_2 satisfying the following inequality

$$(26) \quad P(U > a, V > b) \leq c_2 e^{-\delta_2 b^2} P(U > a).$$

PROOF. Let ε be a positive constant which is less than 1 and let ρ be the correlation coefficient of U and V . It suffices to prove Lemma 4 for sufficiently large a and positive ρ . Then we have

$$\begin{aligned}
 (27) \quad P(U > a, V > b) &= \frac{1}{2\pi(1-\rho^2)^{\frac{1}{2}}} \int_a^\infty \int_b^\infty e^{-\frac{(x^2-2\rho xy+y^2)}{2(1-\rho^2)}} dx dy \\
 &= \frac{1}{2\pi(1-\rho^2)^{\frac{1}{2}}} \int_a^{(1+\varepsilon)^{\frac{1}{2}}b} \int_b^\infty e^{-\frac{(x-\rho y)^2}{2(1-\rho^2)}} e^{-\frac{y^2}{2}} dx dy \\
 &+ \frac{1}{2\pi(1-\rho^2)^{\frac{1}{2}}} \int_{(1+\varepsilon)^{\frac{1}{2}}b}^\infty \int_b^\infty e^{-\frac{(x-\rho y)^2}{2(1-\rho^2)}} e^{-\frac{y^2}{2}} dx dy
 \end{aligned}$$

$$\begin{aligned} &\leq \frac{1}{2\pi(1-\rho^2)^{\frac{1}{2}}} \int_a^{(1+\varepsilon)\frac{1}{2}b} \int_b^\infty e^{-\frac{(x-(1+\varepsilon/2)\frac{1}{2}b)^2}{2(1-\rho^2)}} e^{-\frac{y^2}{2}} dx dy \\ &\quad + \frac{1}{(2\pi)^{\frac{1}{2}}} \int_{(1+\varepsilon)\frac{1}{2}b}^\infty e^{-\frac{y^2}{2}} dy \\ &= O(1)\{e^{-(1-(1+\varepsilon/2)\frac{1}{2})^2 b^2/2} + e^{-\varepsilon/2 b^2}\} P(U > a). \end{aligned}$$

If we take the minimum of $(1-(1+\varepsilon/2)\frac{1}{2})^2/2$ and $\varepsilon/2$ for δ_2 then Lemma 4 follows from (27) immediately. Q. E. D.

LEMMA 5. Let U and V be random variables as in Lemma 3. Denoting the correlation coefficient of U and V by ρ , there exist two positive constants c_3 and δ_3 such that

$$(28) \quad P(U > a, V > a) \leq c_3 e^{-\delta_3(1-\rho^2)a^2} P(U > a) \quad \text{for } a > 0.$$

PROOF. By the definition of Gaussian distribution, we have

$$P(U > a, V > a) = \frac{1}{2\pi(1-\rho^2)^{\frac{1}{2}}} \int_a^\infty \int_a^\infty e^{-\frac{(x^2-2\rho xy+y^2)}{2(1-\rho^2)}} dy dx.$$

Rotating the axes by $\pi/4$, we obtain

$$\begin{aligned} P(U > a, V > a) &= \frac{1}{2\pi(1-\rho^2)^{\frac{1}{2}}} \int_{\frac{1}{2}a}^\infty \int_{-\frac{1}{2}a}^{\frac{1}{2}a} e^{-\frac{(1-\rho)x^2+(1+\rho)y^2}{2(1-\rho^2)}} dy dx \\ &\leq \frac{1}{(2\pi)^{\frac{1}{2}}} \int_{(2/1+\rho)\frac{1}{2}a}^\infty e^{-\frac{x^2}{2}} dx \\ &= O(1) e^{-\frac{1-\rho^2}{2(1+\rho)^2} a^2} \cdot \frac{1}{a} e^{-\frac{a^2}{2}} \\ &= O(1) e^{-\frac{1-\rho^2}{2(1+\rho)^2} a^2} P(U > a). \end{aligned}$$

If we take $1/8$ for δ_3 , Lemma 5 follows from (31). Q. E. D.

Now we prove that the condition (iii) of Lemma 2 is satisfied by our sequence $\{E_n\}$. For given E_j , recalling that E_j has another expression $E_{k',l}$, we choose a sequence $\{E_{j_i}; i=1, 2, \dots, s\}$ of events with the properties that $j_i > j$, the corresponding superscript p' is less than $(p+5 \log p)$ and E_{j_i} is not independent of E_j . If E_m is independent of E_j then (b) of (ii) holds trivially for $c_2=1$. On the other hand, if E_m is not independent of E_j , we use Lemma 3. Let $E_j = E_{k',l}$ and $E_m = E_{k'',l'}$. If m is not one of the j_i 's then it follows from the definition of $\{E_{j_i}\}$ that $(p+5 \log p) < p'$. Considering only the case of $l > \frac{p}{2}$, we have by Lemma 1 and for large p ,

$$(30) \quad E \left\{ \frac{\left(X\left(\frac{k+l}{2^p}\right) - X\left(\frac{k}{2^p}\right) \right)}{\left(\frac{l}{2^p}\right)^{\frac{1}{2}}} \cdot \frac{\left(X\left(\frac{k'+l'}{2^{p'}}\right) - X\left(\frac{k'}{2^{p'}}\right) \right)}{\left(\frac{l'}{2^{p'}}\right)^{\frac{1}{2}}} \right\} \leq \left(\frac{p'}{2^{p'}}\right)^{\frac{1}{2}} \left(\frac{2^{p+1}}{p}\right)^{\frac{1}{2}} \\ \leq \frac{1}{\psi\left(\frac{2^p}{p}\right)\psi\left(\frac{2^{p'}}{p'}\right)}.$$

Since the joint distribution of the two random variables appearing in (30) is a Gaussian distribution in 2-dimension's, we may use Lemma 3. Thus there exists a positive constant c such that

$$(31) \quad P(E_j E_m) = P\left\{ X\left(\frac{k+l}{2^p}\right) - X\left(\frac{k}{2^p}\right) > \varphi\left(\frac{l}{2^p}\right), X\left(\frac{k'+l'}{2^{p'}}\right) - X\left(\frac{k'}{2^{p'}}\right) > \varphi\left(\frac{l'}{2^{p'}}\right) \right\} \\ \leq c P(E_j) P(E_m).$$

If we take the maximum of c and 1 for c_2 in (b) of (iii) then (b) holds.

In order to verify (a) of (iii), we use the other expressions of the E_j 's. Let us denote E_j by $E_{k,l}$ and each one of E_{j_i} by E_{k_i, l_i} . Dividing the sum of $P(E_j E_{j_i})$ according to the magnitude of the correlation coefficient of $\left(X\left(\frac{k+l}{2^p}\right) - X\left(\frac{k}{2^p}\right)\right)$ and $\left(X\left(\frac{k'+l'}{2^{p'}}\right) - X\left(\frac{k'}{2^{p'}}\right)\right)$ we have

$$(32) \quad \sum_{i=1}^s P(E_j E_{j_i}) = \Sigma' P(E_j E_{j_i}) + \Sigma'' P(E_j E_{j_i}),$$

where Σ' denotes the summation over i 's such that the correlation coefficient of the corresponding random variables is larger than $\frac{1}{\sqrt{2}}$ and Σ'' denotes the summation of the remainder. Since the correlation is at most

$$\min\left(\frac{l}{2^p}, \frac{l'}{2^{p'}}\right) \left(\frac{ll'}{2^{p+p'}}\right)^{-1/2}$$

and since $l'2^{-p'} \leq l2^{-p}$ by the limitation on the ranges of l and l' , we see that the largest superscript of E_{j_i} 's appearing in Σ' is at most $p+2$. Moreover, without loss of generality, we may assume in the computation of $P(E_j E_{j_i})$ that $\frac{k}{2^p} \leq \frac{k'}{2^{p'}}$. If $\frac{k+l}{2^p} \leq \frac{k'+l'}{2^{p'}}$, we have

$$(33) \quad P(E_j E_{j_i}) = P\left(\frac{X\left(\frac{k+l}{2^p}\right) - X\left(\frac{k}{2^p}\right)}{\left(\frac{l}{2^p}\right)^{\frac{1}{2}}} > \psi\left(\frac{2^p}{l}\right), \frac{X\left(\frac{k'+l'}{2^{p'}}\right) - X\left(\frac{k'}{2^{p'}}\right)}{\left(\frac{l'}{2^{p'}}\right)^{\frac{1}{2}}} > \psi\left(\frac{2^{p'}}{l'}\right)\right) \\ \leq P\left(\frac{X\left(\frac{k+l}{2^p}\right) - X\left(\frac{k}{2^p}\right)}{\left(\frac{l}{2^p}\right)^{\frac{1}{2}}} > \psi\left(\frac{2^p}{l}\right), \frac{X\left(\frac{k+l}{2^p}\right) - X\left(\frac{k'}{2^{p'}}\right)}{\left(\frac{k+l}{2^p} - \frac{k'}{2^{p'}}\right)^{\frac{1}{2}}} > \psi\left(\frac{2^p}{l}\right)\right)$$

The inequality follows from the definition of ordering and the fact that the correlation coefficient of two random variables appearing in the last term of (33) is larger than that of the second term. Since $p' \leq p+2$, we obtain by Lemma 1 and Lemma 5 that

$$\begin{aligned}
 P(E_j E_{j_i}) &\leq c e^{-\delta \binom{k'-k2^{p'-p}}{2^{p'-p}} \psi' \left(\frac{2^p}{l} \right)} P(E_j) \\
 &\leq c e^{-\delta' (k'-k2^{p'-p})} P(E_j),
 \end{aligned}
 \tag{34}$$

where c, δ and δ' are positive constants. Here we remark that the number of E_{j_i} appearing in the present case is less than $(k'-k2^{p'-p})$ for fixed pair (p', k') because $\frac{l}{2^p} \geq \frac{l'}{2^{p'}}$. Similarly, for the case of $\frac{k}{2^p} > \frac{k'}{2^{p'}}$ we have

$$P(E_j E_{j_i}) \leq c e^{-\delta' (l2^{p'-p}-l')} P(E_j).
 \tag{35}$$

Considering the same situation for $\frac{k}{2^p} > \frac{k'}{2^{p'}}$, we have

$$\begin{aligned}
 \sum' P(E_j E_{j_i}) &\leq 2c P(E_j) \sum_{p'-p}^{p+2} \left\{ \sum_{k'=k2^{p'-p}}^{(k+l)2^{p'-p}} (k'-k2^{p'-p}) e^{-\delta' (k'-k2^{p'-p})} \right. \\
 &\quad \left. + \sum_{l'=1}^{l2^{p'-p}} (l2^{p'-p}-l') e^{-\delta' (l2^{p'-p}-l')} \right\} \\
 &\leq \alpha P(E_j),
 \end{aligned}
 \tag{36}$$

where α is an absolute constant.

For the computation of $P(E_j E_{j_i})$ where E_{j_i} appears in the summation of \sum'' , we apply Lemma 4. Using the same expression for E_i and E_{j_i} as before, for the case of $\frac{k}{2^p} \leq \frac{k'}{2^{p'}} < \frac{k+l}{2^p} \leq \frac{k'+l'}{2^{p'}}$, we have

$$\begin{aligned}
 P(E_j E_{j_i}) &\leq P \left(\frac{X \left(\frac{k+l}{2^p} \right) - X \left(\frac{k}{2^p} \right)}{\left(\frac{l}{2^p} \right)^{\frac{1}{2}}} > \psi \left(\frac{2^p}{l} \right), \frac{X \left(\frac{k+l}{2^p} \right) - X \left(\frac{k'}{2^{p'}} \right)}{\left(\frac{k+l}{2^p} - \frac{k'}{2^{p'}} \right)^{\frac{1}{2}}} > \psi \left(\frac{2^{p'}}{l'} \right) \right) \\
 &\leq c e^{-\delta \psi' \left(\frac{2^{p'}}{l'} \right)} P(E_j) \\
 &\leq c e^{-\delta' p'} P(E_j)
 \end{aligned}
 \tag{37}$$

where c, c', δ and δ' are positive constants. Similarly, for the case of $\frac{k}{2^p} \leq \frac{k'}{2^{p'}} < \frac{k'+l'}{2^{p'}} \leq \frac{k+l}{2^p}$, we obtain the same result. Combining all the cases, we have

$$\begin{aligned}
 \sum'' P(E_j E_{j'}) &\leq cP(E_j) \sum_{j'=j}^{j+\delta \log p} p'^{1/2} 2^{p'-j} l e^{-\delta' p'} \\
 (38) \qquad \qquad &\leq cP(E_j) \sum_{j'=j}^{j+\delta \log p} p'^{1/2} e^{-\delta' p'} \\
 &= \beta P(E_j),
 \end{aligned}$$

where β is an absolute constant. (32), (36) and (38) establish the validity of (iii a). Therefore we may apply Lemma 2 in our case and Theorem 1 is proved completely.

Q. E. D.

Syracuse University, Technion, Israel,
and Shizuoka University.

References

- [1] P. Lévy, *Théorie de l'addition des variables aléatoires*, 1937.
- [2] T. Sirao, On the uniform continuity of Wiener process, *J. Math. Soc. Japan*, **6**, (1954), 332-335.
- [3] P. Lévy, Le mouvement brownien, *Mém. Sci. Math.*, fasc. 126, 1954.
- [4] K. L. Chung and P. Erdős, On the application of the Borel-Cantelli Lemma, *Trans. Amer. Math. Soc.*, **72**, (1952), 179-189.