

## CONCERNING PERIODICITY IN THE ASYMPTOTIC BEHAVIOUR OF PARTITION FUNCTIONS

Dedicated to George Szekeres on his 65th birthday

P. ERDÖS and B. RICHMOND

(Received 31 December 1974)

Communicated by J. Wallis

### Abstract

Let  $P_A(n)$  denote the number of partitions of  $n$  into summands chosen from the set  $A = \{a_1, a_2, \dots\}$ . De Bruijn has shown that in Mahler's partition problem ( $a_\nu = r^\nu$ ) there is a periodic component in the asymptotic behaviour of  $P_A(n)$ . We show by example that this may happen for sequences that satisfy  $a_\nu \sim \nu$  and consider an analogous phenomena for partitions into primes. We then consider corresponding results for partitions into distinct summands. Finally we obtain some weaker results using elementary methods.

### 1. Introduction

Let  $A = \{a_0, a_1, \dots\}$  be an infinite set of monotone increasing integers. Let  $p_A(n)$  denote the number of ways of representing  $n$  as the sum of summands chosen from  $A$ . Mahler (1940) showed that when  $a_\nu = r^\nu$  as  $n \rightarrow \infty$

$$\log P_A(rn) = \frac{1}{2 \log r} (\log(n(\log n)))^2 + \left( \frac{1}{2} + \frac{1}{\log r} + \frac{\log \log r}{\log r} \right) \log n - \left( 1 + \frac{\log \log r}{\log r} \right) \log \log n + o(1).$$

De Bruijn (1943) has shown that this 0-term is actually of the form

$$U \left( \frac{\log n - \log \log n + \log \log r}{\log r} \right) + o \left\{ \frac{(\log \log n)^2}{\log n} \right\}$$

where  $U$  is a periodic function with period one and de Bruijn determined the Fourier expansion of  $U$ . This result has been generalized by various authors, for example, Pennington (1953) and Schwarz (1967) and it seems to be common with sets  $A$  which have  $\liminf (\log a_\nu)/\nu > 0$  for there to be a periodic or almost

periodic function analogous to de Bruijn's  $U$ . In this paper we consider the question of whether partition functions  $p_A(n)$  with  $\liminf_{\nu \rightarrow \infty} (\log a_\nu)/\nu = 0$  may also exhibit the de Bruijn–Mahler phenomenon. We first show that  $p_A(n)$  with  $A$  defined to be the set of integers which are not positive powers of a fixed integer has a periodic term very similar to that of de Bruijn (see theorem 1). We then investigate how large this term corresponding to de Bruijn's  $U$  function may be. It is seen that when  $A$  is the set of primes it may be large indeed and we obtain a direct connection with the Riemann hypothesis. In this case there is a close analogy with the number of primes less than a given limit.

Finally we consider these questions for  $q_A(n)$  the number of partitions into distinct summands chosen from the set  $A$  and obtain some results using elementary methods.

## 2

Let  $A$  be the set of integers which are not powers of the fixed integer  $r$ . Then the number  $p_A(n)$  of partitions of  $n$  into summands from  $A$  satisfied (from the theorem of Roth and Szekeres (1954))

$$(2.1) \quad p_A(n) = (2\pi A_2)^{-\frac{1}{2}} \exp \left\{ \alpha n - \sum_{a \in A} \log(1 - e^{-\alpha a}) \right\} [1 + O\{n^{-\frac{1}{2}}\}]$$

where

$$(2.2) \quad A_2 = \sum_{a \in A} a^2 \frac{e^{-\alpha a}}{(e^{-\alpha a} - 1)^2}$$

and  $\alpha$  is defined by

$$(2.3) \quad n = \sum_{a \in A} \frac{a}{e^{\alpha a} - 1}.$$

From Mellin's transformation formula

$$\sum_{a \in A} \frac{a}{e^{\alpha a} - 1} = \frac{1}{2\pi i} \int_{\sigma - i\infty}^{\sigma + i\infty} \alpha^{-t} \Gamma(t) \zeta(t) \zeta_A(t-1) dt \quad (\sigma > 2)$$

where  $\Gamma(t)$  denotes the usual gamma-function,  $\zeta(t)$  denotes the Riemann zeta-function and

$$\zeta_A(t) = \sum_{a \in A} a^{-t}.$$

Now

$$\zeta_A(t) = \sum_{n=1}^{\infty} n^{-t} - \sum_{j=1}^{\infty} r^{-jt} = \zeta(t) - \frac{1}{r^t - 1},$$

thus  $\zeta_A(t)$  is defined in the entire plane except for simple poles at  $t = 1$  and  $t = 2\pi ik/\log r$  ( $k = 0, 1, 2, \dots$ ). The arguments of Pennington (1953) show that we may shift the contour of integration to a line  $\sigma < 0$  and obtain (using  $\zeta(0) = -\frac{1}{2}$ ) that

$$(2.4) \quad n = \alpha^{-2} \pi^2/6 - \alpha^{-1} \frac{\log(1/\alpha)}{\log r} - \frac{1}{\alpha \log r} \sum'_\nu \Gamma\left(1 + \frac{2\pi i\nu}{\log r}\right) \zeta\left(1 + \frac{2\pi i\nu}{\log r}\right) \exp\left(-\frac{2\pi i\nu \log \alpha}{\log r}\right) + O\{1\}$$

where  $\gamma$  denotes Euler's constant and  $\sum'_\nu$  denotes summation over nonzero  $\nu$ . We may solve this for  $\alpha$  to obtain

$$(2.5) \quad \alpha = n^{-1/2} \frac{\pi}{\sqrt{6}} \left[ 1 - n^{-1/2} \frac{\sqrt{6}}{2\pi \log r} \left\{ \log \frac{\sqrt{6}}{\pi} + \sum'_\nu \Gamma\left(1 + \frac{2\pi i\nu}{\log r}\right) \zeta\left(1 + \frac{2\pi i\nu}{\log r}\right) \exp\left(2\pi i\nu \log\left(\frac{\sqrt{6}n}{\pi}\right)\right) \right\} \right] + O\{n^{-2/3} \log^2 n\}.$$

From Mellin's transformation formula

$$-\sum_{a \in A} \log(1 - e^{-\alpha a}) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \alpha^{-t} \zeta(t+1) \Gamma(t) \zeta_A(t) dt \quad (\sigma > 1).$$

Again shifting the contour of integration to  $\sigma < 0$ , using  $\zeta'(0) = \frac{1}{2} \log 2\pi$  and that (with  $z\zeta(z+1) = 1 + \gamma z + \gamma_2 z^2 + \dots$ )

$$\zeta(t+1) \Gamma(t) \zeta_A(t) = \frac{-1}{t^3 \log r} - \left\{ \frac{\log^2 r}{12} - \frac{1}{2} \log 2\pi + \frac{\gamma_2 + \frac{\gamma^2}{2} + \frac{\pi^2}{12}}{\log r} \right\} \frac{1}{t} + O\{1\}$$

yields

$$(2.6) \quad -\sum_{a \in A} \log(1 - e^{-\alpha a}) = \alpha^{-1} \frac{\pi^2}{6} - \frac{\log^2(1/\alpha)}{\log r} + \frac{\log r}{12} + \frac{1}{2} \log 2\pi - \frac{\gamma_2 + \frac{\gamma^2}{2} + \frac{\pi^2}{12}}{\log r} - \frac{1}{\log r} \sum'_\nu \zeta\left(1 + \frac{2\pi i\nu}{\log r}\right) \Gamma\left(\frac{2\pi i\nu}{\log r}\right) \times \exp\left(\frac{2\pi i\nu \log \alpha}{\log r}\right) + O\{\alpha\}.$$

From (2.4) and (2.6) we obtain

$$\begin{aligned}
 \alpha n - \sum_{a \in A} \log \{1 - e^{-\alpha a}\} &= \alpha^{-1} \frac{\pi^2}{3} - \frac{\log^2(1/\alpha)}{\log r} - \frac{\log(1/\alpha)}{\log r} + \frac{1}{2} \log 2\pi \\
 (2.7) \quad &- \frac{\log r}{12} - \frac{\gamma_2 + \frac{\gamma^2}{2} + \frac{\pi^2}{12}}{\log r} - \frac{1}{\log r} \sum'_\nu \zeta \left(1 + \frac{2\pi i \nu}{\log r}\right) \\
 &\times \left(1 + \frac{2\pi i \nu}{\log r}\right) \Gamma \left(\frac{2\pi i \nu}{\log r}\right) \exp \left(\frac{2\pi i \nu \log(\alpha^{-1})}{\log r}\right) \\
 &+ O\{\alpha\}.
 \end{aligned}$$

Note that

$$(2.8) \quad A_2 = \frac{d}{\alpha a} \left[ \sum_a \frac{a}{e^{\alpha a} - 1} \right] = \frac{\alpha^{-3} \pi^2}{6} + O\{\alpha^{-2}\}.$$

Thus from (2.7), (2.8) and (2.5) applied to (2.1) we obtain

**THEOREM 1.** *Let  $A$  be the set of integers which are not positive powers of the fixed integer  $r$ . Then*

$$\begin{aligned}
 \log p_A(n) &= \pi n^{\frac{1}{2}} \sqrt{\frac{2}{3}} - \frac{\log^2(\sqrt{6n}/\pi)}{\log r} + 3 \log \left(\frac{\sqrt{6n}}{\pi}\right) \\
 &- \frac{\log r}{12} - \frac{1}{2} \log \left(\frac{\pi^2}{3}\right) - \frac{\gamma_2 + \gamma + \frac{\gamma^2}{2} + \frac{\pi^2}{12}}{\log r} - \frac{1}{\log r} \\
 &\times \sum'_\nu \zeta \left(1 + \frac{2\pi i \nu}{\log r}\right) \Gamma \left(\frac{2\pi i \nu}{\log r}\right) \exp \left(\frac{2\pi i \nu \log \left(\frac{\sqrt{6n}}{\pi}\right)}{\log r}\right) \\
 &+ O\{n^{-1/2} \log^2 n\}.
 \end{aligned}$$

This theorem shows that even sequences for which  $a_\nu/\nu = 1 + O\{\nu^{-1} \log \nu\}$  may exhibit the de Bruijn–Mahler phenomenon. The asymptotic formula is in terms of elementary functions. In our attempt to determine how large this oscillatory component may be we shall consider an example in which the asymptotic formula cannot be so expressed.

**THEOREM 2.** *Let  $A$  be the set of primes. Let  $\alpha = \alpha(n)$  be defined by*

$$n = \sum_{a \in A} \frac{a}{e^{\alpha a} - 1},$$

then

$$\alpha \sim \frac{\pi}{\sqrt{3}} n^{-1/2} \log^{-1/2} n.$$

Let

$$F(\alpha) = \alpha^{-1} \int_0^{\infty} e^{-u \log(1/\alpha)} (2+u) \Gamma(1+u) \zeta(2+u) du.$$

Thus by standard results on the Laplace Transform

$$F(\alpha) = \alpha^{-1} \sum_{r=0}^N \frac{C_r n!}{\log^{r+1} \frac{1}{\alpha}} + O \left\{ \alpha^{-1} \log^{-N+2} \frac{1}{\alpha} \right\}$$

where

$$(2+u) \Gamma(1+u) \zeta(2+u) = \sum_{n=0}^{\infty} C_n u^n, \quad C_0 = \frac{\pi^2}{3}.$$

a) There exists a constant  $C > 0$  such that

$$\log p_A(n) = F(\alpha) + O \left\{ \alpha^{-1} \exp \left( -C \log^{\frac{2}{3}} \left( \frac{1}{\alpha} \right) \log \log^{-\frac{1}{3}} \frac{1}{\alpha} \right) \right\}.$$

b) Let  $\theta \equiv$  l.u.b. of the real parts of the imaginary roots of the Riemann zeta-function. Then for every  $\varepsilon > 0$

$$\log p_A(n) - F(\alpha) = O\{\alpha^{-\theta+\varepsilon}\}.$$

c) Conversely if

$$\log p_A(n) - F(\alpha) = O\{\alpha^{-\psi}\},$$

then  $\theta \leq \psi$  where  $\theta$  is defined in part b).

PROOF. The proof of parts a) and b) is very similar to classical proofs in the theory of primes. It is also very similar to the proof of Theorem 7.1 of Richmond. In particular it is shown in Richmond that

$$\alpha \sim \frac{\pi}{\sqrt{3}} n^{-\frac{1}{2}} \log^{-\frac{1}{2}} n.$$

From the Roth-Szekeres (1954) theorem and the Mellin transformation formula it follows that

$$\log p_A(n) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \alpha^{-t} (1+t) \Gamma(t) \zeta(t+1) \zeta_A(t) dt + O\{\log n\}, \quad (\sigma > 1)$$

where

$$\zeta_A(t) = \sum_{a \in A} a^{-t}.$$

It is well-known that (see p. 12 of Titchmarsh (1951))

$$(2.9) \quad \zeta_A(t) = \log \zeta(t) + h(t), \quad h(t) = \sum_{m=2}^{\infty} \frac{\zeta_A(mt)}{m}$$

where  $h(t)$  is holomorphic for  $Re t > \frac{1}{2}$ . One obtains with standard residue arguments that

$$\begin{aligned} \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \alpha^{-t}(1+t)\Gamma(t)\zeta(t+1) \log \frac{1}{t-1} dt \\ = F(\alpha) + O\left\{ \log \frac{1}{\alpha} \right\}. \end{aligned}$$

There is a constant  $C > 0$  such that  $\zeta(t)$  has no zeros in (p. 87 of Montgomery (1971))

$$Re t = 1 - \frac{C}{\log^{2/3}|t| \log \log^{1/3}|t|}$$

Thus by a classical argument (pp. 77-88 of Prachar (1957) by the  $\Gamma$ -function or see Richmond)

$$\begin{aligned} \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \alpha^{-t}(1+t)\Gamma(t)\zeta(t+1) \left[ \zeta_A(t) - \log \frac{1}{t-1} \right] dt \\ = O\left\{ \alpha^{-1} e^{-C} \log^{\frac{3}{2}}\left(\frac{1}{\alpha}\right) \log \log^{-1/5}\left(\frac{1}{\alpha}\right) \right\}. \end{aligned}$$

This proves part a) of the theorem and part b) follows in the same way.

To prove part c) note that

$$\begin{aligned} \log p_A(n) - F(\alpha) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \alpha^{-t}(1+t)\Gamma(t)\zeta(1+t) \left( \zeta_A(t) - \log \frac{1}{t-1} \right) dt \\ + O\left\{ \log \frac{1}{\alpha} \right\}. \end{aligned}$$

We can think of  $p_A(n)$  as a function of  $\alpha$  and it is known that (see Roth and Szekeres (1954) or Richmond)

$$\alpha(n+1) - \alpha(n) = O\left\{ \alpha^2 \log \frac{1}{\alpha} \right\}$$

$$\log p_A(n+1) - \log p_A(n) = O\left\{ \alpha \log \frac{1}{\alpha} \right\}.$$

From this one may readily deduce (Richmond) that  $F(\alpha(n+1)) - F(\alpha(n)) = O\{\alpha \log 1/\alpha\}$  hence we suppose the hypothesis of c) to hold for all  $\alpha$ .

By the Mellin inversion formula

$$(1+t)\zeta(1+t)\Gamma(t)\left\{\zeta_A(t) - \log \frac{1}{t-1}\right\} = \int_0^\infty \alpha^{t-1} [p_A(n(\alpha)) - F(\alpha)] d\alpha.$$

The integral on the right converges for all  $t$  with  $Rt > \psi$ , hence represents a holomorphic function for  $Rt > \psi$ . By eq. (2.9) we obtain our result.

The proof of part c) of Theorem 2 shows that one cannot have

$$\log p_A(n) - F(\alpha) = O\{\alpha^{1-\theta}\}$$

for any positive constant  $\theta$  since  $\zeta_A(t) - \log(1/(t-1))$  is not bounded at  $t = 1/2$ . Also parts b) and c) show that the size of  $\log p_A(n) - F(\alpha)$  is directly related to the question of where the roots of  $\zeta(t)$  lie.

To see that  $\log p_A(n) - F(\alpha)$  corresponds to the de Bruijn-Mahler phenomenon we consider the following representation:

Let  $\rho = \beta + i\gamma$  run through the complex zeros of  $\zeta(t)$ . Bracket all zeros such that any two zeros for which

$$|\gamma - \gamma'| < \exp(-A\gamma/\log \gamma) + \exp(-A\gamma'/\log \gamma')$$

are included in the same bracket. With the bracketing above by the arguments of Titchmarsh (1951) (on pp. 185-187)

$$\log p_A(n) - F(\alpha) = \sum_{\rho} \text{Res} \left\{ \alpha^{-s} (1+S)\Gamma(S)\zeta(S+1)\zeta_A(S) \right\}_{s=\rho} + O\left\{ \log \frac{1}{\alpha} \right\}.$$

### 3

In this section we consider the corresponding problems for  $q_A(n)$ , the number of partitions of  $n$  into distinct summands chosen from  $A$ . The Roth-Szekeres results and the Mellin transformation techniques in §2 apply with minor modifications and we have

**THEOREM 3.** *Let  $A$  be the set of integers which are not positive powers of a fixed integer  $r$ . Then*

$$\begin{aligned} \log q_A(n) &= \frac{\pi}{\sqrt{3}} n^{1/2} - \frac{\log 2}{\log r} \log \left( \frac{\sqrt{12n}}{\pi} \right) - \frac{3}{4} \log n + \frac{\log^2 s}{2 \log r} - \frac{1}{4} \log(48) \\ &+ \frac{\log 2}{2} - \frac{1}{\log r} \sum_{\nu}' (1 - 2^{-2\pi i \nu / \log r}) \\ &\times \zeta \left( 1 + \frac{2\pi i \nu}{\log r} \right) \exp \left( \frac{2\pi i \nu}{\log r} \log \left( \frac{\sqrt{12n}}{\pi} \right) \right) \Gamma \left( \frac{2\pi i \nu}{\log r} \right) + O\{n^{-1/2} \log^2 n\}. \end{aligned}$$

Note that when  $r = 2$  the oscillatory component drops out and the next theorem shows that this must be expected.

THEOREM 4. Let  $A$  be the set of integers which are not positive powers of 2. Let  $q(n)$  denote the number of partitions of  $n$  into distinct integers. Then

$$q_A(n) = q(n) - q(n-2).$$

PROOF. This theorem follows from generating series techniques however we give the following proof due to H. Shank. We wish to prove that  $q(n-2)$  is the number of partitions of  $n$  into distinct summands at least one of which is a power of 2. Let  $2^i$  be the smallest power of 2 in such a partition of  $n$ . Then  $2^i - 2 = 2 + 2^2 + \dots + 2^{i-1}$  and we obtain a partition of  $n-2$  into distinct summands. Clearly different partitions of  $n$  give different partitions of  $n-2$ . On the other hand suppose we have a partition of  $n-2$  into distinct parts. If it contains a 2 let  $2, 2^2, \dots, 2^i$  be the longest string of consecutive powers of 2 it contains. We may replace this string of  $2^i$ 's by  $2^{i+1}$  to obtain a partition of  $n$ . In this way different partitions of  $n-2$  give rise to different partitions of  $n$  containing a positive power of 2. We now prove

THEOREM 5. Let  $A$  be the set of integers which are not positive powers of a fixed integer  $r$ . Let  $k$  be any constant integer. Then for all sufficiently large  $n$  the  $k$ -th differences of  $p_A(n)$  and  $q_A(n)$  are positive.

PROOF. The result for  $p_A(n)$  follows at once from the work of Bateman and Erdős (1956) since if one removes an arbitrary subset of  $A$  having elements, the remaining elements of  $A$  have greatest common divisor unity.

To prove the result for  $q_A(n)$  we first of all note that it is sufficient to show that  $\sum q_A(n)x^n$  can be written as

$$\sum q_A(n)x^n = \sum b_n x^n \cdot \sum C_n x^n$$

where the  $k$ -th difference of the  $b_n$  is positive for  $n$  sufficiently large and where  $C_n \geq 0$ . Suppose  $r$  has an odd factor  $d$ . We write

$$\begin{aligned} \sum q_A(n)x^n &= \prod_{i=1}^{\infty} (1-x^i) \cdot \prod_{i=1}^{\infty} (1+xr^i)^{-1} \\ &= \prod_{i=0}^{\infty} (1-x^{2i+1})^{-1} \cdot \prod_{i=1}^{\infty} (1+x^{2r^i})^{-1} \cdot \prod_{i=1}^{\infty} (1-x^i) \\ &= \prod_{\substack{j=1(\text{mod } 2) \\ j \neq d^i}}^{\infty} (1-x^j)^{-1} \\ &\quad \times \prod_{i=1}^{\infty} (1+x^{d^i} + \dots + x^{d^i(r/d)^i}) \cdot \prod_{i=1}^{\infty} (1-x^{2r^i})^{-1} \end{aligned}$$



However the  $k$ -difference of the coefficients of the first product are positive from some point on by the results of Bateman and Erdős (1956) and we obtain our result from the remark above. Suppose  $r = 2^s$ . Then

$$\sum q_A(n)x^r = \prod_{j=1}^{\infty} (1 - x^{2^{j+1}})^{-1} \cdot \frac{1 - x^r}{1 - x} \prod_{j=1}^{\infty} \frac{1 - x^{r^{j+1}}}{1 - x^{2r^j}}$$

Clearly  $2r^j/r^{j+1}$  and we again have that the  $k$ -difference of the coefficients of the first product are positive from some point on by the results of Bateman and Erdős (1956).

Finally we show by elementary arguments

**THEOREM 6.** *Let  $A$  be the set of integers which are not positive powers of a fixed integer  $r$ .*

$$\log p_A(n) = \sqrt{\frac{2}{3}} \pi n^{\frac{1}{2}} + O\{\log^2 n\}$$

$$\log q_A(n) = \frac{\pi}{\sqrt{3}} n^{\frac{1}{2}} + O\{\log n\}.$$

**PROOF.** We first prove

$$(3.1) \quad q(n) > q_A(n) > n^{-1}q(n).$$

The first inequality is obvious. Also

$$\prod_{i=1}^{\infty} (1 + x^{r^i}) \cdot \sum q_A(n)x^n = \sum q(n)x^n.$$

Since the coefficients of the Taylor series expansion of the infinite product are zero or one and since the  $q_A(n)$  are monotone increasing we readily obtain the second inequality.

Note that

$$\prod_{i=1}^{\infty} (1 - x^{r^i})^{-1} \cdot \sum p_A(n)x^n = \sum p(n)x^n.$$

The infinite product is  $\sum p_r(n)x^n$  where  $p_r(n)$  is the number of partitions of  $n$  into powers of  $r$ . Erdős (1942) has shown that  $\log p_r(n) \sim \log^2 n / 2 \log r$ . From this and the fact that  $p_A(n)$  and  $p_r(n)$  are monotone increasing we obtain that  $p(n) > p_A(n) \exp - (\varepsilon + (2 \log r)^{-1}) \log^2 n$ . Erdős (1942) has shown using elementary arguments that  $p(n) \sim Cn^{-1} \exp(n^{\frac{1}{2}} \pi \sqrt{2/3})$  (Newman (1951) showed also by elementary arguments that  $C = 1/(4\sqrt{3})$ ). The first part of theorem 6 follows immediately. Since the number of partitions of  $n$  into distinct summands equals the number of partitions of  $n$  into odd summands one can apply the method of Erdős to obtain  $q(n) \sim Cn^{-\frac{1}{2}} \exp(n^{\frac{1}{2}} \pi / \sqrt{3})$  and this with (3.1) gives the second part of Theorem 6.

## References

- K. Mahler (1940), 'On a Special Functional Equation', *J. London Math. Soc.* **15**, 115–123.
- N. G. de Bruijn (1943), 'On Mahler's Partition Problem', *Indag. Math.* **10**, 210–220.
- W. B. Pennington (1953), 'On Mahler's Partition Problem', *Annals of Math.* (2) **57**, 531–546.
- W. Schwarz (1967), 'C. Mahler's Partitions Problem', *J. reine angew. Math* **229**, 182–188.
- K. F. Roth and G. Szekeres (1954), 'Some Asymptotic formulae in the Theory of Partitions', *Quart. J. Math., Oxford* (2), 244–259.
- L. B. Richmond, 'Moments of Partitions II', *Acta Arith.* (to appear).
- E. C. Titchmarsh (1951), *The Theory of the Riemann Zeta-function* (Oxford, 1951).
- H. L. Montgomery (1971), *Topics in Multiplicative Number Theory* (Springer-Verlag, 1971).
- K. Prachar (1957), *Primzahlverteilung* (Springer-Verlag, 1957).
- P. T. Bateman and P. Erdős (1956), 'Monotonicity of Partition Functions', *Mathematica* **3**, 1–14.
- P. Erdős (1943), 'On an Elementary Proof of Some Asymptotic Formulas in the Theory of Partitions', *Annals of Math.* (2) **43**, 437–450.
- D. J. Newman (1951), 'The Evaluation of the Constant in the Formula for the Number of Partitions of  $n$ ', *Amer. J. of Math.*, **73**, 599–601.

University of Waterloo  
University Avenue  
Waterloo  
Ontario, Canada