

DIVISIBILITY THEORY OF SEMI-HEREDITARY RINGS

P. N. ÁNH AND M. SIDDOWAY

(Communicated by Birge Huisgen-Zimmermann)

ABSTRACT. The semigroup of finitely generated ideals partially ordered by inverse inclusion, i.e., the *divisibility theory* of semi-hereditary rings, is precisely described by semi-hereditary Bezout semigroups. A *Bezout semigroup* is a commutative monoid S with 0 such that the divisibility relation $a|b \iff b \in aS$ is a partial order inducing a distributive lattice on S with multiplication distributive on both meets and joins, and for any $a, b, d = a \wedge b \in S$, $a = da_1$ there is $b_1 \in S$ with $a_1 \wedge b_1 = 1$, $b = db_1$. S is *semi-hereditary* if for each $a \in S$ there is $e^2 = e \in S$ with $eS = a^\perp = \{x \in S \mid ax = 0\}$. The dictionary is therefore complete: abelian lattice-ordered groups and semi-hereditary Bezout semigroups describe divisibility of Prüfer (i.e., semi-hereditary) domains and semi-hereditary rings, respectively. The construction of a semi-hereditary Bezout ring with a pre-described semi-hereditary Bezout semigroup is inspired by Stone's representation of Boolean algebras as rings of continuous functions and by Gelfand's and Naimark's analogous representation of commutative C^* -algebras.

INTRODUCTION

In considering the structure of rings, classical number theory suggests a careful study of the *semigroup of divisibility*, defined as the semigroup of principal (or more generally finitely generated) ideals partially ordered by reverse inclusion. We refer to this special semigroup of ideals in a ring R as the *divisibility theory of R* , or more informally as its *divisibility*. This approach led to a general valuation theory developed by Krull, establishing a dictionary between valued fields and totally ordered abelian groups. Later, valuation theory is extended to Bezout domains by showing that the *groups of divisibility* (consisting of fractional ideals) of such domains are precisely lattice-ordered abelian groups. However, the task of describing a divisibility theory for Bezout rings (rings whose finitely generated ideals are principal) with zero-divisors appears to be far more difficult and is still open, although it was quite simple to settle the problem for valuation rings (rings, possibly with zero-divisors, in which any two ideals are comparable).

Received by the editors December 11, 2009 and, in revised form, February 25, 2010.

2000 *Mathematics Subject Classification*. Primary 13A05, 13D05, 13F05; Secondary 06F05.

Key words and phrases. Bezout rings, Stone space, semi-hereditary, rings of continuous functions.

The first author was partially supported by the Hungarian National Foundation for Scientific Research grants no. K61007 and NK72523, Colorado College and UC-Colorado Springs during his stay at Colorado College in the Fall of 2006.

The second author was supported as the Verner Z. Reed Professor of Natural Science at Colorado College from 2007 to the present.

©2010 American Mathematical Society
Reverts to public domain 28 years from publication

In this short paper we provide a stronger but partial answer to this question by showing that for semi-hereditary rings the divisibility theory can be described by way of semi-hereditary Bezout semigroups. As a by-product, this result describes both the divisibility theory of semi-hereditary rings and also of certain sub-direct products of GCD domains, including localizations of polynomial rings in any set of variables over fields. It also provides an overview of the nature of semi-hereditary (Bezout) rings which form a proper subclass between rings with global dimension at most 1 and rings with weak dimension at most 1. Notably, the compactness of the minimal spectrum describes semi-hereditary rings inside rings with weak dimension at most 1. In other words, a ring of weak dimension at most 1 is semi-hereditary if and only if its minimal spectrum is compact. Note that the *divisibility theory in a GCD domain* is a semigroup of principal ideals together with the binary operations GCD and LCM. Therefore semi-hereditary B-semigroups describe divisibility in much larger classes of rings, in particular for semi-hereditary rings.

In the first part we characterize semi-hereditary B-semigroups by their geometric structure, that is, the space of minimal \mathfrak{m} -prime filters endowed with the Zariski topology. The corresponding versions of these results (with one exception) are well-known for semiprime rings [14], [10]. However, the focus of our work is on semigroups, and most of the proofs found in [14] are not direct, are mainly homological in nature, and involve localization techniques. We provide direct elementary proofs in the setting of semigroups which also hold for semiprime rings.

In the second part, the geometry of a semi-hereditary B-semigroup S and its subsemigroup S_1 of non-zero-divisors is used to obtain a semi-hereditary Bezout ring R with a given divisibility theory S . The ring R is a well-determined subring of the ring of continuous functions from the minimal spectrum of S into the quotient field of the semigroup algebra of S_1 endowed with the discrete topology. In the tradition of work by Arens and Kaplansky (see Chapter IX.6 [12]) on the representation of certain biregular rings, our construction provides results in the spirit of the representation of commutative C^* -algebras by Gelfand and Naimark (see Theorem 4.29 [4]) as rings of continuous functions. Our work naturally suggests future efforts to characterize the semi-hereditary Bezout rings obtained by our construction and belongs to the rich theory describing the divisibility of rings. For the divisibility of various classes of rings we refer to the recent book [8] and papers [7], [9]. The question as to whether Bezout semigroups characterize the divisibility theory of Bezout rings is still open.

A word about terminology. All structures are commutative, and rings have the identity element 1. A ring is called *Bezout* if every finitely generated ideal is principal. The extension of a semigroup S by a zero element will be denoted by S^\bullet .

1. BASIC NOTIONS AND PRELIMINARY RESULTS

For the benefit of the reader and for the sake of completeness we recall some basic definitions and notation and include some crucial results from [2]. The development of and comments on this supporting material can be found in full detail in [2].

Definition 1.1 (see Section 4 [3] and Definition 1.1 [2]). A *Bezout semigroup* S (in short a *B-semigroup*) is a commutative monoid S with 0 such that the divisibility relation $a|b \iff aS \supseteq bS$ is a partial order, called a natural partial order, inducing a distributive lattice on S , with a multiplication distributive on both meets and

joins, and S is *hyper-normal*, meaning that for any $a, b, d = a \wedge b \in S$, $a = da_1$ there is $b_1 \in S$ satisfying $a_1 \wedge b_1 = 1$, $b = db_1$.

A Bezout semigroup S is *semi-hereditary* if for each $a \in S$ there is $e^2 = e \in S$ with $eS = a^\perp = \{x \in S \mid ax = 0\}$.

A B-semigroup is called *cancellative* if nonzero elements form a cancellative monoid.

Since divisibility is a partial order, 1 is the unique invertible element in a B-semigroup. The most natural example of B-semigroups is the extension P^\bullet of the positive cone P of an abelian lattice-ordered group by an extra zero element. Therefore divisibilities of Bezout domains are cancellative B-semigroups. More generally, Proposition 1.4 [2] establishes

Proposition 1.1. *If R is a Bezout ring, then the divisibility theory S_R of R is a B-semigroup.*

For the sake of completeness and due to the importance of this result, we include a direct, detailed proof which does not appear in [2].

Proof. For any $a, b, z \in R$ and arbitrary $c \in aR \cap bR$, let $dR = aR + bR$, $d = ax + by$, $a = da_1$, $b = db_2$, $c = ac_1$ with some $a_1, b_2, c_1, d, x, y \in R$. Then $ab_2 = da_1b_2 = a_1b \in aR \cap bR$ and $bR = bR + cR + bc_1R = db_2R + ac_1R + bc_1R = db_2R + c_1(aR + bR) = db_2R + dc_1R = d(b_2R + c_1R)$, and hence $a_1bR = da_1(b_2R + c_1R) \supseteq da_1c_1R = cR$. Therefore $aR \cap bR$ is the principal ideal a_1bR . Consequently S_R is a sublattice of the ideal lattice of R which is a distributive lattice under reverse inclusion. Thus S_R is a distributive lattice under the natural partial order. It is trivial that ideal multiplication is distributive on meets, i.e. sums of ideals. In view of $(zR)(aR) + (zR)(bR) = zaR + zbR = z(aR + bR) = zdR$ we have $(zR)(aR) \cap (zR)(bR) = zaR \cap zbR = za_1bR = (zR)(a_1bR) = (zR)(aR \cap bR)$, which shows that multiplication is also distributive on joins. Moreover, if b_1 is any generator of the principal ideal $(1 - a_1x)R + b_2R$, then $d(1 - a_1x) = d - ax = by \in bR$ implies $bR = (dR)(b_1R)$, $b_1R + a_1R = R$, from which hyper-normality follows. \square

Remark 1.2. It is important to emphasize in the above proof that the ideals generated by b and db_1 coincide although b and db_1 are not, in general, equal. That is, Bezout rings are not necessarily Hermitean rings. Moreover, ideal multiplication is obviously distributive on sums (on meets of ideals) but generally not on joins (on intersections of ideals). In fact, by Lemma 2 [13] a domain is a Prüfer domain if and only if ideal multiplication is distributive on intersections of ideals. Note that Jensen’s result also holds in the more general setting of semiprime rings. Namely, a ring without nonzero nilpotent elements has a distributive ideal lattice if and only if ideal multiplication is distributive on intersections of ideals. Recall that an *arithmetical ring* is a ring with a distributive ideal lattice. The semiprime condition is essential. For example, the factor ring $k[x, y]/(x, y)^2$ of a polynomial ring in two variables over a field is obviously not an arithmetical ring, but ideal multiplication is clearly distributive over intersections of ideals. For further comments we refer to Remarks 1.4 [2].

Examples 1.2. One can obtain further classical examples of B-semigroups (see Examples 1.3 and Proposition 1.4 [2]) by considering

- (1) semigroups of principal ideals in unique factorization domains where meets (given as principal ideals generated by greatest common divisors) are not necessarily sums of ideals, e.g., in $k[x, y]$ one has $(x) \wedge (y) = (1) \neq (x, y)$;
- (2) semigroups of ideals in Dedekind domains;
- (3) semigroups of finitely generated ideals in Prüfer domains, or more generally, in arithmetical rings.

Corollary 1.3. *The divisibility theory S_R of a semi-hereditary ring R is a semi-hereditary B-semigroup.*

Proof. By Theorem VI.2.9 [5] a semi-hereditary ring R has weak dimension at most 1, implying (in view of the main result in [13]) that R is an arithmetical ring. Therefore S_R is order-isomorphic to the divisibility theory of an appropriate Bezout ring according to Corollary 8.1 [1]. Hence S_R is a B-semigroup by Proposition 1.1. Since the annihilator of a finitely generated ideal is a finite intersection of the annihilators of principal ideals, it is generated by an idempotent, provided that the ring is semi-hereditary. It follows that S_R is a semi-hereditary B-semigroup. \square

An obvious consequence of hyper-normality is

Corollary 1.4. *In a Bezout semigroup, idempotents form a Boolean algebra.*

Proof. For any idempotent $e = e^2$ hyper-normality implies an element f with $e \wedge f = 1, ef = 0$, whence $f^2 = f, e \vee f = 0$; i.e., f is the *complement* of e , completing the proof. \square

When working with B-semigroups, one has to deal with *filters* and the so-called *m-prime filters* instead of the usual ideals and prime ideals in semigroup theory. In view of the lattice structure, m-prime filters reveal additional information via a *hidden addition* induced by the meet operation. For example, the multiplicative semigroup of the natural numbers has uncountably many ideals, but the ring of integers has only countably many ideals. Recall that a *filter* in a B-semigroup S is a subset F closed under \wedge such that if $a \in F, b \in S$, and $a \leq b$, then $b \in F$. Therefore every filter is also an ideal of the semigroup S (in the usual sense) with respect to multiplication. A filter I is said to be *m-prime* if $ab \in I$ implies $a \in I$ or $b \in I$. Thus m-prime filters are particular cases of prime ideals in weak ideal systems (Aubert's x-systems), and Theorem 1.5 is related to the development in Section 4.4 of Halter-Koch's book [8]. An element $s \in S$ is a *zero-divisor* if $st = 0$ for some $0 \neq t \in S$. An element $a \in S$ is a *non-zero-divisor* if it is not a zero-divisor, i.e., if $a^\perp = 0$.

The following result is basic in our approach.

Theorem 1.5 (see Theorem 2.13 [2]). *For every m-prime filter I a relation Φ_I*

$$\forall x, y \in S : x \cong y \iff \exists s \notin I : x \leq ys \ \& \ y \leq xs$$

defines a congruence whose factor $S_\varphi = S/\Phi_I$ is a naturally totally ordered B-semigroup, and the congruence class of 0 is $K = \{x \in S \mid \exists s \notin I : sx = 0\}$.

Proof. It is routine to verify that the relation Φ_I is a congruence and $S \setminus I$ and K are sets of elements congruent to 1 and 0, respectively. Images in S_φ of $s \in S$ are denoted by s_φ . Given any two elements $x, y \in I$, hyper-normality allows us to write $d = x \wedge y, x = dx_1, y = dy_1, x_1 \wedge y_1 = 1$ for some $x_1, y_1 \in S$. Therefore at least one of x_1, y_1 does not belong to I . Consequently images of x, y in S_φ are

comparable; i.e., S_φ is totally ordered. Moreover, $x_\varphi \leq y_\varphi$ implies $x_\varphi = (x \wedge y)_\varphi$, whence $y_\varphi = x_\varphi z_\varphi$ for $z \in S$ with $y = (x \wedge y)z$. If the equality $y_\varphi = x_\varphi z_\varphi$ holds for some $x, y, z \in S$, then there exists $s \notin I$ with $x \leq xz \leq ys$, hence $x_\varphi \leq y_\varphi$. This shows that the induced order in S_φ is natural. Furthermore, if $a \in I, b \in S$ have the property that a, ab have the same nonzero image in S_φ , then by definition there is some $s \notin I$ with $ab \leq as$. If $b \in I$, then by putting $s \leq b \vee s = c = sc_1 \in I$ for some $c_1 \in S$ one has that $c, c_1 \in I$ and $0 \neq as = ab \vee as = a(b \vee s) = ac = asc_1$. Therefore hyper-normality implies that there is an element $x \in S$ with $asx = 0, c_1 \wedge x = 1$. This shows that $x \in I$ and thus $c_1 \notin I$ and so b cannot be an element of I , a contradiction. It follows that $b \notin I$. This implies that S_φ is hyper-normal, and it is now easy to check that S_φ is a B-semigroup. \square

Remark 1.6. For an m-prime filter I the factor of S by the congruence

$$\forall x, y \in S : x \cong y \iff \exists s \notin I : x \leq ys \ \& \ y \leq xs$$

is called the *localization* of S at I (sending $S \setminus I$ to 1).

The *nil radical* of a B-semigroup S is the (set-theoretical) intersection of all m-prime filters. It is routine to see that the nil radical of S is just the set of all nilpotent elements. S is called *semiprime* or *reduced* if its nil radical is 0. An m-prime filter I of S is called *minimal* if it is minimal in the set of all m-prime filters of S with respect to the set-theoretical inclusion. Throughout this paper \mathcal{X} will denote the set of all minimal m-prime filters of S . It is obvious that the nil radical is also the intersection of all minimal m-prime filters.

Definition 1.3. The *minimal spectrum* of S is the set \mathcal{X} of all minimal m-prime filters endowed with the *Zariski topology* by taking $\{D_s = \{I \in \mathcal{X} \mid s \notin I\}, s \in S\}$ as a basis of open sets.

It is routine to check that $\{D_s = \{I \in \mathcal{X} \mid s \notin I\}, s \in S\}$ satisfies the axioms for a basis of open sets in a topological space. For the sake of completeness we provide the proofs of the following obvious but important assertions.

Proposition 1.7. *Let S be a reduced B-semigroup.*

- (1) [14, Proposition 1.1 (3)] $\bigcup_{I \in \mathcal{X}} I$ is the set of all zero-divisors of R .
- (2) [14, Proposition 1.2 (1)] An m-prime filter I is minimal iff for all $s \in I, s^\perp \not\subseteq I$.

Proof. Let I be a minimal m-prime filter and let $0 \neq s \in I$. By Theorem 1.5 the localization S_I is a naturally totally ordered B-semigroup; hence it must be a Boolean algebra of two elements 1 and 0 because S is reduced and I is a minimal m-prime filter. This implies that there is an element $t \notin I$ with $ts = 0$. $t \notin I$ implies $t \neq 0$. Thus $\bigcup_{I \in \mathcal{X}} I$ consists entirely of zero-divisors. On the other hand, if $0 \neq s$ is a zero-divisor, then there exists $0 \neq t$ with $ts = 0$. Since S is reduced, there is a minimal m-prime filter I with $t \notin I$. Therefore $s \in I$ holds because $ts = 0 \in I$. This finishes the verification of (1).

If I is a minimal m-prime filter and $0 \neq s \in I$, then we have already seen that there is some $t \notin I$ with $ts = 0$, i.e, $s^\perp \neq 0$ and $s^\perp \not\subseteq I$. Conversely, let I be an m-prime filter such that $s^\perp \not\subseteq I$ holds for all $s \in I$. Assume indirectly that there is another m-prime filter J properly contained in I . Then for $s \in I \setminus J$ we have $s^\perp \subseteq J \subseteq I$, a contradiction. Thus, statement (2) holds. \square

Remark 1.8. Although $\bigcup_{I \in \mathcal{X}} I$ is the set of all zero-divisors of R , there exist in general m-prime filters consisting of zero-divisors which are not minimal.

2. SEMI-HEREDITARY B-SEMIGROUPS

For every element $a \in S$ let $S_a = \{s \in S \mid s^\perp = a^\perp\}$. The following simple but basic result is proved in [2]. For the sake of clarity and because of its importance for our development, we include the proof.

Proposition 2.1 (cf. Theorem 3.5 [2]). *If S is a semiprime B-semigroup, then for every element $a \in S$ the subsemigroup S_a is a cancellative semigroup closed under \wedge and \vee . The extension S_a^\bullet of S_a by a new extra zero element 0 is a B-semigroup if and only if its induced order is natural, i.e., if $S_a \ni b \leq c \in S_a$ implies $c = bu$ for some $u \in S_a$.*

Proof. Assume $x, y \in S_a$. Then $(x \wedge y)^\perp = x^\perp \cap y^\perp = a^\perp$. If $t(x \vee y) = 0$, then $txy = 0$. Hence $tx \in y^\perp = x^\perp$; thus $(tx)^2 = 0 \implies tx = 0$, i.e., $t \in x^\perp$. Thus S_a is a subsemigroup closed under both \wedge and \vee . If now $xz = yz$ with $x, y, z \in S_a$, then $(x \wedge y)z = (x \vee y)z$ holds. Hence one can assume without loss of generality that $x < y = dx$ and it follows that $zx = dzx$. Thus $x \neq y$ would imply $d \neq 1$. If there is some $c \in x^\perp = a^\perp$ with $c \wedge d = 1$, then we have $x = x \cdot 1 = x(c \wedge d) = xd = y$, a contradiction. Thus there is an m-prime filter I such that in the 0-cancellative naturally totally ordered factor S/Φ_I we have $d_\varphi \neq 1$, $(xz)_\varphi \neq 0$ and the equality $(xz)_\varphi = d_\varphi(xz)_\varphi$, a contradiction.

Assume now that S_a^\bullet is a B-semigroup and $b, c \in S_a$, $b \leq c$. Then S_a has an identity; i.e., there is an idempotent $e \in S_a$ and we have $c = bd$ for some $d \in S$. This implies $c = bd = bed = bu$ with $u = de \in S_a$, as is easy to check. Conversely, $a \leq a$ implies that $a = ae$ for some $e \in S_a$, and hence $e = e^2$ is the identity of S_a . Consequently S_a , endowed with 0, is a B-semigroup. \square

Corollary 2.2. *Let S be a B-semigroup.*

- (1) *If for every $a \in S$, S_a contains an idempotent, then S is semiprime.*
- (2) *S is semi-hereditary iff S_a contains an idempotent for each $a \in S$. In particular, semi-hereditary B-semigroups are disjoint unions of positive cones of lattice-ordered groups, whence reduced.*

Proof. If S_a contains an idempotent e , then $a^\perp = e^\perp = fS$ where f is the orthogonal complement of e . This shows the equivalence claimed in assertion (2) and hence claim (1). \square

Remarks 2.3. (1) If R is a semi-hereditary Bezout ring, then it is well-known (see [13]) as a consequence of a long homological argument that R is semiprime. Our proof provides a short, direct elementary justification. Note that semi-hereditary associative rings are not necessarily semiprime. For instance, upper triangular matrix rings over fields are hereditary and have nonzero nilpotent radicals.

- (2) It is well-known that non-trivial lattice-ordered abelian groups are torsion-free. However, in our treatment of semi-hereditary B-semigroups, the trivial group consisting only of the identity element is lattice-ordered in the obvious way. The reason for this convention is the case of Boolean algebras which are obviously B-semigroups S where S_a is the trivial group for any element $a \in S$.

Now we turn to the geometric structure of semi-hereditary B-semigroups. With the exception of claim (2), the ring-theoretical statements corresponding to assertions in the following theorem are well-known even for the larger class of reduced rings (see Section 1 in [14]).

Theorem 2.4. *For a B-semigroup S the following statements are equivalent:*

- (1) S is semi-hereditary.
- (2) S is reduced and for each element $a \in S$ there exists $b \in S$ such that $a^\perp = bS$.
- (3) S is reduced and for each $a \in S$ there exists $b \in a^\perp$ such that $a \wedge b$ is a non-zero-divisor.
- (4) The minimal spectrum \mathcal{X} of S is compact with respect to the Zariski topology.

Remarks 2.5. (1) The condition in claim (2) that a^\perp is a cyclic filter for each $a \in S$ means precisely that every principal filter is finitely presented. Bezout rings with this condition are exactly coherent Bezout rings.
 (2) Any $e^2 = e \in S$ together with its complement f induces a direct decomposition

$$S_e \times S_f \longrightarrow S_1 : (S_e \ni s, t \in S_f) \longmapsto s \wedge t \in S_1 \ni u \longmapsto (eu, fu) \in S_e \times S_f.$$

Consequently, the structure of a semi-hereditary B-semigroup S is completely described by its subsemigroup S_1 , which is a positive cone of a lattice-ordered abelian group G , and a Boolean algebra of idempotents in S such that each idempotent of S induces a direct decomposition of G with obvious compatibility conditions. Thus one might think that this Boolean algebra can be realized as a Boolean algebra of commuting idempotents in the endomorphism ring of the abelian group G . However, this is not necessarily the case in view of Remark 2.3 (2). Here is another simple example to the contrary. Let R be the subring in the direct product $\prod_{i \in \mathbb{N}} \mathbb{Q}$ of countably infinitely many copies of the field \mathbb{Q} of rationals generated by their direct sum; i.e., R is a subring of sequences, all but finitely many of whose entries are equal to an integer. Such a ring R is a semi-hereditary Bezout ring. Let S be a semigroup of divisibility of R . Then one can verify that a Boolean algebra of idempotents of S is isomorphic to a Boolean algebra generated by finite subsets of \mathbb{N} with respect to unions and intersections and S_1 is just the multiplicative semigroup \mathbb{N} . However, every idempotent e of S induces the same identity map of \mathbb{N} , and S_e consists either of one element or is isomorphic to S_1 .

Proof. The implication $1 \implies 2$ is trivial by Corollary 2.2 (2) and the definition of semiheredity.

The implication $2 \implies 3$ is obvious.

$3 \implies 4$. To show the compactness of \mathcal{X} , it is enough to see that every open covering of \mathcal{X} consisting of open sets of the form D_{a_i} contains a finite subcollection that also covers \mathcal{X} because $\{D_s \mid s \in S\}$ forms a basis of open sets for the Zariski topology. Let F be the filter generated by all of the a_i . One can assume that all the a_i are zero-divisors because $D_a = \mathcal{X}$ holds for a non-zero-divisor $a \in S$. If F does not contain non-zero-divisors, then F can be extended to an m-prime filter I disjoint from S_1 . Then I must be a minimal m-prime filter by Proposition 1.7 (2).

In fact, for any element $s \in I$ there is $t \in s^\perp$ with the non-zero-divisor $s \wedge t$, hence $t \notin I$. Consequently $s^\perp \not\subseteq I$ holds. Therefore we have shown that $I \in \mathcal{X}$. However, $I \notin \bigcup D_{a_i}$ holds because all the a_i belong to I ; i.e., $\{D_{a_i}\}$ is not an open covering of \mathcal{X} , a contradiction. Henceforth F must contain a non-zero-divisor d ; consequently, there are finitely many a_i 's, say a_1, \dots, a_n , such that d is contained in the filter generated by a_1, \dots, a_n . This shows $\bigcup_{i=1}^n D_{a_i} = \mathcal{X}$, which completes the verification that \mathcal{X} is compact.

4 \implies 1. For an arbitrary element $a \in S$ and $I \in \mathcal{X}$ the equality $aa^\perp = 0 \in I$ implies either $a \in I$ or $a \notin I$. The latter case implies $I \in D_a$. The first case shows that there is one $b \in a^\perp$ with $b \notin I$ in view of Proposition 1.7(2), thus $I \in D_b$. Therefore $\{D_a, D_b \mid s \in a^\perp\}$ is an open covering of \mathcal{X} . By the compactness of \mathcal{X} there are finitely many $s \in a^\perp$, and so, by taking their meet $b \in a^\perp$, we have $D_a \cup D_b = \mathcal{X}$. Consequently $d = a \wedge b$ is a non-zero-divisor. By the hyper-normality condition there are $a_1, b_1 \in S$ such that $a_1 \wedge b_1 = 1, a = da_1, b = db_1$. On the other hand, $0 = ab = d^2 a_1 b_1$ implies $a_1 b_1 = 0$ because d is a non-zero-divisor. Therefore $a_1^2 = a_1, b_1^2 = b_1$ and $b_1 S = a^\perp$ hold. This completes the proof of the theorem. \square

From the proof of the implication 3 \implies 4 one obtains the following assertion.

Corollary 2.6. *If F is a filter consisting of zero-divisors in a semi-hereditary B-semigroup, then F is contained in a minimal m -prime filter. In particular, minimal m -prime filters in semi-hereditary B-semigroups are maximal m -prime filters consisting of zero-divisors. Therefore, a reduced B-semigroup is semi-hereditary if and only if the set of minimal m -prime filters coincides with the set of maximal m -prime filters of zero-divisors.*

Remark 2.7. If a minimal m -prime filter F in a B-semigroup S is at the same time a maximal m -prime filter, then for each $a \notin F$ there is obviously $b \in F$ such that $a \wedge b$ is a non-zero-divisor but b is, in general, not annihilated by a .

It is well-known (see [13]) that a ring has weak dimension at most 1 if and only if its localization at each prime ideal is a valuation domain. The following result provides the topological property of the minimal spectrum that characterizes semi-hereditary rings among commutative rings with weak dimension at most 1.

Corollary 2.8. *A commutative ring R has weak dimension at most 1 and its minimal spectrum is compact if and only if R is semi-hereditary.*

Proof. By Corollary 1.3 and Theorem 2.4 it is enough to prove sufficiency. Let S be the divisibility of R . Then it is obvious that the minimal spectra of S and R are canonically homeomorphic. Therefore, again by Theorem 2.4, S is a semi-hereditary B-semigroup. This implies that the annihilator I^\perp of any finitely generated ideal of R is also a finitely generated idempotent ideal. Consequently, I^\perp is generated by an idempotent. In particular, every cyclic ideal of R is projective. Moreover, any finitely generated ideal I is flat because its localization I_P at any prime ideal P is either a free module of rank 1 over a valuation domain R_P or 0 in view of the equality $I^\perp = eR$ for some $e^2 = e \in R$. Therefore R is semi-hereditary by Proposition 2.7 [14]. \square

Proposition 2.9. *For every open and closed subset U in the minimal spectrum \mathcal{X} of a semi-hereditary B-semigroup there is precisely one idempotent $e^2 = e$ with $U = D_e$.*

Proof. Since U is open, $U = \bigcup D_{s_i}$ for appropriate elements s_i . Therefore $\{X \setminus U, D_{s_i}\}$ is an open covering of X because U is closed. Hence the compactness of X shows that there are finitely many s_i 's, say s_1, \dots, s_n , such that $U = \bigcup_{i=1}^n D_{s_i}$ which implies $U = D_s$ if $s = \bigwedge_{i=1}^n s_i$. By assumption there is an idempotent e with $s^\perp = e^\perp$. The inclusion $D_s \subseteq D_e$ obviously holds because $e \leq s$. If $I \in D_e \subseteq \mathcal{X}$, i.e., $e \notin I$ and $s \in I$, then for the orthogonal complement $f^2 = f$ of e the element $s \wedge f \in I$ is a non-zero-divisor, a contradiction by Proposition 1.7 (2). Thus $s \notin I$, which shows $D_e \subseteq D_s$. Hence $D_e = D_s$. If $D_e = D_g$ holds for some other idempotent $g^2 = g$, then $D_e = D_g = D_{e \wedge g} = D_{e \vee g}$ and hence $e = e \wedge g = e \vee g = g$, which completes the proof. \square

Theorem 2.10. *The minimal spectrum \mathcal{X} of a semi-hereditary B-semigroup S is a Stone space, i.e., a totally disconnected compact Hausdorff space. If B is the Boolean algebra of idempotents in S , then the map $I \in \mathcal{X} \mapsto I \cap B$ is a homeomorphism between \mathcal{X} and the prime spectrum \mathcal{M} of B .*

Proof. Let $\mathcal{X} \ni I \neq J \in \mathcal{X}$. Then there is an element $a \in I \setminus J$. By assumption there is an idempotent $e \in S$ such that $a^\perp = e^\perp$. If $f^2 = f$ is the orthogonal complement of e , then $I \in D_f$ and $J \in D_e$ imply that \mathcal{X} is Hausdorff and totally disconnected. It is clear that $I \cap B$ is a prime ideal of the Boolean algebra B ; i.e., it is easy to check that the map $I \in \mathcal{X} \mapsto J = I \cap B \in \mathcal{M}$ is well-defined with inverse the map which carries $J \in \mathcal{M}$ to $\bigcup_{e \in J} S_e \in \mathcal{X}$. The last assertion then follows immediately from the definitions of the topologies of \mathcal{X} and \mathcal{M} . \square

3. THE REPRESENTATION THEOREM

The main aim of this section is to prove the following theorem.

Theorem 3.1. *A semigroup S is the divisibility theory of a semi-hereditary Bezout ring if and only if it is a semi-hereditary B-semigroup.*

Proof. The necessity that the divisibility theory of a semi-hereditary Bezout ring is a semi-hereditary B-semigroup is obvious by Proposition 1.1. For the converse, we will construct a semi-hereditary Bezout ring with an arbitrary pre-described semi-hereditary B-semigroup S . For this aim, let L be an arbitrary field. It is well-known (see Theorem 4.5.1 [6]) that the localisation A of the semigroup algebra LS_1 at the set

$$\left\{ \sum_{i=1}^{i=n} k_i s_i \mid 0 \neq k_i \in L, s_i \in S_1, \bigwedge_{i=1}^n s_i = 1, n \in \mathbb{N} \right\}$$

of primitive elements is a Bezout domain whose divisibility semigroup is just S_1 extended with the new extra zero element 0. Moreover, S_1 is identified as a sub-semigroup of A under multiplication. Furthermore, let K be the quotient field of A . For every $e^2 = e \in S$ let $f^2 = f \in S$ be its orthogonal complement. Then $(fS)_1$ is order isomorphic to $\{e \wedge t \mid t \in S_f = (e^\perp)_1 \wedge e\}$, which is a multiplicatively closed subset of $S_1 \subseteq A$. Let A_e be the localisation of A at this subset. Thus we have the equality $A_1 = A$, and moreover A_e is a subring of A_g for every idempotent $g \in S$ satisfying $ge = g$. We further claim

Lemma 3.2. *The divisibility semigroup of A_e is isomorphic to S_e^\bullet under the map $f \wedge s \mapsto s \in S_e$.*

Proof. For any $t \in (e^\perp)_1 = (Sf)_1 = S_f$ and $s \in S_e$ in A , the invertibility of $t \wedge e \in A_e$ and the equalities

$$(t \wedge e)(f \wedge s) = (tf \wedge ts) \wedge (ef \wedge es) = (t \wedge 0) \wedge (0 \wedge s) = t \wedge s$$

imply that $(t \wedge s)A_e = (f \wedge s)A_e$, from which the assertion follows. \square

We are now in position to construct a semi-hereditary Bezout ring R whose semigroup of divisibility is S . Let $C_K(\mathcal{X})$ denote the ring of all continuous functions from the minimal spectrum \mathcal{X} of S into the field K endowed with the discrete topology. If U is an open and closed subset of \mathcal{X} , then a characteristic function χ_U which is 1 on U and 0 on $\mathcal{X} \setminus U$ is trivially an element of $C_K(\mathcal{X})$. Since \mathcal{X} is compact, Hausdorff and totally disconnected, a function from \mathcal{X} to K is continuous if it is a step function associated to a partition of \mathcal{X} as a finite disjoint union of open subsets,

i.e., a linear combination $\sum_{i=0}^{n-1} a_i \chi_{U_i}$ of characteristic functions χ_{U_i} with respect

to some finite partition $\mathcal{X} = \bigcup_{i=0}^{n-1} U_i$, $n \in \mathbb{N}$, of \mathcal{X} into pairwise disjoint open (and hence also closed) subsets U_i where the $\{a_i \in K\}$ are n distinct elements of K . The partition is finite because the topology on K is discrete. On the other hand, every open and closed subset U of \mathcal{X} is of the form D_e for a unique idempotent $e^2 = e \in S$. Therefore, for the sake of simplicity, the characteristic function χ_{D_e} will be denoted as χ_e . Hence a function $\varphi : \mathcal{X} \rightarrow K$ is continuous if and only if there are finitely many pairwise orthogonal idempotents e_i with $\bigwedge_{i=0}^{n-1} e_i = 1$ such

that φ is equal to the linear combination $\sum_{i=0}^{n-1} a_i \chi_{e_i}$. For the sake of convenience

we always set $a_0 = 0$; i.e., $\bigcup_{i=1}^{n-1} D_{e_i} = (\varphi)^{-1}(K \setminus \{0\})$ is the *support* of φ and D_{e_0} denotes the subset $(\varphi)^{-1}(0) \subseteq \mathcal{X}$. Note that the subset D_{e_0} can be the empty set.

We define R to be a subset of $C_K(\mathcal{X})$ consisting of linear combinations $\sum_{i=0}^{n-1} a_i \chi_{e_i}$

satisfying $\bigwedge_{i=0}^{n-1} e_i = 1$ where e_i are pairwise orthogonal idempotents and $a_i \in A_{e_i}$ for every index i .

Lemma 3.3. *R is a subring of $C_K(\mathcal{X})$.*

Proof. It is clear that R contains the identity element $\chi_{\mathcal{X}}$, its scalar multiples by elements of A , and the zero element χ_\emptyset of $C_K \mathcal{X}$. Therefore we have to show only that R is closed under addition and multiplication; i.e., $\varphi, \theta \in R$ imply $\varphi + \theta, \varphi \theta \in R$.

Write $\varphi = \sum_{i=0}^{n-1} a_i \chi_{e_i}$ and $\theta = \sum_{j=0}^{m-1} b_j \chi_{f_j}$ as linear combinations of characteristic

functions such that $a_0 = 0 = b_0$, $a_i \in A_{e_i}$, $b_j \in A_{f_j}$, and e_i as well as f_j are pairwise orthogonal idempotents of S with $\bigwedge_{i=0}^{n-1} e_i = 1 = \bigwedge_{j=0}^{m-1} f_j$, respectively. Note that some of the subsets D_{e_0} , D_{f_0} can be empty. It is obvious that $\{D_{e_i} \cap D_{f_j} = D_{e_i \vee f_j} = D_{e_i f_j}\}$ is a partition of \mathcal{X} induced by pairwise orthogonal idempotents $e_i \vee f_j = e_i f_j$. It follows that R is closed under multiplication and addition by noting that A_{e_i} , A_{f_j} are subrings of $A_{e_i f_j}$, which completes the proof. \square

We are now in position to show that R is a semi-hereditary Bezout ring whose semigroup of divisibility is S . Consider an arbitrary element $\varphi = \sum_{i=0}^{i=n-1} a_i \chi_{e_i} \in R$ where $a_0 = 0$, $a_i \in A_{e_i}$, and e_i are pairwise orthogonal idempotents of S with $\bigwedge_{i=0}^{n-1} e_i = 1$. Note that $(\varphi)^{-1}(0) = D_{e_0}$ can be empty. Let $e = \bigwedge_{i=1}^{n-1} e_i$. Since A_{e_i} is a Bezout domain whose semigroup of divisibility is $S_{e_i}^\bullet$, there is $s_i \in S_{e_i}$ such that $a_i A_{e_i} = (f_i \wedge s_i) A_{e_i}$ where f_i is the orthogonal complement of e_i in S . Let $s = \bigwedge_{i=1}^{n-1} s_i$. The equalities

$$(s \wedge e_0) \chi_e = \sum_{i=1}^{i=n-1} (s \wedge e_0) \chi_{e_i}, \quad (s \wedge e_0) \chi_{e_i} A_{e_i} = (f_i \wedge s_i) A_{e_i} = a_i A_{e_i}$$

imply $\varphi R = (s \wedge e_0) \chi_e R$. Therefore to finish the proof of the theorem, it is enough to see that R is a Bezout ring, i.e., to show that every ideal generated by two elements φ, θ is principal. As in the proof of Lemma 3.3 write $\varphi = \sum_{i=0}^{i=n-1} a_i \chi_{e_i}$

and $\theta = \sum_{j=0}^{j=m-1} b_j \chi_{f_j}$ as linear combinations of characteristic functions such that $a_0 = 0 = b_0$, $a_i \in A_{e_i}$, $b_j \in A_{f_j}$, where e_i and f_j are sets of pairwise orthogonal idempotents of S with $\bigwedge_{i=0}^{n-1} e_i = 1 = \bigwedge_{j=0}^{m-1} f_j$, respectively. Note that some of the subsets D_{e_0}, D_{f_0} can be empty. It is obvious that $\{D_{e_i} \cap D_{f_j} = D_{e_i \vee f_j} = D_{e_i f_j}\}$ is a partition of \mathcal{X} induced by pairwise orthogonal idempotents $e_i \vee f_j = e_i f_j$. In each ring, the ideal $A_{e_i f_j}$ generated by a_i and b_j is principal. This implies that $\varphi R + \theta R$ is a principal ideal, and the proof of Theorem 3.1 is complete. \square

By Proposition 1.3 the divisibility theory of semi-hereditary rings is described in the following corollary.

Corollary 3.4. *The divisibility theory of a semi-hereditary ring is a semi-hereditary Bezout semigroup, and any semi-hereditary Bezout semigroup can be realized as the divisibility theory of a semi-hereditary Bezout ring.*

ACKNOWLEDGMENT

The authors are grateful to the referee for calling their attention to Halter-Koch’s work on the divisibility theory of rings.

REFERENCES

1. D. D. Anderson, Multiplication ideals, multiplication rings, and the ring $R(X)$, *Canad. J. Math.* **28** (1976), 760 – 768. MR0424794 (54:12752)
2. P. N. Ánh, L. Márki, P. Vámos, Divisibility theory in commutative rings: Bezout semigroups, manuscript.
3. B. Bosbach, Representable divisibility semigroups, *Proc. Edinburgh Math. Soc.* **34** (1991), 45–64. MR1093175 (92m:20056)
4. R. G. Douglas, *Banach algebra techniques in operator theory*, GTM 179, 2nd ed., Springer-Verlag, New York, 1998. MR1634900 (99c:47001)
5. H. Cartan and S. Eilenberg, *Homological algebra*, Princeton, 1956. MR0077480 (17:1040e)
6. A. M. W. Glass and W. Charles Holland (eds), *Lattice-ordered groups: advances and techniques*, Kluwer Academic Publishers, 1989. MR1036072 (91i:06017)
7. F. Halter-Koch, A characterization of Krull rings with zero divisors, *Archivum Math.* **29** (1993), 119 – 122. MR1242634 (94h:13012)
8. F. Halter-Koch, *Ideal systems. An introduction to multiplicative ideal theory*, Marcel Dekker, 1998. MR1828371 (2001m:13005)

9. F. Halter-Koch, Construction of ideal systems with nice Noetherian properties, in *Non-Noetherian commutative ring theory* (eds. S. T. Chapman and S. Glaz), pp. 271 – 285, Kluwer, 2000. MR1858166 (2002h:13002)
10. M. Henriksen and M. Jerison, The space of minimal prime ideals of a commutative ring, *Trans. Amer. Math. Soc.* **115** (1965), 110 – 130. MR0194880 (33:3086)
11. N. Jacobson, *Structure of rings*, revised edition, AMS, 1964. MR0222106 (36:5158)
12. P. Jaffard, *Les système d'idéaux*, Dunod, Paris, 1960. MR0114810 (22:5628)
13. C. U. Jensen, A remark on arithmetical rings, *Proc. Amer. Math. Soc.* **15** (1964), 951–953. MR0179197 (31:3446)
14. E. Matlis, The minimal prime spectrum of a reduced ring, *Ill. J. Math.* **27** (1983), 353–391. MR698302 (84h:13010)
15. M. H. Stone, Applications of the theory of Boolean rings to general topology, *Trans. Amer. Math. Soc.* **41** (1937), 375–481. MR1501905

RÉNYI INSTITUTE OF MATHEMATICS, HUNGARIAN ACADEMY OF SCIENCES, 1364 BUDAPEST,
PF. 127, HUNGARY

E-mail address: `anh@renyi.hu`

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, COLORADO COLLEGE, COLORADO
SPRINGS, COLORADO 80903

E-mail address: `msiddoway@coloradocollege.edu`