

ISTVÁN VINCZE (1912–1999)
AND HIS CONTRIBUTION TO
LATTICE PATH COMBINATORICS
AND
STATISTICS

Endre Csáki¹

*A. Rényi Institute of Mathematics, Hungarian Academy of Sciences,
P.O.B. 127, H-1364, Budapest, Hungary
E-mail address: csaki@renyi.hu*

Abstract

A brief account of the life and work of István Vincze, a prominent Hungarian statistician is given. His contributions in various topics are discussed. They include: empirical distribution; Kolmogorov-Smirnov statistics; information theory; Cramér-Fréchet-Rao inequality; estimation of density; and a characterization problem.

MSC: primary 60C05; secondary 30D20; 60G50; 62B10; 62F10; 62G05; 62G30; 94A17

Keywords: Empirical distribution; Power function; Entropy; Cramér-Fréchet-Rao inequality; Density estimation; Entire function

1. Introduction

István Vincze was born in Szeged, Hungary, on February 26, 1912. After his graduation from the University of Szeged in 1935, he worked for a Hungarian Insurance Company until 1945. The second world war interrupted his career. After the war he worked for the Ministry of Education until 1950. Then, he was invited by the late Alfréd Rényi to join an Institute, whose main duty was set to do both theoretical and applied Mathematics. In this way, he became one of the founders of Mathematical Institute of the Hungarian Academy, whose director was Alfréd Rényi. Vincze was the Head of Statistics Department until his retirement

¹Supported by the Hungarian National Foundation for Scientific Research, Grant No. T 029621, T 037886 and T 043037.

in 1980 and, during 1950–1964, he also served as Deputy Director of the Institute. He was also a Professor in Statistics at the Eötvös Loránd University, Budapest. I had the privilege to be one of his numerous students in Statistics. He was considered as one of the main experts in both Theoretical and Applied Statistics in Hungary and also all over the world. Though in early stage of his research activity he was interested in Geometry, a subject on which he wrote several papers, including joint papers with Erdős, he has made significant contributions to several branches of Statistics, such as Quality Control, Nonparametric Statistics, Empirical distributions, Cramér–Rao inequality, Information Theory, etc. He is author of more than 100 research papers and 10 books.

He was awarded a number of honors in his life, including Hungarian State Prize in 1966, Gauss Ehrenplakette in 1978.

Except for the last two years of his life, he was very active even when he was over 80. He came and worked regularly in the Mathematical Institute, gave seminar talks, participated in conferences, such as Probastat, Bratislava, 1991 and 1994, Stochastic Modeling and Lattice Path Combinatorics, Delhi, 1994, Stability Problems, Kazan, 1995, Approximation Theory, Budapest, 1995, Statistical Conference, Poland, 1996. He was invited to the Combinatorial Methods Conference, Hamilton, Canada, 1997. He wrote a paper for the occasion (see Vincze and Törös, 1997), but an unfortunate accident prevented him from participating.

Professor István Vincze has visited many Universities and Institutes all over the world. He spent several months in China, GDR, USA, Canada, Argentina, etc. He was invited speaker on several Conferences, including three Berkeley Symposiums: 1960, 1965, 1970. He also organized a number of Conferences: European Meeting of Statisticians in Budapest, 1972, Nonparametric Statistical Inference in Budapest, 1980, Pannonian Symposiums on Mathematical Statistics in Bad Tatzmannsdorf in 1979, 1981 and 1983 and in Visegrád, Hungary, 1982. He was the director of the Unesco courses on Probability and Statistics, held in the Mathematical Institute, Budapest in 1964 and 1968.

Professor Vincze was a very kind man, his hospitality was legendary. He would walk with his guests through Budapest an entire day to show them the most important tourist attractions and serve as a real guide to explain the history of Hungary attached to the particular building and place. He was physically vigorous in all his life.

István Vincze will be remembered by the statistical community for his warmth, his humanity and his friendliness.

In this paper we summarize the most important contributions of Professor István Vincze in the following areas, focusing mainly in the first topic, but mentioning briefly his contributions in other subjects as well.

- Empirical distribution, random walk, lattice paths
- Information Theory
- Cramér–Fréchet–Rao inequality
- Estimation of density and its derivatives
- A characterization problem

2. Empirical distribution, random walk, lattice paths

Professor Vincze was a main contributor to the theory of empirical distributions and random walks (lattice paths), which were among his favourite topics. Consider a random sample

$$(X_1, X_2, \dots, X_n)$$

of size n , coming from a population with (theoretical) distribution function $F(x) = \mathbf{P}(X_1 \leq x)$. Empirical or sample distribution function is defined by

$$F_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbf{I}\{X_i \leq x\},$$

where $\mathbf{I}\{A\}$ stands for the indicator of the event A . Empirical distribution functions are widely used in statistics, nonparametric statistics in particular. In the two-sample case Gnedenko and Korolyuk (1951) developed a method based on random walk models. Let (X_1, X_2, \dots, X_n) and (Y_1, Y_2, \dots, Y_m) be two samples coming from continuous distributions. Let $F(x)$ and $G(x)$, resp. be their theoretical distribution functions and let $F_n(x)$ and $G_m(x)$, resp. be their empirical distribution functions. Testing the null hypothesis $H_0 : F(x) = G(x)$, a number of statistics has been investigated and their distributions, limiting distributions and other characteristics have been determined in the statistical literature. The idea of Gnedenko and Korolyuk was as follows: let

$$Z_1^* < Z_2^* < \dots < Z_{n+m}^*$$

denote the order statistics of the union of the two samples and define

$$\theta_i = \begin{cases} +1 & \text{if } Z_i^* = X_j \text{ for some } j, \\ -1 & \text{if } Z_i^* = Y_j \text{ for some } j \end{cases}$$

$i = 1, 2, \dots, n + m$. Put

$$S_0 = 0, \quad S_i = \theta_1 + \dots + \theta_i, \quad i = 1, 2, \dots, n + m.$$

Then $(S_0, S_1, \dots, S_{n+m})$ is a random walk path with $S_{n+m} = n - m$ and under H_0 each of them has the same probability. This idea of Gnedenko and Korolyuk enables one to determine the distributions of certain statistics by reducing the problems to combinatorial enumeration.

In a series of papers Professor Vincze and his collaborators presented a number of results on this subject. His first result concerns the joint distribution of the maximum and its location in the case $n = m$. Define

$$B_n^+ = n \max_{(x)} (F_n(x) - G_n(x)) = \max_{1 \leq i \leq 2n} S_i$$

and let R_n^+ be the first index i for which this maximum is achieved.

Moreover, put

$$B_n = n \max_{(x)} |F_n(x) - G_n(x)| = \max_{1 \leq i \leq 2n} |S_i|$$

and let R_n be the first index i for which this maximum is achieved.

Then, under H_0 , Vincze (1958) showed

$$\mathbf{P}(B_n^+ = k, R_n^+ = r) = \frac{k(k+1)}{r(2n-r+1)} \frac{\binom{r}{\frac{r+k}{2}} \binom{2n-r+1}{n-\frac{r+k}{2}}}{\binom{2n}{n}},$$

$k = 1, 2, \dots, n; r = k, k+2, \dots, 2n-k$.

Concerning the joint limiting distribution, it was shown that

$$\begin{aligned} & \lim_{n \rightarrow \infty} \mathbf{P} \left(\frac{B_n^+}{\sqrt{2n}} < y, \frac{R_n^+}{n} < z \right) = \\ & = \sqrt{\frac{2}{\pi}} \int_0^y \int_0^z \frac{u^2}{(v(1-v))^{3/2}} \exp \left(-\frac{u^2}{v(1-v)} \right) du dv. \end{aligned}$$

Furthermore,

$$\mathbf{P}(B_n = k, R_n = r) = \frac{2A_r^{(k)}A_{2n-r+1}^{(k+1)}}{\binom{2n}{n}},$$

with

$$A_r^{(k)} = \sum_{j=0}^{\infty} (-1)^j \frac{(2j+1)k}{r} \binom{r}{\frac{r+k}{2} + jk}$$

and

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathbf{P} \left(\frac{B_n}{\sqrt{2n}} < y, \frac{R_n}{n} < z \right) &= \\ &= \sqrt{\frac{8}{\pi}} \int_0^y \int_0^z f(u, v) f(u, 1-v) du dv, \end{aligned}$$

where

$$f(y, z) = \frac{y}{z^{3/2}} \sum_{j=0}^{\infty} (-1)^j (2j+1) e^{-\frac{(2j+1)^2 y^2}{2z}}.$$

Reimann and Vincze (1960) studied the case of different sample sizes. Define

$$B_{n,m}^+ = \max_{(x)} (nF_n(x) - mG_m(x)) = \max_{1 \leq i \leq n+m} S_i,$$

and let $R_{n,m}^+$ be the first index i for which this maximum is achieved.

Furthermore, put

$$\begin{aligned} B_{n,m} &= \max_{(x)} \left| nF_n(x) - mG_m(x) + \frac{m-n}{2} \right| - \frac{m-n}{2} = \\ &= \max_{1 \leq i \leq n+m} \left| S_i + \frac{m-n}{2} \right| - \frac{m-n}{2}, \end{aligned}$$

and let $R_{n,m}$ be the first index i for which this maximum is achieved. Let $m > n$. It was shown that

$$\mathbf{P}(B_{n,m}^+ = k) = \frac{2k+1+m-n}{m+k+1} \frac{\binom{m+n}{n-k}}{\binom{m+n}{n}}$$

and

$$\begin{aligned} \mathbf{P}(B_{n,m} = k) &= \\ &= \frac{1}{\binom{m+n}{n}} \sum_{j=-\infty}^{\infty} \left(\binom{m+n}{m+js} - \binom{m+n}{m+k+js} \right) = \end{aligned}$$

$$= \frac{2^{m+n+1}}{s \binom{m+n}{n}} \sum_{l=1}^{\infty} \cos^{m+n} \frac{l\pi}{s} \sin \frac{kl\pi}{s} \sin \frac{(s-k)l\pi}{s},$$

with $s = 2k + m - n$. Joint distributions of (B, R) and limiting distributions were also given.

These Reimann-Vincze statistics are different from the usual Kolmogorov-Smirnov statistics $\sup_{(x)} (F_n(x) - G_m(x))$, or $\sup_{(x)} |F_n(x) - G_m(x)|$, but have the advantage of easier computations of their distributions. Koul and Quine (1974) have shown that when the sample sizes are slightly different only, then the Bahadur efficiency of Reimann-Vincze statistics relative to the Kolmogorov-Smirnov statistics is 1.

In Vincze (1959, 1963) he proposed the use of generating functions to determine the above distributions and joint distributions. It was shown that

$$\begin{aligned} & \sum_{n=k}^{\infty} \sum_{r=k}^{2n-k} \binom{2n}{n} \mathbf{P}(B_n^+ = k, R_n^+ = r) v^r w^n = \\ & = \frac{2^{2k+1} v^k w^k}{(1 + \sqrt{1 - 4v^2 w})^k (1 + \sqrt{1 - 4w})^{k+1}}, \end{aligned}$$

and

$$\begin{aligned} & \sum_{n=k}^{\infty} \sum_{r=k}^{2n-k} \binom{2n}{n} \mathbf{P}(B_n = k, R_n = r) z^r w^{n-k} = \\ & = 2 \frac{(1 + \omega(w))^{k+1} (1 + \omega(z^2 w))^k}{(1 + (\omega(w))^{k+1}) (1 + (\omega(z^2 w))^k)}, \end{aligned}$$

with

$$\omega(z) = \frac{1 - \sqrt{1 - 4z}}{1 + \sqrt{1 - 4z}} = \frac{4z}{(1 + \sqrt{1 - 4z})^2}.$$

Now let

$$(X_1^* < X_2^* < \dots < X_n^*), \quad (Y_1^* < Y_2^* < \dots < Y_n^*)$$

denote the ordered samples. Then

$$\gamma_n = \sum_{i=1}^n \mathbf{I}\{X_i^* > Y_i^*\} = \frac{1}{2} \sum_{i=1}^{2n} (\mathbf{I}\{S_i > 0\} + \mathbf{I}\{S_{i-1} = +1, S_i = 0\}),$$

is the so-called Galton statistics. Chung and Feller (1949) showed that γ_n is uniformly distributed, i.e.,

$$\mathbf{P}(\gamma_n = g) = \frac{1}{n+1}, \quad g = 0, 1, 2, \dots, n.$$

Csáki and Vincze (1961) considered the number of crosses

$$\lambda_n = \sum_{i=1}^{2n-1} \mathbf{I}\{S_i = 0, S_{i-1}S_{i+1} < 0\},$$

and showed that

$$\mathbf{P}(\lambda_n = \ell - 1) = \frac{2\ell}{n} \frac{\binom{2n}{n-\ell}}{\binom{2n}{n}}, \quad \ell = 1, 2, \dots, n$$

and

$$\lim_{n \rightarrow \infty} \mathbf{P}(\lambda_n < y\sqrt{2n}) = 1 - e^{-2y^2}, \quad y \geq 0.$$

The joint exact and limiting distributions of (γ_n, λ_n) were also given. Namely,

$$\begin{aligned} \mathbf{P}(\gamma_n = g, \lambda_n = \ell - 1) &= \\ &= \frac{1}{\binom{2n}{n}} \frac{\ell^2}{2g(n-g)} \binom{2g}{g-\ell/2} \binom{2n-2g}{n-g-\ell/2} \end{aligned}$$

for ℓ even, and

$$\begin{aligned} \mathbf{P}(\gamma_n = g, \lambda_n = \ell - 1) &= \\ &= \frac{1}{\binom{2n}{n}} \frac{\ell^2 - 1}{4g(n-g)} \left(\binom{2g}{g-(\ell+1)/2} \binom{2n-2g}{n-g-(\ell-1)/2} \right) + \\ &\quad + \left(\binom{2g}{g-(\ell-1)/2} \binom{2n-2g}{n-g-(\ell+1)/2} \right) \end{aligned}$$

for ℓ odd, while concerning the joint limiting distribution, it was shown that

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathbf{P}(\gamma_n \leq zn, \lambda_n \leq y\sqrt{2n}) &= \\ &= \sqrt{\frac{2}{\pi}} \int_0^y \int_0^z \frac{u^2}{(v(1-v))^{3/2}} \exp\left(-\frac{u^2}{2v(1-v)}\right) du dv. \end{aligned}$$

Another use of the generating function method is found in Csáki and Vincze (1963b), where the joint distribution of the maximum and the number of crosses was given in the form

$$\sum_{n=1}^{\infty} \binom{2n}{n} \mathbf{P}\left(\max_x |F_n(x) - G_n(x)| = \frac{k}{n}, \lambda_n = \ell - 1\right) z^n =$$

$$= 2 \left(\frac{w - w^k}{1 - w^{k+1}} \right)^\ell, \quad \ell, k = 1, 2, \dots,$$

where

$$w = \frac{1 - \sqrt{1 - 4z}}{1 + \sqrt{1 - 4z}}, \quad |z| < \frac{1}{4}.$$

These and related results were presented by I. Vincze on a number of occasions in conferences, including the 4th, 5th and 6th Berkeley Symposium on Mathematical Statistics and Probability.

Vincze (1961) considered two-dimensional samples $(X_i^{(1)}, X_i^{(2)})$ and $(Y_i^{(1)}, Y_i^{(2)})$, $i = 1, 2, \dots, n$ from distributions having theoretical distribution functions $F(x, y)$ and $G(x, y)$, resp. Empirical distribution functions are denoted by $F_n(x, y)$ and $G_n(x, y)$, resp.. Testing $F = G$ he proposed to choose $\eta = y$ randomly according to the distribution function $H(y) = F(\infty, y)$ and consider the maximum deviation between $F_n(x, \eta)$ and $G_n(x, \eta)$. The following distributions were determined:

$$\begin{aligned} & \mathbf{P} \left(\max_{(x)} (F_n(x, \eta) - G_n(x, \eta)) < \frac{k}{n} \right) = \\ &= \frac{1}{(2n+1) \binom{2n}{n}} \sum_{i=0}^n \sum_{j=\max(0, i-k)}^n \binom{2n-i-j}{n-i} \left(\binom{i+j}{i} - \binom{i+j}{i-k} \right) \end{aligned}$$

and

$$\begin{aligned} & \mathbf{P} \left(\max_{(x)} |F_n(x, \eta) - G_n(x, \eta)| < \frac{k}{n} \right) = \\ &= \frac{1}{(2n+1) \binom{2n}{n}} \sum_{i=0}^n \sum_{j=\max(0, i-k)}^{\min(n, i+k)} \binom{2n-i-j}{n-i} \sum_{h=-\infty}^{\infty} (-1)^k \binom{i+j}{i+hk}. \end{aligned}$$

Vincze's idea in determining joint distributions was to construct tests based on a pair of statistics (instead of one single statistic) in order to improve the power of the tests. In Vincze (1965, 1967, 1968a) he studied the power function of several tests based on empirical distribution function. He computed the power function of two-sample Smirnov test and showed by numerical examples that the power can be increased by using a pair of statistics instead of one statistic. In particular, he considered the maximum and its location as a pair of statistics.

In Vincze (1972) several problems are treated concerning Kolmogorov-Smirnov statistics. Among others an explicit formula is given for two-sample case for discontinuous random variables and a discussion is given in the two-dimensional case.

Finally, we mention several results in Csáki and Vincze (1961, 1963a, 1964), concerning equivalence relations, proved by bijections. Define the following quantities:

$$\begin{aligned}\lambda'_n &= \sum_{i=1}^{2n-1} \mathbf{I}\{S_{i-1} = 0, S_i = +1\}, \\ \pi_n &= \sum_{i=1}^{2n} (\mathbf{I}\{S_i > 0\}), \\ \tau_\ell &= \min\{i : S_i = \ell\}.\end{aligned}$$

Then we have

$$\begin{aligned}\{\lambda_n = \ell - 1, S_1 = +1\} &\iff \{\tau_{2\ell} = 2n\}, \\ \{B_n^+ = \ell, R_n^+ = r\} &\iff \{\lambda'_n = \ell, \pi_n = r\}.\end{aligned}$$

Here $\{\dots\} \iff \{\dots\}$ means that there is a bijection between the two sets of random walk paths.

These relations are valid in the case when $S_{2n} = 0$. Now consider the general case, i.e., no restriction on the terminal point. Define

$$2\gamma_{2n}^{(2k)} = \sum_{i=1}^{2n} (\mathbf{I}\{S_i > 2k\} + \mathbf{I}\{S_{i-1} = 2k + 1, S_i = 2k\}).$$

Then

$$\{\gamma_{2n}^{(2k)} = g\} \iff \{S_{2n-2g} = S_{2n} = 2k\}.$$

The last relation implies the following distribution result:

$$\mathbf{P}(\gamma_{2n}^{(2k)} = g) = \frac{1}{2^{2n}} \binom{2g}{g} \binom{2n-2g}{n-g+k}, \quad g = 1, \dots, n-k.$$

In case $k = 0$, we recover the finite arcsine law of Chung and Feller (1949).

3. Information Theory

Vincze (1960) gave an interpretation of the I -divergence as below, concerning the information of a continuous random variable relative to "the distribution of our interest".

The entropy of a system of events A_1, A_2, \dots, A_n , in the case $P(A_i) = p_i$ ($i = 1, 2, \dots, n$), is $E_n = \sum_{i=1}^n p_i \log(1/p_i)$. Consider the quantity

$$I_n = \log n - E_n = \sum_{i=1}^n p_i \log np_i, \quad 0 \leq I_n \leq \log n,$$

which is called the "information of the system of events". Let now ξ be a continuous random variable with distribution function $F(x)$ and density function $F'(x) = f(x) > 0$ for any value of x . Vincze introduces a distribution function ϕ called the "distribution of our interest". Consider a system of divisions of the real line such that the points of division are regarded as the quantiles of a distribution function, that is, $\phi(x_k^{(n)}) = k/n$, $k = 1, 2, \dots, n-1$; $n = 1, 2, \dots$, $\phi(-\infty) = 0$, $\phi(\infty) = 1$. For each n the information of the system of discrete events $A_i^{(n)} = \{x_{i-1}^{(n)} \leq \xi < x_i^{(n)}\}$ ($i = 1, 2, \dots, n$) is

$$I_{n,\phi}(\xi) = \sum_{i=1}^n (F(x_i^{(n)}) - F(x_{i-1}^{(n)})) \log(n(F(x_i^{(n)}) - F(x_{i-1}^{(n)}))),$$

or, after the substitution $1/n = \phi(x_i^{(n)}) - \phi(x_{i-1}^{(n)})$,

$$I_{n,\phi}(\xi) = \sum_{i=1}^n (F(x_i^{(n)}) - F(x_{i-1}^{(n)})) \log \frac{F(x_i^{(n)}) - F(x_{i-1}^{(n)})}{\phi(x_i^{(n)}) - \phi(x_{i-1}^{(n)})}.$$

Assuming that $\phi'(x) = \varphi(x) > 0$ for all real x , by a limiting process one obtains the relation

$$\lim_{n \rightarrow \infty} I_{n,\phi}(\xi) = \int_{-\infty}^{\infty} f(x) \log \frac{f(x)}{\varphi(x)} dx = I_\phi(\xi),$$

which is the so-called I -divergence (Kullback, 1959). The expression $I_\phi(\xi)$ can be regarded as the information of the random variable relative to the distribution ϕ of our interest.

For another use of I -divergence as treated in Vincze (1974), we state the following result of Sanov (1957): Let X be a random variable whose distribution function is $\phi(x)$ and $F(x)$ be any distribution function such that the Borel measures μ_ϕ and μ_F induced, respectively, by $\phi(x)$ and $F(x)$ are equivalent; let $\phi_N(x)$ be the empirical distribution function of X after a large number N of independent trials; then for every $\varepsilon > 0$

$$\mathbf{P}(\sup_x |\phi_N(x) - F(x)| < \varepsilon) \cong \exp(-NI),$$

where $I = \int \log[dF(x)/d\phi(x)] dF(x)$ is the I -divergence. On the basis of this result, Vincze (1974) gives a correct formulation of the maximum-probability principle valid for both continuous and discrete random quantities. The principle consists in finding the distribution function $F(x)$ which minimizes the I -divergence under the constraints $\int x dF(x) = m \neq \int x d\phi(x)$ and $\int dF(x) = 1$. It is then shown that the I -divergence is the natural extension to continuous random variables of Shannon's formula for discrete entropy. Consequently, the maximum-probability principle provides an information-theoretical foundation of statistical mechanics, as suggested by Jaynes (1957), who considered only the discrete case.

Vincze (1975) also extends the maximum-probability principle to the case where the components of the system are not statistically independent, and for illustration he discusses the derivation of the Bose-Einstein and the Fermi-Dirac distributions.

Further results of I. Vincze in this area can be found in Puri and Vincze (1989, 1990, 1992), Vincze and Törös (1997) and Vincze (1962, 1968b, 1975).

4. Cramér–Fréchet–Rao inequality

Let $X = (X_1, X_2, \dots, X_n)$ be a sample from a distribution having (joint) density $p(x; \theta) = p(x_1, x_2, \dots, x_n; \theta)$ with respect to a measure μ , where θ is a parameter. Let $t(X)$ be an unbiased estimator of $g(\theta)$, i.e. $E_\theta(t(X)) = g(\theta)$. Cramér (1946), Fréchet (1943), Rao (1945) concluded the following inequality:

$$\text{Var}_\theta(t(X)) \geq \frac{(g'(\theta))^2}{I(\theta)},$$

with

$$I(\theta) = \int \left(\frac{\partial p}{\partial \theta} \right)^2 p(x; \theta) dx.$$

For fixed θ, θ' , Vincze (1979, 1981) considered the mixture

$$p_\alpha = p_\alpha(x; \theta, \theta') = (1 - \alpha)p(x; \theta) + \alpha p(x; \theta'), \quad 0 < \alpha < 1$$

with α being a new parameter. Then

$$\hat{\alpha} = \frac{t(X) - g(\theta)}{g(\theta') - g(\theta)}$$

is an unbiased estimator of α .

It follows that

$$\text{Var}_\alpha(\hat{\alpha}) \geq \frac{1}{J_\alpha(\theta, \theta')},$$

where

$$J_\alpha(\theta, \theta') = \int \frac{(p(x; \theta') - p(x; \theta))^2}{p_\alpha(x; \theta, \theta')} d\mu.$$

Then

$$(1 - \alpha)\text{Var}_\theta(t(X)) + \alpha\text{Var}_{\theta'}(t(X)) \geq \frac{1}{J_\alpha(\theta, \theta')} - \alpha(1 - \alpha)$$

and in the case when $Var_\theta(t(x))$ does not depend on θ , Vincze concluded the following lower bound:

$$Var(t(X)) \geq \sup_{\alpha} \sup_{\theta'} \alpha(1 - \alpha)(g(\theta') - g(\theta))^2 \left(\frac{1}{\alpha(1 - \alpha)J_{\alpha}} - 1 \right).$$

In certain cases this gives a reasonably good bound. This problem was further investigated by Puri and Vincze (1985), Govindarajulu and Vincze (1989) and Vincze (1992, 1996). It was shown among others that for the translation parameter of the uniform distribution this lower bound is of order n^{-2} , which is attainable.

5. Estimation of density and its derivatives

Let $f(x)$ be a density on the interval (a, b) and for positive integer consider a partition $a = x_0^{(n)} < x_1^{(n)} < x_2^{(n)} < \dots < x_n^{(n)} = b$. Put

$$s_k^{(n)} = \frac{\int_{x_k^{(n)}}^{x_{k+1}^{(n)}} t f(t) dt}{\int_{x_k^{(n)}}^{x_{k+1}^{(n)}} f(t) dt}, \quad k = 0, 1, 2, \dots, n - 1.$$

Rényi (1952, personal communication) raised the question whether $f(x)$ can be determined if for each n we know $\{s_0^{(n)}, s_1^{(n)}, s_2^{(n)}, \dots, s_{n-1}^{(n)}\}$ for a partition, such that

$$\lim_{n \rightarrow \infty} \max_{1 \leq i \leq n} (x_i^{(n)} - x_{i-1}^{(n)}) = 0.$$

Vincze (1954) answered this question in the affirmative. His idea was to show that

$$\frac{s(u, v) - \frac{u+v}{2}}{(v-u)^2} \rightarrow \frac{1}{12} \frac{f'(x)}{f(x)}, \quad \text{when } u, v \rightarrow x,$$

where

$$s(u, v) = \frac{\int_u^v t f(t) dt}{\int_u^v f(t) dt}.$$

This was extended by A.K. Gupta and Vincze (1991) as follows.

$$2^r (2r + 1)!! \frac{\int_u^v L_r(t; u, v) f(t) dt}{(v-u)^{r+1}} \rightarrow f^{(r)}(x), \quad \text{when } u, v \rightarrow x,$$

where $f^{(r)}$ denotes the r -th derivative of f and $L_r(t; u, v)$ denotes the Legendre polynomial of degree r belonging to the interval (u, v) normalized such that

$$\int_u^v L_r^2(t; u, v) dt = \frac{v - u}{2r + 1}.$$

Now suppose we want to estimate

$$I = \int_a^b \psi \left(\frac{f'(x)}{f(x)} \right) f(x) dx,$$

where $\psi(y)$ is a given function. For example, $\psi(y) = y^2$.

Assume that (X_1, \dots, X_N) is a random sample taken from a population with the distribution function having the density $f(x)$. For a partition $x_0 = a < x_1 < x_2 < \dots < x_n = b$ introduce the following notations:

$$\begin{aligned} \nu_i &= \sum_{j=1}^N I\{x_{i-1} \leq X_j < x_i\}, \\ \bar{X}_{(i)} &= \frac{1}{\nu_i} \sum_{j=1}^N X_j I\{x_{i-1} \leq X_j < x_i\}, \\ m_i &= \frac{x_{i-1} + x_i}{2}, \quad d_i = \frac{1}{\sqrt{12}}(x_i - x_{i-1}). \end{aligned}$$

Csáki and Vincze (1977) showed that under certain regularity conditions,

$$I_n = \sum_{i=1}^n \psi \left(\frac{\bar{X}_{(i)} - m_i}{d_i^2} \right)$$

is a consistent estimator of I , as $N \rightarrow \infty$.

A related question was treated in Csáki and Vincze (1978) as follows. Under the previous notation for $\bar{X}_{(i)}$, consider

$$\bar{\chi}_n^2 = \sum_{i=1}^n \left(\frac{\bar{X}_{(i)} - E_i}{\sigma_i} \right)^2 \nu_i,$$

where

$$\begin{aligned} E_i &= \mathbf{E}(X_1 | x_{i-1} \leq X_1 < x_i), \\ \sigma_i^2 &= \text{Var}(X_1 | x_{i-1} \leq X_1 < x_i). \end{aligned}$$

It was shown that (for fixed n), as $N \rightarrow \infty$, the limiting distribution of the above defined $\bar{\chi}^2$ statistics is *chi-square* with n degrees of freedom. This provides an alternative method for a goodness of fit test instead of the usual Pearson's chi-square test.

6. A characterization problem

Rényi and Vincze posed the following question:

Let

$$f(t) = 1 + a_1 t + a_2 t^2 + \dots + a_n t^n + \dots$$

be an entire function. Suppose that, on the one hand,

$$(6.1) \quad p_i = \frac{a_i t^i}{f(t)}, \quad i = 1, 2, \dots$$

is a probability distribution for all fixed $t > 0$, i.e.,

$$\sum_{i=0}^{\infty} \frac{a_i t^i}{f(t)} = 1, \quad t > 0,$$

and, on the other hand, for each $i = 0, 1, 2, \dots$

$$(6.2) \quad \frac{a_i t^i}{f(t)}$$

is a density, i.e.,

$$\int_0^{\infty} \frac{a_i t^i}{f(t)} dt = 1, \quad i = 0, 1, 2, \dots$$

Is it true that $f(t) = e^t$?

It is clear that for $f(t) = e^t$, $a_i = 1/i!$, (6.1) is the Poisson distribution with parameter t , while (6.2) is a gamma density. The converse, i.e. to show that $f(t) = e^t$ is the only solution, proved to be a rather hard problem. This (open) problem was also mentioned in a book by Hayman (1967).

In attacking this problem, Vincze and his coauthors (Hayman and Vincze, 1978; Hall and Vincze, 1981; Vincze, 1984, 1988; Csordás and Vincze, 1992), though did not solve the problem completely, made significant steps toward the solution and obtained many interesting side-results. It was shown in Hayman and Vincze (1978) that

$$e^{t-c\sqrt{t}} < f(t) < e^{t+c\sqrt{t}}$$

with some constant $c > 0$. Based on this result, a final answer (i.e., under the given conditions, it follows that $f(t) = e^t$) was given by Miles and Williamson (1986).

Acknowledgements

The author is grateful to the referees for their valuable comments.

References

- Chung, K.L., Feller, W., 1949. Fluctuations in coin tossing. *Proc. Nat. Acad. Sci. USA* 35, 605–608.
- Cramér, H., 1946. *Mathematical Methods of Statistics*. Princeton University Press, Princeton.
- Csáki, E., Vincze, I., 1961. On some problems connected with the Galton test. *Publ. Math. Inst. Hungar. Acad. Sci.* 6, 97–109.
- Csáki, E., Vincze, I., 1963a. On some combinatorial relations concerning the symmetric random walk. *Acta Sci. Math. (Szeged)* 24, 231–235.
- Csáki, E., Vincze, I., 1963b. Two joint distribution laws in the theory of order statistics. *Mathematica (Cluj)* 5, 27–37.
- Csáki, E., Vincze, I., 1964. On some distributions connected with the arcsine law. *Publ. Math. Inst. Hungar. Acad. Sci.* 8, 281–291.
- Csáki, E., Vincze, I., 1977. A lemma on real functions and its application in the nonparametric theory. In: R. Bartoszyński, E. Fidelis, W. Klonecki (Eds.), *Proceedings of the Symposium to honour Jerzy Neyman*. Polish Scientific Publishers, Warsaw, pp. 83–92.
- Csáki, E., Vincze, I., 1978. On limiting distribution laws of statistics analogous to Pearson's chi-square. *Math. Operationsforsch. Statist., Ser. Statistics* 9, 531–548.
- Csordás, G., Vincze, I., 1992. Convexity properties of power series with logarithmically s-concave coefficients. *Analysis Mathematica* 18, 3–13.
- Fréchet, M., 1943. Sur l'extension de certaines évaluations statistiques au case de petits échantillons. *Rev. Inst. Internat. Statist.* 11, 182–205.
- Gnedenko, B.V., Korolyuk, V.S., 1951. On the maximum discrepancy between two empirical distribution functions (Russian). *Dokl. Akad. Nauk SSSR* 80, 525–528; English translation: *Selected Transl. Math. Statist. Probab., Amer. Math. Soc.* 1, 13–16.

- Govindarajulu, Z., Vincze, I., 1989. The Cramér–Fréchet–Rao inequality for sequential estimation in non-regular case. In: Y. Dodge (Ed.), *Statistical Data Analysis and Inference*. North Holland, Amsterdam, pp. 257–268.
- Gupta, A.K., Vincze, I., 1991. Approximation of higher order derivatives of a real function. In: J. Szabados, K. Tandori (Eds.), *Approximation Theory* (Kecskemét, Hungary, 1990). Coll. Math. Soc. J. Bolyai, Vol. 58, pp. 339–342.
- Hall, R.R., Vincze, I., 1981. On a simultaneous characterization of the Poisson law and the gamma distribution. In: D. Dugué, E. Lukács, V.K. Rohatgi (Eds.), *Analytical Methods in Probability Theory*. Lecture Notes in Mathematics, Vol. 861. Springer, Berlin, pp. 54–59.
- Hayman, W.K., 1967. *Research Problems in Function Theory*. The Athlone Press University of London, London.
- Hayman, W.K., Vincze, I., 1978. A problem on entire functions. In: N.N. Bogoljubov, M.A. Lavrent’ev, A.V. Bicadze (Eds.), *Complex Analysis and its Applications*, dedicated to I.N. Vekua on his 70th birthday. Izd. Nauka, Moscow, pp. 591–594.
- Hayman, W.K., Vincze, I., 1979. Markov–type inequalities and entire functions. In: B. Gyires (Ed.), *Analytic Function Methods in Probability Theory* (Debrecen, Hungary, 1977). Coll. Math. Soc. J. Bolyai, Vol. 21, North-Holland, Amsterdam, pp. 153–163.
- Jaynes, E.T., 1957. Information theory and statistical mechanics. *Phys. Rev.* (2) 106, 620–630.
- Koul, H.L., Quine, M.P., 1974. The Bahadur efficiency of the Reimann-Vincze statistics. *Studia Sci. Math. Hungar.* 9, 399–403.
- Kullback, S., 1959. *Information Theory and Statistics*. Wiley, New York.
- Miles, J., Williamson, J., 1986. A characterization of the exponential function. *J. London Math. Soc.* (2) 33, 110–116.
- Puri, M.L., Vincze, I., 1985. On the Cramér–Fréchet–Rao inequality for translation parameter in the case of finite support. *Statistics* 16, 495–506.
- Puri, M.L., Vincze, I., 1989. Information and mathematical statistics. In: P. Mandl, M. Hušková (Eds.), *Proceedings of the Fourth Prague Symposium on Asymptotic Statistics* (Prague, 1988). Charles University, Prague, pp. 447–456.
- Puri, M.L., Vincze, I., 1990. Measure of information and contiguity. *Statist. Probab. Lett.* 9, 223–228.
- Puri, M.L., Vincze, I., 1992. The Neyman-Pearson probability ratio and information. In: S. Schach, G. Trenkler (Eds.), *Data Analysis and Statistical Inference*. Verlag Josef Eul, Bergisch Gladbach, pp. 53–64.

- Reimann, J., Vincze, I., 1960. On the comparison of two samples with slightly different sizes. *Publ. Math. Inst. Hungar. Acad. Sci.* 5, 293–309.
- Rao, C.R., 1945. Information and accuracy attainable in the estimation of statistical parameter. *Bull. Calcutta Math. Soc.* 37, 81–91.
- Sanov, I.N., 1957. On the probability of large deviations of random magnitudes (Russian). *Mat. Sbornik (N.S.)* 42 (84), 11–44; English translation: *Selected Transl. Math. Stat. Probab., Amer. Math. Soc.* 1, 213–244.
- Vincze, I., 1954. Determination of distributions by means of their conditional expectations (Hungarian). *Magyar Tud. Akad. Mat. Fiz. Oszt. Közl.* 4, 513–523.
- Vincze, I., 1958. Einige zweidimensionale Verteilungs- und Grenzverteilungssätze in der Theorie der geordneten Stichproben. *Publ. Math. Inst. Hungar. Acad. Sci.* 2, 183–209.
- Vincze, I., 1959. On some joint distribution and joint limiting distribution in the theory of order statistics, II. *Publ. Math. Inst. Hungar. Acad. Sci.* 4, 29–47.
- Vincze, I., 1960. An interpretation of the I-divergence of information theory. In: *Transactions of the Second Prague Conference on Information Theory, Statistical Decision Functions and Random Processes.* Publ. House Czechoslovak Acad. Sci., Prague, pp. 681–684.
- Vincze, I., 1961. On two sample tests based on order statistics. In: *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability.* University of California Press, Berkeley, Vol. I, pp. 695–705.
- Vincze, I., 1962. Some questions concerning the probabilistic concept of information (Hungarian). *Magyar Tud. Akad. Mat. Fiz. Oszt. Közl.* 12, 7–14.
- Vincze, I., 1963. A generating function in the theory of order statistics. *Publicationes Mathematicae* 10, 82–87.
- Vincze, I., 1965. On the power function of the Kolmogorov–Smirnov and other non-parametric tests (Hungarian). *Magyar Tud. Akad. Mat. Fiz. Oszt. Közl.* 15, 97–105.
- Vincze, I., 1967. Some questions connected with two sample tests of Smirnov type. In: *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability.* University of California Press, Berkeley, Vol. I, pp. 654–666.
- Vincze, I., 1968a. On the power of the Kolmogorov–Smirnov two-sample test and related nonparametric tests. In: K. Sarkadi, I. Vincze (Eds.), *Studies in Mathematical Statistics: Theory and Applications.* Publishing House of the Hung. Acad. Sci., Budapest, pp. 201–210.
- Vincze, I., 1968b. On the information–theoretical foundation of mathematical statistics. In: A. Rényi (Ed.), *Proceedings of the Colloquium on Information Theory.* J. Bolyai

- Mathematical Society, Budapest, Vol II, pp. 503–509.
- Vincze, I., 1970. On Kolmogorov–Smirnov type distribution theorems. In: M.L. Puri (Ed.), *Nonparametric Techniques in Statistical Inference*. Cambridge University Press, London, pp. 385–401.
- Vincze, I., 1972. On some results and problems in connection with statistics of the Kolmogorov–Smirnov type. In: *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability*. University of California Press, Berkeley, Vol. I, pp. 459–470.
- Vincze, I., 1974. On the maximum probability principle in statistical physics. In: J. Gani, K. Sarkadi, I. Vincze (Eds.), *Progress in Statistics*. Coll. Math. Soc. J. Bolyai 9, pp. 869–893.
- Vincze, I., 1975. On some problems in connection with the Bose-Einstein statistic. *Sankhyā* 37, 355–362.
- Vincze, I., 1979. On the Cramér–Fréchet–Rao inequality in the non-regular case. In: J. Jurečková (Ed.), *Contributions to Statistics. J. Hájek Memorial Volume*. Reidel, Dordrecht, pp. 253–262.
- Vincze, I., 1981. Remark to the derivation of the Cramér–Fréchet–Rao inequality in the regular case. In: P. Révész, L. Schmetterer, V.M. Zolotarev (Eds.), *Proceedings of the First Pannonian Symposium on Mathematical Statistics. Lecture Notes in Statistics*, Vol. 8, Springer, Berlin, pp. 284–289.
- Vincze, I., 1984. Contribution to a characterization problem. J. Mogyoródi, I. Vincze, W. Wertz (Eds.), *Proceedings of the Third Pannonian Symposium on Mathematical Statistics (Visegrád, Hungary, 1982)*. Reidel, Dordrecht, pp. 353–361.
- Vincze, I., 1988. On a probabilistic characterization theorem. In: W. Grossmann, J. Mogyoródi, I. Vincze, W. Wertz (Eds.), *Proceedings of the Fifth Pannonian Symposium on Mathematical Statistics (Visegrád, Hungary, 1985)*. Reidel, Dordrecht, pp. 213–219.
- Vincze, I., 1992. On nonparametric Cramér–Rao inequalities. In: P.K. Sen, I.A. Salama (Eds.), *Order Statistics and Nonparametrics: Theory and Applications*. North Holland, Amsterdam, pp. 439–454.
- Vincze, I., 1996. Cramér–Rao type inequality and a problem of mixture of distributions. *ProbaStat '94 (Smolenice Castle, 1994)*. Tatra Mt. Math. Publ. 7, 237–245.
- Vincze, I., Törös, R., 1997. The joint energy distributions of the Bose-Einstein and of the Fermi-Dirac particles. In: N. Balakrishnan (Ed.), *Advances in Combinatorial Methods and Applications to Probability and Statistics*. Birkhäuser, Boston, pp. 441–449.