

A supplement to Chebotarev's density theorem

On the 50th Anniversary of Chen's Theorem and the 100th Anniversary of Chebotarev's Theorem

Gergely Harcos^{1,*} & Kannan Soundararajan²

¹*Alfréd Rényi Institute of Mathematics, Budapest 1364, Hungary;*

²*Department of Mathematics, Stanford University, Stanford, CA 94305, USA*

Email: gharcos@renyi.hu, ksound@stanford.edu

Received October 26, 2022; accepted April 17, 2023; published online June 26, 2023

Abstract Let L/K be a Galois extension of number fields with Galois group G . We show that if the density of prime ideals in K that split totally in L tends to $1/|G|$ with a power saving error term, then the density of prime ideals in K whose Frobenius is a given conjugacy class $C \subset G$ tends to $|C|/|G|$ with the same power saving error term. We deduce this by relating the poles of the corresponding Dirichlet series to the zeros of $\zeta_L(s)/\zeta_K(s)$.

Keywords Chebotarev's density theorem, Artin L -functions, Heilbronn characters

MSC(2020) 11R42, 11M41

Citation: Harcos G, Soundararajan K. A supplement to Chebotarev's density theorem. *Sci China Math*, 2023, 66: 2749–2753, <https://doi.org/10.1007/s11425-022-2141-1>

1 Introduction

This paper arose from an amusing observation by MathOverflow user Lucia [11], which informally says that if the density of primes congruent to 1 modulo q tends to $1/\phi(q)$ rapidly, then for all $(a, q) = 1$ the density of primes congruent to a modulo q also tends to $1/\phi(q)$ rapidly. Here, the density refers to the actual proportions. More precisely, for any $\sigma \geq 1/2$, the asymptotic

$$\psi(x; q, 1) = \psi(x)/\phi(q) + O_\varepsilon(x^{\sigma+\varepsilon}) \quad (1.1)$$

implies that

$$\psi(x; q, a) = \psi(x)/\phi(q) + O_\varepsilon(x^{\sigma+\varepsilon}) \quad \text{for all } (a, q) = 1. \quad (1.2)$$

The reason is simple. The relation (1.1) implies that the function

$$\sum_{\substack{\chi \bmod q \\ \chi \neq \chi_0}} \frac{L'}{L}(s, \chi) = \int_1^\infty (\psi(x, \chi_0) - \phi(q)\psi(x; q, 1)) \frac{s}{x^{s+1}} dx$$

extends analytically to the half-plane

$$H_\sigma := \{s \in \mathbb{C} : \Re(s) > \sigma\}.$$

* Corresponding author

That is, the product of the Dirichlet L -functions $L(s, \chi)$ ($\chi \neq \chi_0$) has no zero or pole in H_σ . As these L -functions are entire, none of them has a zero or pole in H_σ , and (1.2) follows easily.

Our goal is to show that this phenomenon persists in the context of Chebotarev’s density theorem [4–6] (see [12, Chapter VII, Theorem 13.4]), even though the underlying Artin L -functions are only conjectured to be entire (Artin’s conjecture).

Corollary. *Let L/K be a Galois extension of number fields, with G denoting the Galois group. Suppose $\sigma \geq 1/2$ is such that for any $\varepsilon > 0$ and $x \geq 2$ the asymptotic formula*

$$\sum_{\substack{N(\mathfrak{p}) \leq x \\ \text{Frob}(\mathfrak{p})=\{1\}}} \log N(\mathfrak{p}) = \frac{1}{|G|} \sum_{N(\mathfrak{p}) \leq x} \log N(\mathfrak{p}) + O(x^{\sigma+\varepsilon})$$

holds. Here the sums are over unramified prime ideals of \mathfrak{o}_K , and $N(\mathfrak{p}) := |\mathfrak{o}_K/\mathfrak{p}|$ is the absolute norm of \mathfrak{p} . Then, the Artin L -functions $L(s, \chi)$ associated to the non-trivial irreducible characters of G are analytic in the half-plane H_σ and have no zeros there. Consequently, for any conjugacy class C of G we have

$$\sum_{\substack{N(\mathfrak{p}) \leq x \\ \text{Frob}(\mathfrak{p})=C}} \log N(\mathfrak{p}) = \frac{|C|}{|G|} \sum_{N(\mathfrak{p}) \leq x} \log N(\mathfrak{p}) + O(x^{\sigma+\varepsilon}).$$

Remark 1.1. If $H \triangleleft G$ is a normal subgroup, then the above result applies equally well to the Galois extension L^H/K in place of L/K . It follows that if the density of prime ideals in K whose Frobenius lies in H tends to $|H|/|G|$ with a power saving error term, then the density of prime ideals in K whose Frobenius lies in CH tends to $|CH|/|G|$ with the same power saving error term. We are grateful to the referee for pointing out this formal generalization.

Lucia’s observation pertains to the special case when $K = \mathbb{Q}$ and $L = \mathbb{Q}(e^{2\pi i/q})$. In the next section we shall formulate a more precise result from which the corollary above will follow. As we shall see, the phenomenon admits a clean explanation by Heilbronn characters [7, 9], and leads to a transparent proof of the density theorem itself. In particular, we shall use a result of Foote and Murty [8, Section 3, Proposition] (see [7, Proposition 2.1]), which refines Aramata’s theorem [1] (see [10, Chapter XVIII, Theorem 8.4]) and ultimately relies on Artin’s reciprocity law [2] (see [12, Chapter VI, Theorem 5.5]).

2 Statement of the result

As above, let L/K be a Galois extension of number fields with Galois group $G := \text{Gal}(L/K)$. For each conjugacy class $C \subset G$, consider the Dirichlet series

$$F(s, C) := \sum_{\text{Frob}(\mathfrak{p})=C} \frac{\log N(\mathfrak{p})}{N(\mathfrak{p})^s} - \frac{|C|}{|G|} \sum_{\mathfrak{p}} \frac{\log N(\mathfrak{p})}{N(\mathfrak{p})^s},$$

which converges absolutely in the half-plane H_1 . Here \mathfrak{p} runs through the unramified prime ideals of \mathfrak{o}_K , and $N(\mathfrak{p}) := |\mathfrak{o}_K/\mathfrak{p}|$ is the absolute norm of \mathfrak{p} . Chebotarev’s density theorem, in its weakest analytic form, states that $F(s, C)$ tends to a finite limit as $s \rightarrow 1+$.

For our purposes, $F(s, C)$ is essentially the same as

$$H(s, C) := -\frac{|C|}{|G|} \sum_{\substack{\chi \in \text{Irr}(G) \\ \chi \neq \chi_0}} \frac{L'}{L}(s, \chi) \bar{\chi}(g_C), \quad s \in H_1,$$

where χ runs through the non-trivial irreducible characters of G . Here $L(s, \chi)$ is the Artin L -function of χ , and g_C is any element of the conjugacy class C . Indeed, the Schur orthogonality relation [10, Chapter XVIII, Theorem 5.5] shows that $F(s, C) - H(s, C)$ is represented by an absolutely convergent Dirichlet series in $H_{1/2}$, hence it extends to a bounded analytic function in H_σ for any $\sigma > 1/2$.

Recall Brauer’s theorem [3, Theorem 1] (see [10, Chapter XVIII, Theorem 10.13]), which shows that each $L(s, \chi)$ is meromorphic on \mathbb{C} . Hence $H(s, C)$ is meromorphic on \mathbb{C} with simple poles, and it follows that $F(s, C)$ extends meromorphically to $H_{1/2}$ with (at most) simple poles. A natural question arises: where are the poles and how large are the corresponding residues? Chebotarev’s density theorem tells us that $s = 1$ is not a pole. Artin’s conjecture and the Riemann hypothesis for Artin L -functions imply that $F(s, C)$ has no pole at all in the half-plane $H_{1/2}$.

Our result relates the poles and residues of $F(s, C)$ in the half-plane $H_{1/2}$ to the zeros of $\zeta_L(s)/\zeta_K(s)$ in $H_{1/2}$. Since L/K is Galois, Aramata’s theorem shows that $\zeta_L(s)/\zeta_K(s)$ is an entire function. Since $\zeta_K(s)$ and $\zeta_L(s)$ both have simple poles at $s = 1$, we know moreover that $\zeta_L(s)/\zeta_K(s)$ is non-zero at $s = 1$. Thus the result stated below can be read as a supplement to (or refinement of) Chebotarev’s density theorem.

Theorem. For any point $s_0 \in H_{1/2}$, the following statements are equivalent:

- (a) s_0 is a zero of $\zeta_L(s)/\zeta_K(s)$;
- (b) s_0 is a pole of $F(s, \{1\})$;
- (c) s_0 is a pole of $F(s, C)$ for some conjugacy class $C \subset G$;
- (d) s_0 is a zero or pole of $L(s, \chi)$ for some non-trivial $\chi \in \text{Irr}(G)$;
- (e) s_0 is a zero of $L(s, \chi)$ for some non-trivial $\chi \in \text{Irr}(G)$.

Moreover,

$$\sum_C \frac{|G|}{|C|} \left| \text{res}_{s=s_0} F(s, C) \right|^2 = \sum_{\substack{\chi \in \text{Irr}(G) \\ \chi \neq \chi_0}} \left(\text{ord}_{s=s_0} L(s, \chi) \right)^2 \leq \left(\text{ord}_{s=s_0} \zeta_L(s) \right)^2 - \left(\text{ord}_{s=s_0} \zeta_K(s) \right)^2. \tag{2.1}$$

3 Proof of the Theorem

We begin by proving the key relation (2.1). By our initial remarks,

$$\text{res}_{s=s_0} F(s, C) = \text{res}_{s=s_0} H(s, C) = -\frac{|C|}{|G|} \sum_{\substack{\chi \in \text{Irr}(G) \\ \chi \neq \chi_0}} \left(\text{ord}_{s=s_0} L(s, \chi) \right) \bar{\chi}(g_C), \tag{3.1}$$

hence,

$$\begin{aligned} \sum_C \frac{|G|}{|C|} \left| \text{res}_{s=s_0} F(s, C) \right|^2 &= \sum_C \frac{|C|}{|G|} \left| \sum_{\substack{\chi \in \text{Irr}(G) \\ \chi \neq \chi_0}} \left(\text{ord}_{s=s_0} L(s, \chi) \right) \bar{\chi}(g_C) \right|^2 \\ &= \frac{1}{|G|} \sum_{g \in G} \left| \sum_{\substack{\chi \in \text{Irr}(G) \\ \chi \neq \chi_0}} \left(\text{ord}_{s=s_0} L(s, \chi) \right) \bar{\chi}(g) \right|^2. \end{aligned}$$

Now the first part of (2.1) follows from the Schur orthogonality relation [10, Chapter XVIII, Theorem 5.2], and the second part of (2.1) is a result of Foote and Murty [8, Section 3, Proposition] (see [7, Proposition 2.1]) upon noting that $\zeta_K(s) = L(s, \chi_0)$. See also Remark 3.2 below.

The rest of the Theorem is straightforward. The implication (a) \Rightarrow (b) follows from the factorization

$$\zeta_L(s)/\zeta_K(s) = \prod_{\substack{\chi \in \text{Irr}(G) \\ \chi \neq \chi_0}} L(s, \chi)^{\chi(1)}. \tag{3.2}$$

Indeed, if s_0 is a zero of this meromorphic function, then it is a pole of the logarithmic derivative of the function, which is $-|G|H(s, \{1\})$. Hence s_0 is a pole of $H(s, \{1\})$, and of $F(s, \{1\})$ as well.

The implication (b) \Rightarrow (c) is trivial.

The identity (3.1), or alternatively the first part of (2.1), ensures that if some $F(s, C)$ has a pole at s_0 , then $\text{ord}_{s=s_0} L(s, \chi)$ must be non-zero for some non-trivial $\chi \in \text{Irr}(G)$. Thus (c) implies (d).

If $L(s, \chi)$ has a zero or pole at s_0 for some non-trivial $\chi \in \text{Irr}(G)$, then the second part of (2.1) shows that the right-hand side of (2.1) is strictly positive. Hence s_0 is a zero of $\zeta_L(s)/\zeta_K(s)$, and by (3.2) it is a zero of $L(s, \chi)$ for some non-trivial $\chi \in \text{Irr}(G)$ as well. Thus (d) implies (e), and the same argument also gives that (e) implies (a).

The proof of the Theorem is complete.

Remark 3.1. There are other natural ways to see the equivalence of the five statements in the Theorem. For example, (2.1) can be perceived as the analytic embodiment of $(c) \Leftrightarrow (d) \Rightarrow (a)$. We have seen that $(a) \Rightarrow (b)$ follows from (3.2), while $(b) \Rightarrow (e)$ clearly follows from (3.1), and $(e) \Rightarrow (d)$ is trivial.

Remark 3.2. The crucial input in the proof above is the result of Foote and Murty that we quote. This result can be formulated as the bound $\langle \theta_G, \theta_G \rangle_G \leq \theta_G(1)^2$, where

$$\theta_G := \sum_{\chi \in \text{Irr}(G)} \left(\text{ord}_{s=s_0} L(s, \chi) \right) \chi$$

is the Heilbronn character of G relative to s_0 . Foote and Murty [8] assume that $s_0 \neq 1$, and they actually prove that $|\theta_G(g)| \leq \theta_G(1)$ holds for all $g \in G$. This stronger bound relies on two fundamental properties. First, the restriction of θ_G to a subgroup $H \leq G$ equals θ_H . Second, the Heilbronn character of a cyclic group is a character (not just a virtual character) by Artin’s reciprocity law. Indeed, putting $H = \langle g \rangle$, and using the first property, then the second property, and then again the first property, we obtain $|\theta_G(g)| = |\theta_H(g)| \leq \theta_H(1) = \theta_G(1)$. For $s_0 = 1$ the same argument works if we replace θ_G by

$$\theta_G + \chi_0 = \sum_{\substack{\chi \in \text{Irr}(G) \\ \chi \neq \chi_0}} \left(\text{ord}_{s=1} L(s, \chi) \right) \chi.$$

The conclusion is that this modified Heilbronn character is zero, which is equivalent to Chebotarev’s density theorem. Of course Artin’s reciprocity law is also present in our overall discussion via Brauer’s theorem that Artin L -functions are meromorphic.

Remark 3.3. One can also apply Heilbronn’s idea to the familiar function

$$U_G(s, g) := \sum_{\chi \in \text{Irr}(G)} \frac{L'}{L}(s, \chi) \bar{\chi}(g), \quad s \in H_1, \quad g \in G.$$

Namely, by Frobenius reciprocity, the restriction of $U_G(s, *)$ to a subgroup $H \leq G$ equals $U_H(s, *)$. Hence, by Artin’s reciprocity law, $U_G(s, g)$ is meromorphic on \mathbb{C} with simple poles, and it has a simple pole at $s = 1$ with residue -1 . Using also

$$\frac{L'}{L}(s, \chi) = \langle U_G(s, *), \bar{\chi} \rangle,$$

one obtains a weak form of Brauer’s theorem: each $L'(s, \chi)/L(s, \chi)$ is meromorphic on \mathbb{C} with simple poles, and for $\chi \neq \chi_0$, the point $s = 1$ is not a pole. From this, one can derive a slightly weaker version of the main Theorem without using Brauer’s theorem: (a), (b), (c) are equivalent to each other and to the statement that s_0 is a pole of $L'(s, \chi)/L(s, \chi)$ for some non-trivial $\chi \in \text{Irr}(G)$. For a self-contained argument along these lines, see the lecture slides at <https://ntrg.math.unideb.hu/GH2023Talk.pdf>.

4 Deducing the Corollary

Note that

$$F(s, \{1\}) = \int_1^\infty \left(\sum_{\substack{N(\mathfrak{p}) \leq x \\ \text{Frob}(\mathfrak{p}) = \{1\}}} \log N(\mathfrak{p}) - \frac{1}{|G|} \sum_{N(\mathfrak{p}) \leq x} \log N(\mathfrak{p}) \right) \frac{s}{x^{s+1}} dx,$$

so that the hypothesis in the Corollary implies that $F(s, \{1\})$ extends analytically to the half-plane H_σ . By the implication $(d) \Rightarrow (b)$ of our Theorem, it follows that all the Artin L -functions $L(s, \chi)$ with

non-trivial $\chi \in \text{Irr}(G)$ extend analytically to this half-plane H_σ and have no zeros there. Arguing as in the proof of the prime number theorem, it follows that for every non-trivial $\chi \in \text{Irr}(G)$ one has

$$\sum_{N(\mathfrak{p}) \leq x} \chi(\text{Frob}(\mathfrak{p})) \log N(\mathfrak{p}) = O(x^{\sigma+\varepsilon})$$

for any $\varepsilon > 0$. The Schur orthogonality relation [10, Chapter XVIII, Theorem 5.5] gives

$$\sum_{\substack{N(\mathfrak{p}) \leq x \\ \text{Frob}(\mathfrak{p})=C}} \log N(\mathfrak{p}) - \frac{|C|}{|G|} \sum_{N(\mathfrak{p}) \leq x} \log N(\mathfrak{p}) = \frac{|C|}{|G|} \sum_{\substack{\chi \in \text{Irr}(G) \\ \chi \neq \chi_0}} \bar{\chi}(g_C) \sum_{N(\mathfrak{p}) \leq x} \chi(\text{Frob}(\mathfrak{p})) \log N(\mathfrak{p}),$$

and the conclusion of the Corollary follows.

Acknowledgements The first author was supported by the Rényi Intézet Lendület Automorphic Research Group and NKFIH (National Research, Development and Innovation Office) (Grant No. K 143876). The second author was supported in part by a grant from the National Science Foundation, and a Simons Investigator Award from the Simons Foundation. MathOverflow user Lucia is aware of this paper and agrees that it be published in the current state and authorship. The authors are grateful to Lucia. Finally, the authors thank Emmanuel Kowalski and the anonymous referees for valuable comments.

Open access funding provided by ELKH Alfréd Rényi Institute of Mathematics.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made.

The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.

To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- 1 Aramata H. Über die Teilbarkeit der Dedekindschen Zetafunktionen. *Proc Imp Acad Tokyo*, 1933, 9: 31–34
- 2 Artin E. Beweis des allgemeinen Reziprozitätsgesetzes. *Abh Math Sem Univ Hamburg*, 1927, 5: 353–363
- 3 Brauer R. On Artin's L -series with general group characters. *Ann of Math (2)*, 1947, 48: 502–514
- 4 Chebotarev N. Détermination de la densité des nombres premiers appartenants à une classe donnée de substitutions, I (in Russian). *Bull Acad Sci Russie*, 1923, 17: 205–230
- 5 Chebotarev N. Détermination de la densité des nombres premiers appartenants à une classe donnée de substitutions, II (in Russian). *Bull Acad Sci Russie*, 1923, 17: 231–250
- 6 Chebotarev N. Die Bestimmung der Dichtigkeit einer Menge von Primzahlen, welche zu einer gegebenen Substitutionsklasse gehören. *Math Ann*, 1925, 95: 191–228
- 7 Foote R, Ginsberg H, Murty V K. Heilbronn characters. *Bull Amer Math Soc*, 2015, 52: 465–496
- 8 Foote R, Murty V K. Zeros and poles of Artin L -series. *Math Proc Cambridge Philos Soc*, 1989, 105: 5–11
- 9 Heilbronn H. On real zeros of Dedekind ζ -functions. *Canadian J Math*, 1973, 25: 870–873
- 10 Lang S. *Algebra*, revised 3rd ed. Graduate Texts in Mathematics, vol. 211. New York: Springer-Verlag, 2002
- 11 MathOverflow. Is there a von Koch-type theorem for the generalized Riemann hypothesis? <https://mathoverflow.net/q/181241>
- 12 Neukirch J. *Algebraische Zahlentheorie*. Berlin: Springer-Verlag, 1992