

# THE INFIMUM OF THE VOLUMES OF CONVEX POLYTOPES OF ANY GIVEN FACET AREAS IS 0

N. V. ABROSIMOV\*, E. MAKAI, JR.\*\* , A. D. MEDNYKH\* ,  
YU. G. NIKONOROV\* , G. ROTE

ABSTRACT. We prove the theorem mentioned in the title for  $\mathbb{R}^n$  where  $n \geq 3$ . The case of the simplex was known previously. Also the case  $n = 2$  was settled, but there the infimum was some well-defined function of the side lengths. We also consider the cases of spherical and hyperbolic  $n$ -spaces. There we give some necessary conditions for the existence of a convex polytope with given facet areas and some partial results about sufficient conditions for the existence of (convex) tetrahedra.

2010 Mathematical Subject Classification: 52B11 (primary), 52A38, 52A55 (secondary).

Keywords and phrases: convex polytopes, volume, Euclidean, spherical, hyperbolic spaces.

Studia Scientiarum Mathematicarum Hungarica 51 (4) (2014), 466-519

DOI: 10.1556/SSc.Math.2014.1292

<http://arxiv.org/abs/1304.6579v3>

## 1. PRELIMINARIES

Minimum-area convex polygons with given side lengths are characterized by the following theorem of Böröczky–Kertész–Makai, Jr.

**Theorem A** ([11]). *Let  $m \geq 3$  and  $s_m \geq s_{m-1} \geq \dots \geq s_1 > 0$  and  $s_m < s_{m-1} + \dots + s_1$ . Then the infimum of the areas of convex  $m$ -gons in  $\mathbb{R}^2$  that have side lengths  $s_i$  equals the following number  $A$ . This number  $A$  is the minimum area of all triangles with side lengths  $\sum_{i \in I_1} s_i$ ,  $\sum_{i \in I_2} s_i$ ,  $\sum_{i \in I_3} s_i$ . The minimum is taken over all partitions  $\{I_1, I_2, I_3\}$  of  $\{1, \dots, m\}$  into non-empty parts for which the three resulting side lengths satisfy the non-strict triangle inequality. If the cyclic order of the sides is fixed then an analogous statement holds, where the sides with indices in each of the sets  $I_1, I_2, I_3$  form an arc of the polygonal curve.*

When we investigate simple polygons instead of convex polygons, we have the following result, due to Böröczky–Kertész–Makai, Jr. and Nikonorova.

**Theorem B** ([11, 42]). *Let  $m \geq 3$  and  $s_m \geq s_{m-1} \geq \dots \geq s_1 > 0$  and  $s_m < s_{m-1} + \dots + s_1$ . Then the infimum of the areas of simple  $m$ -gons in  $\mathbb{R}^2$  that have side lengths  $s_i$  equals the following number  $B$ . This number  $B$  is the minimum area*

---

\* The project was supported in part by the State Maintenance Program for the Leading Scientific Schools of the Russian Federation (Grant NSH-921.2012.1) and by the Federal Target Grant “Scientific and educational personnel of innovative Russia” for 2009–2013 (agreement no. 8206, application no. 2012-1.1-12-000-1003-014).

\*\* Research (partially) supported by Hungarian National Foundation for Scientific Research, grant nos. K68398, K75016, K81146.

of all triangles with side lengths  $\sum_{i \in I_1} \varepsilon_i s_i$ ,  $\sum_{i \in I_2} \varepsilon_i s_i$ ,  $\sum_{i \in I_3} \varepsilon_i s_i$ . The minimum is taken over all partitions  $\{I_1, I_2, I_3\}$  of  $\{1, \dots, m\}$  into non-empty parts, and all signs  $\varepsilon_1, \dots, \varepsilon_m$ , for which the three resulting side lengths are non-negative and satisfy the non-strict triangle inequality.

Moreover, if this minimum is not 0 then we may additionally suppose the following. For each  $j \in \{1, 2, 3\}$  the sum  $\sum_{i \in I_j} \varepsilon_i s_i$  cannot be written as  $\sum_{i \in I'_j} \varepsilon_i s_i + \sum_{i \in I''_j} \varepsilon_i s_i$  where  $\{I'_j, I''_j\}$  is a partition of  $I_j$  and where these partial summands are both positive.

We remark that the proofs in the two papers were different. Moreover, in [42] the result is formulated in a special case only, but all ingredients of the proof of the general case are present in [42] as well.

In our paper we write  $\mathbb{R}^n$ ,  $\mathbb{H}^n$  and  $\mathbb{S}^n$  for the *Euclidean*, *hyperbolic* and *spherical*  $n$ -space, respectively. Theorems A and B extend to  $\mathbb{S}^2$  and  $\mathbb{H}^2$  as follows.

**Theorem C** ([11]). *Let  $m \geq 3$  and  $s_m \geq s_{m-1} \geq \dots \geq s_1 > 0$  and  $s_m < s_{m-1} + \dots + s_1$ . Rather than  $\mathbb{R}^2$  we consider  $\mathbb{H}^2$  and  $\mathbb{S}^2$ , but in case of  $\mathbb{S}^2$  we additionally suppose  $\sum_{i=1}^m s_i \leq \pi$ . Then in both cases the word-for-word analogues of Theorems A and B hold for  $\mathbb{H}^2$  and  $\mathbb{S}^2$ .*

In each of these three theorems the question of finding the infimum is reduced to finding the minimum of a set of non-negative numbers whose cardinality is bounded by a function of  $m$ . In Theorems A and B, this bound is  $3^m$  and  $6^m$ , respectively. In Theorem A with given cyclic order of the sides, the bound is  $\binom{m}{3}$ . In Theorem C, the bounds are the same as for Theorems A and B.

Böröczky et al. [11] posed the question whether it is possible to extend these theorems to dimensions  $n \geq 3$ . Their conjecture was that, analogously to the two-dimensional case, the solutions would be given as the volumes of some simplices. Unfortunately, they were unaware of the fact that the case of simplices already had long ago been solved, namely in 1938, as we will describe below.

The analogous problem about the maximal volume of simplices with given facet areas (an isoperimetric-type problem) was solved by Lagrange [35] in 1773 for  $\mathbb{R}^3$  and by Borchardt [9] in 1866 for  $\mathbb{R}^n$ . A simplex is called *orthocentric* if it has an orthocentre, i.e., a common point of all altitudes. It can be characterized also as a simplex where any two disjoint edges (or, equivalently, any two disjoint faces of dimension at least 1) are orthogonal. For this reason, an orthocentric simplex is sometimes also called *orthogonal*, although orthocentric is the presently used terminology. For a relatively recent exposition of the above-mentioned facts and some other properties of orthocentric simplices, see for example [24]. See also the recent paper [20], whose first part is a comprehensive survey about orthocentric simplices. There it is also stressed that for many elementary geometrical theorems the objects in  $\mathbb{R}^n$  corresponding to triangles are not the general simplices but just the orthocentric ones.

**Theorem D** ([35, 9]). *Let  $n \geq 3$ . Then among the simplices in  $\mathbb{R}^n$  with given facet areas  $S_{n+1} \geq \dots \geq S_1 > 0$  (if such simplices exist) there exists (up to congruence) exactly one simplex of maximal volume. It is also the (up to congruence) unique orthogonal simplex with these facet areas.*

Unaware of the above-mentioned solution of the maximum problem, A. Narasinga Rao [50] posed the following problem in 1937:

“The areas of the four facets of a tetrahedron are  $\alpha, \beta, \gamma, \delta$ . Is the volume determinate? If not, between what limits does it lie?”

This problem was soon solved independently by Venkatachaliengar, Iyengar, Auluck, and Iyengar–Iyengar [61, 30, 6, 31]. In fact, under the above hypothesis the volume is not determined (supposing that such tetrahedra exist). Moreover, they reproved that there is up to congruence exactly one tetrahedron of maximal volume with the given facet areas, which is also orthogonal — and they reproved that it is also the unique orthocentric tetrahedron with these facet areas. Moreover, they proved that there exists a tetrahedron with the given face areas that has an arbitrarily small volume. A generalization of their first mentioned result to multi-dimensional Euclidean spaces was obtained in [61, 30, 31]. We cite only their statement about the infimum of the volumes.

**Theorem E** ([61, 30, 6, 31]). *Let  $n \geq 3$  and  $S_{n+1} \geq S_n \geq \dots \geq S_1 > 0$ . Then there exists a simplex in  $\mathbb{R}^n$  with these  $(n-1)$ -volumes of the facets if and only if*

$$S_{n+1} < S_1 + S_2 + \dots + S_n.$$

*If this inequality holds, then, for any  $\varepsilon > 0$ , there is a nondegenerate simplex in  $\mathbb{R}^n$  with facet areas  $S_1, S_2, \dots, S_{n+1}$  and volume at most  $\varepsilon$ .*

The proof of Theorem E by Iyengar and Iyengar [31] was based on the following statement, which is valid for simplices only [31, p. 306]. Let  $T$  be a simplex with facet areas  $S_1, S_2, \dots, S_{n+1}$  and respective outer unit facet normals  $u_1, \dots, u_{n+1}$ . Then we have  $\sum_{i=1}^{n+1} S_i u_i = 0$ . Let us consider an  $(n+1)$ -gon  $P \subset \mathbb{R}^n$  with side vectors  $S_1 u_1, \dots, S_{n+1} u_{n+1}$ . Its convex hull  $T'$  is then also a simplex, whose volume is invariant under permutations of the side vectors of  $P$ . Moreover, for the volumes of  $T$  and  $T'$ , we have  $V(T)^{n-1} = V(T')[(n-1)!]^2/n^{n-2}$ . Based on this relation and some calculations, Iyengar and Iyengar could make  $V(T')$  arbitrarily small. However, this can also be done by choosing  $u_1, \dots, u_{n+1}$  in a small neighbourhood of the  $x_1 \dots x_{n-1}$ -coordinate hyperplane. See also the first and third proofs of our Theorem 2.

The question of maximal volume of polytopes with given facet areas is much less understood.

For non-degenerate polytopes in  $\mathbb{R}^n$  with given facet areas and given facet outer unit normals, we have the following result of Brunn [14], see also [41, §10.5]. The maximal volume is attained for the (up to translation) unique *convex* polytope with these given facet areas and given facet outer unit normals. (For coinciding facet outer unit normals, one has to add their areas.) This result was rediscovered in [10, Theorems 2 and 3] and applied to solve another problem. In crystallography, this set of maximal volume is called the *Wulff shape* [62]. It minimizes total surface energy of the crystal and is always convex. For a nice description of the interplay of mathematics and crystallography see [12, §10.11].

For any given number  $m$  of facets and fixed total surface area, the polytope of largest volume has an inball and the facets must touch the inball at their centroids (Lindelöf’s theorem, see [56, p. 43] or [22, II.4.3, p. 264; English ed. IX.43, p. 283].

Now let us restrict our attention to  $\mathbb{R}^3$ . L. Fejes Tóth [21, Theorem 1, p. 175] (see also [22, II.4.3, p. 265, Satz; English ed. IX.43, p. 283, Theorem]) asserts that among (convex) polyhedra with given surface area and  $m = 4, 6,$  and  $12$  faces, the largest volume is attained for the regular tetrahedron, cube, and regular dodecahedron, respectively. He gave a bound on the maximum volume [21, p. 175], valid for each  $m \geq 4$ , which is also asymptotically sharp for  $m \rightarrow \infty$ . For  $m = 5$ , the extremal polyhedron is the regular triangular prism that has an inball [56, p. 41]. A recent complete and simple proof of this fact is given in [25, Theorem 5.10]. However, for  $m = 8$  and  $m = 20$ , the extremal polyhedron is not the regular octahedron and icosahedron, respectively [26, p. 234]. For more information about this isoperimetric problem about convex polyhedra in  $\mathbb{R}^3$  with given number of faces, see the old survey in [26] or the recent survey in the introduction of [57]. For recent numerical results (examples) about the isoperimetric problem for polyhedra, with large symmetry groups, see [36].

A different problem is to maximize the volume enclosed by a given surface that may be bent (isometrically) but not stretched. A theorem of S. P. Olovianishnikoff says the following. For convex bodies  $P, Q \subset \mathbb{R}^3$ , where  $P$  is a convex polyhedron, any mapping of  $\partial P$  to  $\partial Q$  preserving the geodesic distance of every pair of points of  $\partial P$  (i.e., the length of the shortest arc in  $\partial P$  joining these points) extends to an isometry of  $\mathbb{R}^3$ . See [3, Ch. 3, §3, 2, p. 150, Satz 1] for a special case, and S. P. Olovianishnikoff [43], p. 441, Theorem for the general case described above. For convex bodies  $P, Q \subset \mathbb{R}^3$  where  $\partial P$  is of class  $C^2$ , the analogous theorem holds. See [2, Ch. 8, §5, p. 337] for a special case and A. V. Pogorelov [46, Introduction, §1, A, p. 8, Theorem 1, and Ch. 3, 3, p. 66, Theorem 1] for the general case described above.

However, this uniqueness theorem does not say that this unique convex polyhedron would have the maximal volume. The opposite is true: the surface of *every* (not necessarily convex) polyhedron can be isometrically deformed to increase the enclosed volume [44]. For example, the cube can be “blown up”: the face centers move outwards and the vertices move closer to the center. The face diagonals maintain their original length, but the original edges of the cube are longer than necessary: they become crumpled, with wrinkles perpendicular to the original edge. Globally, the polyhedron becomes more “ball-like”. This volume-increasing phenomenon for convex bodies was first observed by A. V. Pogorelov in the theory of thin shells in mechanics [47, 48]. (A short summary of the results of [47] and of some other related results is given in [49].) An animation showing a deformation of the cube with a volume increase by a factor of about 1.2567 has been produced by Buchin and Schulz [15]. The problem of enclosing the largest volume with the surface of a given convex polyhedron, possibly under the constraint of preserving the original symmetries, has been treated in many papers [58, 59, 38, 16, 8, 4, 44, 39] (“inextensional” in the title of [58] means “isometric w.r.t. the geodesic distance”). (According to a private communication from the second author of [39], in the tableau summarizing the numerical results in pp. 154 and 181, the values in the middle column for the dodecahedron and the icosahedron are not correct. They are actually smaller than the values in the third column, which are proved in [39], and those are the best published values.) For a recent survey on this and related questions see [51].

**Notations.** In this paper,  $V(\cdot)$  denotes volume of a set,  $S(\cdot)$  its surface area,  $\text{diam}(\cdot)$  its diameter,  $\text{aff}(\cdot)$  its affine hull,  $\text{lin}(\cdot)$  its linear hull, and  $\partial(\cdot)$  its boundary. If we want to indicate also the dimension  $n$  then we will write  $V_n(\cdot)$  for the  $n$ -volume. Sometimes we will refer to the  $(n-1)$ -volume in  $\mathbb{R}^n$ ,  $\mathbb{H}^n$  or  $\mathbb{S}^n$  as *area*. We write  $\kappa_n$  for the volume of the unit ball in  $\mathbb{R}^n$ . For  $x, y$  in  $\mathbb{R}^n$ ,  $\mathbb{H}^n$  or  $\mathbb{S}^n$ , we write  $[x, y]$  for the segment and  $\ell(x, y)$  for the line joining  $x$  and  $y$ . On  $\mathbb{S}^n$ ,  $x$  and  $y$  must not be antipodes, and we mean by  $[x, y]$  the minor arc on the great circle through  $x$  and  $y$ . The line  $\ell(x, y)$  is well-defined only for  $x \neq y$  — writing  $\ell(x, y)$  we suppose  $x \neq y$ . We denote the distance between  $x$  and  $y$  by  $|xy|$ .

For standard facts about convex bodies we refer to [55].

## 2. NEW RESULTS

**2.1. Euclidean Space.** The following theorem can be considered as folklore, but we could not locate a proof. For completeness, we state and prove it.

**Theorem 1.** *Assume that  $m > n \geq 3$  are integers, and consider any sequence of numbers  $S_m \geq S_{m-1} \geq \dots \geq S_1 > 0$ . Then the following statements are equivalent:*

- (i) *There exists a non-degenerate polytope  $P \subset \mathbb{R}^n$  with  $m$  facets and with facet areas  $S_1, S_2, \dots, S_m$ .*
- (ii) *There exists a non-degenerate convex polytope  $P \subset \mathbb{R}^n$  with  $m$  facets and with facet areas  $S_1, S_2, \dots, S_m$ .*
- (iii) *The inequality  $S_m < S_1 + S_2 + \dots + S_{m-1}$  holds.*

*If we also allow degenerate polytopes in (i) or (ii), then they imply, rather than (iii),*

- (iii')  *$S_m \leq S_1 + \dots + S_{m-1}$  with equality if and only if the polytope degenerates into the doubly counted facet with area  $S_m$ .*

**Theorem 2.** *Let  $m > n \geq 3$  be integers. Let  $\varepsilon > 0$  and  $S_m \geq S_{m-1} \geq \dots \geq S_1 > 0$  be a sequence of numbers such that  $S_m < S_1 + S_2 + \dots + S_{m-1}$ . Then there exists a non-degenerate convex polytope  $P \subset \mathbb{R}^n$  with  $m$  facets and with facet areas  $S_1, S_2, \dots, S_m$  and with volume  $V(P) \leq \varepsilon$ .*

*Remark 1.* This theorem shows that for dimension  $n \geq 3$  there are no separate questions for convex and general polytopes. Recall that for dimension  $n = 2$  these questions had different answers, see Theorems A and B.

We give three different proofs of Theorem 2. The first one is independent of Theorem D and reproves the case of the simplex. It is an existence proof by contradiction. The second proof uses Theorem D. It reduces the question to the case of simplices. Both proofs rely on delicate convergence arguments (see Sections 3 and 4). The third proof is geometric. It constructs examples with small volumes that are like “needles”. In particular we will give an explicit upper bound for the volumes of our examples in terms of the “steepness” of their facets (Lemmas 2 and 4). If we consider  $n, m$  and the facet areas as fixed then our estimate is sharp up to a constant factor (see Lemma 4).

Note that there is a very interesting dichotomy. In Theorems A and B for  $\mathbb{R}^2$  (and also in Theorem C for  $\mathbb{H}^2$  and  $\mathbb{S}^2$ ) we have some definite functions of the side lengths as infima. In Theorem 2 for  $\mathbb{R}^n$  with  $n \geq 3$  the infimum does not depend at all on the facet areas.

**2.2. Hyperbolic Space.** For the hyperbolic case we have a word-for-word analog of the implications (ii) $\implies$ (iii) and (ii) $\implies$ (iii') from Theorem 1 (under the respective hypotheses).

**Proposition 1.** *Let  $P \subset \mathbb{H}^n$  be a polytope with facet areas  $S_m \geq S_{m-1} \geq \dots \geq S_1 > 0$ . Then the inequality  $S_m \leq \sum_{i=1}^{m-1} S_i$  holds, with equality if and only if  $P$  degenerates into the doubly counted facet with area  $S_m$ .*

Next we give two statements that show the following. The necessary condition in Proposition 1 together with the inequalities  $S_i \leq \pi$  is not sufficient even for the existence of a tetrahedron in  $\mathbb{H}^3$  with these facet areas. That is, there are some further necessary conditions. Recall that the area of a simple  $k$ -gon in  $\mathbb{H}^2$  is bounded by  $(k-2)\pi$ .

**Proposition 2.** *Let us admit polyhedra in  $\mathbb{H}^3$  whose vertices are all distinct but which possibly have some infinite vertices. Then a non-degenerate polyhedron with facet areas  $S_m, S_{m-1}, \dots, S_3$  maximal (i.e.,  $(k-2)\pi$  for a  $k$ -gonal face) but with facet areas  $S_2, S_1$  not maximal does not exist.*

Proposition 2 would suggest that for polyhedra in  $\mathbb{H}^3$ , if all facets but two have areas nearly maximal (i.e., close to  $(k-2)\pi$  for a  $k$ -gonal face) then the same statement would hold for the remaining two facets as well. However, this is not true. Even in the convex case, these two facets can have areas close to 0, as shown by the following example. Consider a very large circle in  $\mathbb{H}^2 \subset \mathbb{H}^3$  and a regular  $l$ -gon  $p_1 \dots p_l$  inscribed in it ( $l \geq 3$ ). Choose  $p_{l+1}$  on our circle with  $|p_l p_{l+1}| = \varepsilon$ . Then all triangles with vertices among the  $p_i$ 's have areas close to  $\pi$  except those that contain both  $p_l$  and  $p_{l+1}$ , and those have very small areas. Now perturb these points  $p_i$  a little bit in  $\mathbb{H}^3$  so that no four lie in a plane. Then their convex hull is a triangle-faced convex polyhedron, and the perturbation of the segment  $[p_l, p_{l+1}]$  is an edge of it. (To see this, use the collinear model. For any convex polygon with strictly convex angles, its edges will remain edges of the convex hull after a sufficiently small perturbation.) The two facets of our polyhedron incident to this edge have very small areas while all other facets have areas close to  $\pi$ , i.e., are nearly maximal.

However, an analogous statement for all but one facets will be shown in the convex case.

**Proposition 3.** *Assume that we have a convex polyhedron in  $\mathbb{H}^3$  with infinite vertices admitted. Suppose its  $m$  facets are a  $k_m$ -gon,  $\dots$ ,  $k_1$ -gon and have respective areas  $S_m \geq \dots \geq S_2 \geq S_1 > 0$ . Then for any  $i \in \{1, \dots, m\}$  we have*

$$(k_i - 2)\pi - S_i \leq \sum_{\substack{1 \leq j \leq m \\ j \neq i}} ((k_j - 2)\pi - S_j).$$

*If there is a finite vertex whose incident edges do not lie in a plane, then the above inequality is strict.*

In §6 Remark 9, it will be explained that, in a sense, there are no interesting analogues of Proposition 3 for  $\mathbb{R}^3$  and  $\mathbb{S}^3$ .

Now we turn to sufficient conditions for the existence of hyperbolic tetrahedra.

**Theorem 3.** *Assume that  $\pi/2 > S_4 \geq S_3 \geq S_2 \geq S_1 > 0$ ,  $S_4 < S_1 + S_2 + S_3$ , and one of the inequalities*

$$\tan(S_1/2) > \frac{1 - \cos S_4}{2\sqrt{\cos S_4}}, \quad (1)$$

or

$$S_4 \geq S_3 + S_2 \quad (2)$$

holds. Then there exists a non-degenerate tetrahedron  $T \subset \mathbb{H}^3$  with facet areas  $S_1, S_2, S_3, S_4$ .

**2.3. Spherical Space.** For the spherical case, we give some necessary and some sufficient conditions for the existence.

We say that a set  $X \subset \mathbb{S}^n$  (for  $n \geq 2$ ) is *convex* if, for any two non-antipodal  $x, y \in X$ , the connecting minor great- $\mathbb{S}^1$  arc  $[x, y]$  also belongs to  $X$ . This definition classifies an antipodal pair of points as a convex set. But these are the only convex sets which are disconnected, and since the sets we consider contain non-trivial arcs, these exceptional cases play no role for us. By a *nondegenerate simplex in  $\mathbb{S}^n$*  we mean the set of those points of  $\mathbb{S}^n$  that have non-negative coordinates in some (non-orthogonal) coordinate system with origin at 0, with its usual face lattice. A *simplex in  $\mathbb{S}^n$*  is a nondegenerate simplex, or a limiting position of nondegenerate simplices. Thus, for example, we will not consider concave spherical triangles or spherical triangles with sides  $3\pi/2, \pi/4, \pi/4$  or  $2\pi, 0, 0$ , but a spherical triangle with angles  $\pi, \pi/5, \pi/5$  and sides  $\pi, \pi/3, 2\pi/3$  is a (degenerate) simplex. As a point set, this simplex is indistinguishable from a digon. A different division of the digon side, like  $\pi, \pi/4, 3\pi/4$ , is regarded as a different simplex. To emphasize the fact that we do not just regard a simplex as a point set but we consider its face structure, we will often refer to it as a *combinatorial simplex*. All simplices in  $\mathbb{S}^n$ , as well as in  $\mathbb{R}^n$  and  $\mathbb{H}^n$ , are convex. A simplex in an open half- $\mathbb{S}^n$  with non-empty interior is always nondegenerate. (Observe that an open half- $\mathbb{S}^n$  also has a collinear model in  $\mathbb{R}^n$  that also respects convexity. For the open southern half- $\mathbb{S}^n$  in  $\mathbb{R}^{n+1}$  consider the central projection to the tangent space  $\mathbb{R}^n$  at the South Pole.)

**Proposition 4.** *Let  $P \subset \mathbb{S}^n$  be a polytope with facet areas  $S_m \geq \dots \geq S_1 > 0$ , such that each facet lies in some closed half- $\mathbb{S}^{n-1}$ . Then*

$$S_m \leq S_1 + \dots + S_{m-1}.$$

Here strict inequality holds if  $P$  is contained in an open half- $\mathbb{S}^n$  and does not degenerate into the doubly-counted facet with area  $S_m$ .

If  $P$  is a convex polytope contained in some closed half- $\mathbb{S}^n$ , then

$$\sum_{i=1}^m S_i \leq V_{n-1}(\mathbb{S}^{n-1}).$$

Here strict inequality holds if  $P$  is contained in an open half- $\mathbb{S}^n$ .

*Remark 2.* Clearly, in the first part of Proposition 4, the hypothesis that each facet lies in some closed half- $\mathbb{S}^n$  cannot be dispensed. Already for  $n = 2$ , we may even have a degenerate combinatorial simplex lying in some great- $\mathbb{S}^{n-1}$  with one facet strictly containing a half- $\mathbb{S}^{n-1}$ . In the second part of Proposition 4, if  $P$  is contained in a closed half- $\mathbb{S}^n$  but not in an open half- $\mathbb{S}^n$ , then equality can occur:  $P$  can

degenerate so that one facet is a closed half- $\mathbb{S}^{n-1}$ , and the union of the other facets is this closed half- $\mathbb{S}^{n-1}$  or the closure of its complement in this  $\mathbb{S}^{n-1}$ .

*Remark 3.* We do not know how to algorithmically decide whether a simplex with given facet areas  $S_i$  in  $\mathbb{H}^n$  or  $\mathbb{S}^n$  exists, for  $n \geq 3$ . The main difficulty are the transcendental functions that enter into the calculation of volumes. In  $\mathbb{H}^3$  and  $\mathbb{S}^3$ , however, we have a positive answer to a slightly modified question. The question whether there is a tetrahedron (for  $\mathbb{S}^3$  in the sense described above) with facet areas  $S_1, S_2, S_3, S_4$ , is decidable if we are given  $\tan(S_1/2), \dots, \tan(S_4/2)$  as inputs.

We model this question by setting up a system of equations and inequalities in the unknown coordinates  $(x_{ij})$  of the four vertices. (For  $\mathbb{S}^3$  we use its standard embedding into  $\mathbb{R}^4$ , while for  $\mathbb{H}^3$  we use the hyperboloid model in  $\mathbb{R}^4$ .) The equations express the condition that the vertices lie on  $\mathbb{S}^3$  or  $\mathbb{H}^3$ , and that the facet areas of the corresponding tetrahedron should be  $S_1, S_2, S_3, S_4$ . Further inequalities are necessary for  $\mathbb{S}^3$  to ensure our definition of simplices. We are interested in the set of 4-tuples  $(S_1, S_2, S_3, S_4)$  for which *there exist* coordinate vectors  $(x_{ij})$  that fulfill the conditions. These conditions turn out to be polynomial equations and inequalities (these polynomials having rational coefficients) in  $\tan(S_1/2), \dots, \tan(S_4/2)$  and in the coordinates  $x_{ij}$ . By a fundamental result of Tarski [60], this existence question is therefore (in principle) decidable (we can eliminate the variables  $x_{ij}$ ). More specifically, the set of quadruples  $(\tan(S_1/2), \dots, \tan(S_4/2))$  for all tetrahedra in  $\mathbb{H}^3$  or  $\mathbb{S}^3$  (for  $\mathbb{S}^3$  in our sense) can be described by a finite number of polynomial equalities and inequalities, these polynomials having rational coefficients, also using the usual logical connectives “and”, “or”, “not”. In other words, this set forms a *semi-algebraic set*.

*Remark 4.* For simplices in  $\mathbb{S}^n$  and  $\mathbb{H}^n$  (with finite vertices) the case  $S_1 = 0$  and all other  $S_i$ 's positive and sufficiently small can be described. The description is: there exists a partition of the other facets into two classes such that for the two classes the sums of the facet areas are equal. For this we have to use index considerations, like later in the Proofs of Theorems 3 and 4.

**Theorem 4.** *Assume that  $\pi/2 > S_4 \geq S_3 \geq S_2 \geq S_1 > 0$ ,  $S_4 < S_1 + S_2 + S_3$ , and one of the inequalities*

$$\tan(S_1/2) \geq \frac{1 - \cos S_4}{2\sqrt{\cos S_4}}, \quad (3)$$

or

$$S_4 \geq S_3 + S_2 \quad (4)$$

*holds. Then there exists a non-degenerate (convex) tetrahedron  $T \subset \mathbb{S}^3$  with facet areas  $S_1, S_2, S_3, S_4$ .*

Now we turn to sufficient conditions for the existence of spherical polyhedral complexes. The second statement of Proposition 5 says the following. For combinatorial simplices contained in some closed half- $\mathbb{S}^n$ , the two necessary conditions from Proposition 4 are also sufficient for their existence.

**Proposition 5.** (i) *Let  $n \geq 2$  and  $m \geq 3$  be integers and let  $S_m \geq \dots \geq S_1 > 0$  and  $S_m \leq S_1 + \dots + S_{m-1}$  and  $S_1 + \dots + S_m \leq V_{n-1}(\mathbb{S}^{n-1})$ . Then there exists a convex  $n$ -dimensional polyhedral complex in  $\mathbb{S}^n$  that lies in a closed half- $\mathbb{S}^n$  and has facet areas  $S_1, \dots, S_m$ . All of its facets have two  $(n-2)$ -faces. If*



$S_m < S_1 + \cdots + S_{m-1}$  and  $S_1 + \cdots + S_m < V_{n-1}(\mathbb{S}^{n-1})$  then all its dihedral angles are less than  $\pi$ .

- (ii) Let  $n \geq 2$  and assume  $S_{n+1} \geq \cdots \geq S_1 > 0$ ,  $S_{n+1} \leq S_1 + \cdots + S_n$ , and  $S_1 + \cdots + S_{n+1} \leq V_{n-1}(\mathbb{S}^{n-1})$ . Then there exists a convex polyhedral complex in  $\mathbb{S}^n$  lying in a closed half- $\mathbb{S}^n$  that is a combinatorial  $n$ -simplex with facet areas  $S_1, \dots, S_{n+1}$ . Its faces of any dimension (including the complex itself) have their dihedral angles at most  $\pi$ , and are thus convex, but some of their dihedral angles are equal to  $\pi$  for  $n \geq 3$ .

### 3. TOOLS FOR THE EUCLIDEAN CASE: MINKOWSKI'S THEOREMS

We recall some classical concepts and theorems, which are in essence due to Minkowski, but got their final form by A. D. Aleksandrov [1] and W. Fenchel and B. Jessen [23]. We state the results first for arbitrary convex bodies, and then we restrict them to convex polytopes. We will actually need general convex bodies when considering convergent sequences of convex polytopes in our first two proofs of Theorem 2 (Sections 4.3 and 4.4). The third proof uses only Minkowski's Theorem about convex polytopes (Theorem F'). The reader may want to skip directly to Theorem F'.

A *convex body* in  $\mathbb{R}^n$  is a compact convex set  $K \subset \mathbb{R}^n$  with interior points. For  $x \in \partial K$  we say that  $u \in \mathbb{S}^{n-1}$  is an *outer unit normal vector for  $K$  at  $x$*  if  $\max\{\langle k, u \rangle \mid k \in K\} = \langle x, u \rangle$ . In this section we assume  $n \geq 2$  although the theorems of this section will be applied later for  $n \geq 3$  only.

**Definition 1** (Minkowski, Aleksandrov [1], Fenchel–Jessen [23], see also [55, p. 207, (4.2.24) (with  $\tau(K, \omega)$  defined on p. 77)]). Let  $K \subset \mathbb{R}^n$  be a convex body. The *surface area measure*  $\mu_K$  of  $K$  is a finite Borel measure on  $\mathbb{S}^{n-1}$  defined as follows. For a Borel set  $B \subset \mathbb{S}^{n-1}$ ,  $\mu_K(B)$  is the  $(n-1)$ -dimensional Hausdorff measure of the set  $\{x \in \partial K \mid \text{there is an outer unit normal vector } u \text{ to } K \text{ at } x \text{ such that } u \in B\}$ .

Thus,  $\mu_K$  is an element of  $C(\mathbb{S}^{d-1})^*$ , the dual space of the space of real-valued continuous functions  $C(\mathbb{S}^{d-1})$  on  $\mathbb{S}^{d-1}$ , i.e., the finite signed Borel measures on  $\mathbb{S}^{d-1}$ . We will use the weak\* topology of  $C(\mathbb{S}^{d-1})^*$  as the topology for the finite (signed) Borel measures  $\mu_K$ . That is, convergence of a sequence (or more generally of a net) of finite signed Borel measures  $\mu_\alpha \in C(\mathbb{S}^{d-1})^*$  to a finite signed Borel measure  $\mu \in C(\mathbb{S}^{d-1})^*$  means the following. For each  $f \in C(\mathbb{S}^{n-1})$ , we have  $\int_{\mathbb{S}^{n-1}} f(u) d\mu_\alpha(u) \rightarrow \int_{\mathbb{S}^{n-1}} f(u) d\mu(u)$ . Moreover, since  $\mathbb{S}^{n-1}$  is a compact metric space, the space  $C(\mathbb{S}^{n-1})$  is separable, and hence the weak\* topology of  $C(\mathbb{S}^{n-1})^*$  is metrizable. Therefore, it suffices to give the convergent sequences in it (i.e., it is not necessary to consider nets).

For these elementary concepts and facts from functional analysis, we refer to [19].

**Theorem F** (Minkowski, Aleksandrov [1], Fenchel–Jessen [23], see also [55, p. 389, (7.1.1), pp. 389–390, p. 392, Theorem 7.1.2., p. 397, Theorem 7.2.1]). Let  $n \geq 2$  be an integer and  $K \subset \mathbb{R}^n$  a convex body. The measure  $\mu_K$  defined in Definition 1 is invariant under translations of  $K$  and has the following properties.

- (i)  $\int_{\mathbb{S}^{n-1}} u d\mu_K(u) = 0$ , and
- (ii)  $\mu_K$  is not concentrated on any great- $\mathbb{S}^{n-2}$  of  $\mathbb{S}^{n-1}$ .

Conversely, for any finite Borel measure  $\mu$  on  $\mathbb{S}^{n-1}$  satisfying (i) and (ii), there exists a convex body  $K$  such that  $\mu_K = \mu$ . Moreover, this convex body  $K$  is unique up to translations.

Thus, we can consider the map  $K \mapsto \mu_K$  also as a map  $\{\text{translates of } K\} \mapsto \mu_K$ .

**Theorem G** (Minkowski, Aleksandrov [1], Fenchel–Jessen [23], see also [55, p. 198, Theorem 4.1.1, p. 205, pp. 392–393, proof of Theorem 7.1.2]). *Let  $n \geq 2$  be an integer. Then the mapping  $\{\text{translates of } K\} \mapsto \mu_K$  defined in Definition 1 and just before this theorem is a homeomorphism between its domain and its range. Its domain is the quotient topology of the topology on the convex bodies induced by the Hausdorff metric with respect to the equivalence relation of being translates. Its range is the set of finite Borel measures on  $\mathbb{S}^{n-1}$  satisfying (i) and (ii) of Theorem F with the subspace topology of the weak\* topology on  $C(\mathbb{S}^{n-1})^*$ .*

We have to remark that the cited sources, [55, pp. 392–393, proof of Theorem 7.1.2], as well as [1, proof of the theorem on p. 36, on p. 38], contain explicitly only the proof of the continuity of the bijection  $\{\text{translates of } K\} \mapsto \mu_K$ . However, also the continuity of the inverse map is proved at both places although not explicitly stated. In fact, as kindly pointed out to the authors by R. Schneider, one has to make the following addition to his book [55, proof of Theorem 7.1.2]. Let the sequence of surface area measures  $\mu_{K_i}$  of some convex bodies  $K_i \subset \mathbb{R}^n$  converge to the surface area measure  $\mu_K$  of some convex body  $K \subset \mathbb{R}^n$  in the weak\* topology. Then all  $K_i$ 's have a bounded diameter. This is stated there for polytopes only, but the given proof is valid for all convex bodies. By a translation one can achieve that all  $K_i$ 's and also  $K$  are contained in a fixed ball. Recall that the set of non-empty compact convex sets contained in some closed ball is compact in their usual topology (i.e., that of the Hausdorff metric). Therefore, we can choose a convergent subsequence  $K_{i_j}$  of  $K_i$  with limit  $K'$ , say. In the Note added in proof at the end of the paper we will show that  $K'$  is a convex body. Then the surface area measure  $\mu_{K'}$  of  $K'$  is the weak\* limit of the  $\mu_{K_{i_j}}$ 's, i.e., it equals the originally considered  $\mu_K$ . Hence we have that  $K'$  is a translate of  $K$ . Then the entire sequence  $K_i$  converges to  $K$ . Otherwise, we could choose another subsequence  $K_{i_k}$  converging to another convex body  $K''$ , which is not a translate of  $K$ , with  $\mu_{K''} = \mu_K$ . This is a contradiction. Then also the translation equivalence class of  $K_i$  converges to that of  $K$ , by continuity of the quotient map.

It was also proved by Minkowski that a convex body  $K$  is a convex polytope if and only if  $\mu_K$  (that satisfies (i) and (ii) of Theorem F) has a finite support [55, p. 390, Theorem 7.1.1, also considering p. 397, Theorem 7.2.1]. If the support is  $\{u_1, \dots, u_m\}$ , we may write

$$\mu_K = \sum_{i=1}^m \mu_K(\{u_i\}) \delta(u_i),$$

where  $\delta(u_i)$  is the Dirac measure concentrated at  $u_i$ . (I.e., for a Borel set  $B \subset \mathbb{S}^{n-1}$  we have  $\delta(u_i)(B) = 0 \iff u_i \notin B$  and  $\delta(u_i)(B) = 1 \iff u_i \in B$ .) When we write such an equation, we always assume that  $\mu_K(\{u_i\}) \neq 0$  for all  $i \in \{1, \dots, m\}$ . (Thus, the empty sum means the 0 (finite signed Borel) measure; although for a convex body  $K$ , we have  $\mu_K \neq 0$ .) The weak\* topology restricted to the finite signed Borel measures of finite support, where the support has at most  $m$  elements,

is the following. (We will use only the case when we have a finite Borel measure and (i) and (ii) of Theorem F hold.) For  $u_\alpha, u \in \mathbb{S}^{d-1}$  with  $u_\alpha \rightarrow u$  and for  $c_\alpha, c \in \mathbb{R} \setminus \{0\}$  with  $c_\alpha \rightarrow c$ , where the  $u_\alpha$ 's and  $c_\alpha$ 's are nets indexed by  $\alpha$ 's from the same index set, we have  $c_\alpha \delta(u_\alpha) \rightarrow c \delta(u)$ . Moreover, for arbitrary  $u_\alpha \in \mathbb{S}^{n-1}$  and  $c_\alpha \rightarrow 0$ , we have  $c_\alpha \delta(u_\alpha) \rightarrow 0$ . Thus, the convergence is defined for finite signed Borel measures whose supports have at most one point. Then the convergence is defined for finite sums of such sequences as well (and in fact, only for these, see the formal definition in the next paragraph).

More exactly, a sequence (or more generally, a net)  $\mu_\alpha = \sum_{i=1}^{m_\alpha} \mu_\alpha(\{u_i\}) \delta(u_i)$  of finite signed Borel measures on  $\mathbb{S}^{d-1}$  with  $m_\alpha \leq m$  can converge only to a finite signed Borel measure of support of at most  $m$  points. Moreover,  $\mu_\alpha$  tends to a finite signed Borel measure  $\mu = \sum_{i=1}^{m'} \mu(\{u_i\}) \delta(u_i)$  on  $\mathbb{S}^{n-1}$  with  $1 \leq m' \leq m$  if and only if the following holds. For each  $\alpha$ , there exists a partition of  $\{1, \dots, m_\alpha\}$  of cardinality  $m'$ , say  $\{P_{\alpha 1}, \dots, P_{\alpha m'}\}$  (where each  $P_{\alpha j}$  is non-empty), such that

- (A) for any  $j \in \{1, \dots, m'\}$ , the sets  $P_{\alpha j}$  converge to  $u_j$  (i.e., for any neighbourhood  $U_j$  of  $u_j$  and for all sufficiently large  $\alpha$ , we have  $P_{\alpha j} \subset U_j$ ) and
- (B) for any  $j \in \{1, \dots, m'\}$ , the sum  $\sum \{\mu_\alpha(\{u_i\}) \mid i \in P_{\alpha j}\}$  converges to  $\mu(\{u_j\})$ .

The same sequence (or more generally a net)  $\mu_\alpha$  tends to the 0 (finite signed Borel) measure if and only if

- (C)  $\sum_{i=1}^{m_\alpha} |\mu_\alpha(\{u_i\})| \rightarrow 0$ . (This corresponds to the case  $m' = 0$ , and also here, an empty sum means 0.)

For convex polytopes, Theorem F can be rewritten for  $\mu_K = \sum_{i=1}^m \mu_K(\{u_i\}) \delta(u_i)$  as follows.

**Theorem F'** (Minkowski, see also [55, p. 389, (7.1.1), pp. 389–390, p. 390, Theorem 7.1.1, p. 397, Theorem 7.2.1]). *Let  $m > n \geq 2$  be integers, let  $S_1, \dots, S_m > 0$ , and let  $u_1, \dots, u_m \in \mathbb{S}^{n-1}$ . Then there exists a non-degenerate convex polytope  $P$  having  $m$  facets with facet areas  $S_1, \dots, S_m$  and respective facet outer unit normals  $u_1, \dots, u_m$  if and only if*

- (i)  $\sum_{i=1}^m S_i u_i = 0$ , and
- (ii)  $u_1, \dots, u_m$  do not lie in a linear  $(n-1)$ -subspace of  $\mathbb{R}^n$ .

*Moreover, if  $P$  exists, it is unique up to translations.*

For convex polytopes with at most  $m$  facets, Theorem G can be rewritten for  $\mu_K = \sum_{i=1}^m \mu_K(\{u_i\}) \delta(u_i)$  as follows.

**Theorem G'** (Minkowski, see also [55, p. 198, Theorem 4.1.1, p. 205, pp. 392–393, proof of Theorem 7.1.2] and the addition after our Theorem G). *Let  $m > n \geq 2$  be integers. Then the mapping  $\{\text{translates of } K\} \mapsto \mu_K$  defined in Definition 1 and after Theorem F is a homeomorphism between its domain and its range. Its domain is the subspace corresponding to the non-degenerate convex polytopes with at most  $m$  facets of the quotient topology of the topology on the convex bodies (induced by the Hausdorff metric) with respect to the equivalence relation of being translates. Its range is the set of finite Borel measures on  $\mathbb{S}^{n-1}$  with supports of at most  $m$  points satisfying (i) and (ii) of Theorem F', with the subspace topology of the weak\* topology on  $C(\mathbb{S}^{d-1})^*$ . This subspace topology is described in more explicit form before Theorem F'.*

## 4. PROOFS FOR THE EUCLIDEAN CASE

Essentially the following proposition was used in [31] without explicitly stating and proving it. It can be considered as folklore (as part of the proof of the folklore Theorem 1), but we state and prove it for completeness.

**Proposition 6.** *Let  $m > n \geq 3$  be integers. Let  $\delta > 0$  and let  $S_m \geq S_{m-1} \geq \dots \geq S_1 > 0$  be numbers such that  $S_m < S_1 + S_2 + \dots + S_{m-1}$ . Then there are pairwise distinct unit vectors  $v_1, v_2, \dots, v_m \in \mathbb{S}^{n-1}$  with the following properties:*

- (i) *they lie in the open  $\delta$ -neighbourhood of the  $x_1x_2$ -coordinate plane,*
- (ii) *they do not lie in a linear  $(n-1)$ -subspace of  $\mathbb{R}^n$ , and*
- (iii)  $S_1v_1 + S_2v_2 + \dots + S_mv_m = 0$ .

*Proof.* Let  $P$  be the  $x_1x_2$ -coordinate plane in  $\mathbb{R}^n$ . Since  $S_m < S_1 + S_2 + \dots + S_{m-1}$ , there exists a convex polygon  $A_1A_2\dots A_m$  (with angles strictly smaller than  $\pi$ ) in  $P$  such that  $|A_iA_{i+1}| = S_i$  for  $i = 1, \dots, m$  (indices considered modulo  $m$ ), see [32, p. 44], [34, pp. 53–54]. Then the edge directions  $u_i := \overrightarrow{A_iA_{i+1}}/|A_iA_{i+1}| \in \mathbb{S}^{n-1} \cap P$  are distinct unit vectors. We will perturb  $A_1A_2\dots A_m$  to a spatial polygon  $B_1B_2\dots B_m$ , keeping the side lengths equal:  $|B_iB_{i+1}| = |A_iA_{i+1}| = S_i$ . The unit vectors  $v_i := \overrightarrow{B_iB_{i+1}}/S_i$  will then fulfill (iii) by construction.

Clearly, for  $\|v_i - u_i\| < \delta$ , the vector  $v_i$  lies in the open  $\delta$ -neighbourhood of  $P$ , hence (i) is satisfied. Further, for  $\delta$  sufficiently small, the vectors  $v_1, \dots, v_m$  are also pairwise distinct.

Let  $k$  denote the largest integer such that there are arbitrarily small perturbations  $B_i$  of our original points  $A_i$  that satisfy the following: all edges have the right length  $|B_iB_{i+1}| = S_i$ , and the dimension of the affine hull of  $B_1, \dots, B_m$  has dimension  $k$ .

Assume for contradiction that  $k < n$ . Then, by  $m \geq n + 1 \geq k + 2$ , there is an affine dependence among the  $B_i$ 's. Let, for example,  $B_m$  lie in the affine hull  $H$  of  $B_1, \dots, B_{m-1}$ . Then, fixing  $|B_{m-1}B_m|$  and  $|B_mB_1|$ , the point  $B_m$  can move on an  $(n-2)$ -sphere around the axis  $B_{m-1}B_1$  in a hyperplane perpendicular to  $H$ . Hence there is an arbitrarily small perturbation of  $B_m$  lying outside  $H$ , while  $\text{aff}\{B_1, \dots, B_{m-1}\}$  already spans  $H$ . Thus we have obtained a contradiction to the choice of  $k$ .

This proves  $k = n$  and thus (ii). □

**4.1. Proof of Theorem 1.** The implication (ii)  $\implies$  (i) is evident.

The implication (i)  $\implies$  (iii) is well-known, but we give the proof for completeness. Using the notations from Theorem F', we have  $S_m = \|S_mu_m\| = \|\sum_{i=1}^{m-1} S_iu_i\| \leq \sum_{i=1}^{m-1} S_i$ . The only case of equality is the degenerate case given in condition (iii') of the theorem.

Finally, (iii)  $\implies$  (ii) follows from Proposition 6 and Minkowski's Theorem F'.

The degenerate case, with (iii'), follows from the above considerations. □

**4.2. Proofs for Theorem 2.** We need the following relation between the surface area, diameter and volume of a convex body. Here  $\kappa_{n-1}$  is the volume of the unit ball in  $\mathbb{R}^{n-1}$ .

**Proposition 7** (Gritzmann, Wills and Wrase [27]). *Let  $K \subset \mathbb{R}^n$  be a convex body. Then the inequality  $S(K)^{n-1} > \kappa_{n-1} \cdot \text{diam}(K) \cdot (nV(K))^{n-2}$  holds and this inequality is sharp.*  $\square$

We will construct the polytope for Theorem 2 by Minkowski's Theorem F'. We need to choose only an appropriate surface area measure. For a convex polytope, this finite Borel measure is concentrated in finitely many points. Assume that for given facet areas we are far from the degenerate case where this measure is concentrated in a great- $\mathbb{S}^{n-2}$ . Then by compactness, the volume of the convex polytope is bounded from below. Therefore, to get an arbitrarily small volume, we must approach the degenerate case. This will be done in the following proof. Recall also the paragraph after Theorem E citing [31] (p. 3), where also the degenerate case was approximated — however, for simplices only.

**4.3. First proof of Theorem 2.** By Proposition 6, for any  $\delta > 0$ , there are pairwise distinct vectors  $v_1, v_2, \dots, v_m \in \mathbb{S}^{n-1}$  in the open  $\delta$ -neighbourhood of the  $x_1x_2$ -coordinate plane, satisfying the following. They do not lie in a linear  $(n-1)$ -subspace of  $\mathbb{R}^n$ , and  $S_1v_1 + S_2v_2 + \dots + S_mv_m = 0$ . By Minkowski's Theorem F' there exists a non-degenerate convex polytope  $P = P(\delta)$  in  $\mathbb{R}^n$  having  $m$  facets with areas  $S_1, \dots, S_m$  and unit outer normals  $v_1, \dots, v_m$ .

Let us consider the sequence of polytopes  $P_k = P(1/k)$  for  $k = 1, 2, \dots$ . We will show that  $V(P_k) \rightarrow 0$  as  $k \rightarrow \infty$ . Assume the contrary. Then (possibly passing to a subsequence), we may assume without loss of generality that  $V(P_k) \geq \alpha > 0$ .

By Proposition 7, we get the inequality

$$S(P_k)^{n-1} > \kappa_{n-1} \cdot \text{diam}(P_k) \cdot (nV(P_k))^{n-2} \geq \kappa_{n-1} \cdot \text{diam}(P_k) \cdot (n\alpha)^{n-2},$$

where  $S(P_k) = \sum_{i=1}^m S_i$  is a constant. From this inequality, we conclude that  $\text{diam}(P_k)$  is bounded by some constant  $D$  for all  $k$ .

By applying translations, we may assume without loss of generality that all  $P_k$ 's have a common point. Therefore, all polytopes  $P_k$  lie in a ball of radius  $D$ . Using compactness (possibly passing to a subsequence), we may assume even more. The sequence  $P_k$  tends (in the Hausdorff metric) to a non-empty compact convex set  $P_0$  as  $k \rightarrow \infty$  [55, p. 50, Theorem 1.8.6]. Therefore,  $V(P_0) \geq \alpha > 0$ , and hence  $P_0$  is a convex body. Moreover, by Minkowski's Theorem G' (actually only by the continuity of the bijection in that theorem),  $P_0$  is a convex polytope having  $m$  facets with all facet outer unit normals in the  $x_1x_2$ -coordinate plane. However, this is a contradiction to condition (ii) of Theorem F'.  $\square$

*Remark 5.* Instead of Proposition 7, where the multiplicative constant is sharp, we could have used a consequence of the Aleksandrov-Fenchel inequality [55, p. 327, Theorem 6.3.1] to show that the diameter is bounded. Namely: the quermassintegrals  $W_i(K)$  ([55, p. 209]) for  $0 \leq i \leq n$  form a logarithmically concave sequence. Here,  $W_0(K) = V(K)$  and for fixed  $n$ ,  $W_1(K)$  is proportional to  $S(K)$  and  $W_{n-1}(K)$  is proportional to the mean width of  $K$  ([55, p. 210, p. 291, (5.3.12)]). Then apply this logarithmic convexity for volume, constant times surface area and constant times mean width. Finally, use the fact that the quotient of the diameter and the mean width is between two positive numbers (depending only on  $n$ ). This yields the inequality of Proposition 7 with a weaker constant.

*Example 1.* We give an example of a family of tetrahedra ( $n = 3$  and  $m = 4$ ) with constant facet areas and arbitrarily small volume. The tetrahedra look like thin vertical needles and have vertices  $(\pm\varepsilon, 0, -(1/\varepsilon)\sqrt{1 - \varepsilon^4/4})$  and  $(0, \pm\varepsilon, (1/\varepsilon)\sqrt{1 - \varepsilon^4/4})$ . All facets have area 2, and the volume is  $(4\varepsilon/3)\sqrt{1 - \varepsilon^4/4}$ , which tends to zero as  $\varepsilon \rightarrow 0$ .

**4.4. Second proof of Theorem 2. 1.** First we will construct a partition  $\mathcal{P} = \{P_1, \dots, P_{n+1}\}$  of the index set  $\{1, \dots, m\}$  into  $n + 1$  classes. We will achieve that the  $n + 1$  numbers  $\sum_{i \in P_j} S_i$  (for  $1 \leq j \leq n + 1$ ) have the property that

(\*) the largest of these numbers is smaller than the sum of all others.

We start with the partition into  $m$  singleton classes. Suppose that we already have constructed a partition  $\mathcal{Q} = \{Q_1, \dots, Q_k\}$  such that

(\*\*) the largest of the numbers  $T_j := \sum_{i \in Q_j} S_i$ , where  $1 \leq j \leq k$ , is smaller than the sum of all the other numbers  $T_j$ .

If  $k = n + 1$ , then we stop. If  $k > n + 1$ , then let us assume  $T_1 \leq T_2 \leq \dots \leq T_k$ . Now we take the two classes  $Q_1$  and  $Q_2$  with the two smallest sums and form their union while the other classes  $Q_j$  are kept. In the new partition, the partition class that has maximal sum  $T_j$  can be either the same partition class as in the preceding step or the newly constructed union. In the first case, (\*\*) is evident. In the second case, we have  $T_1 + T_2 \leq T_{k-1} + T_k < T_3 + \dots + T_{k-1} + T_k$  before taking the union since  $k \geq n + 2 \geq 5$ . Thus  $T_{\text{new}} := T_1 + T_2 < T_3 + \dots + T_{k-1} + T_k$ . Therefore, (\*\*) holds in this case as well.

This proves that (\*) holds for the final partition.

**2.** We consider the partition  $\mathcal{P} = \{P_1, \dots, P_{n+1}\}$  constructed above. For the sums  $R_j := \sum_{i \in P_j} S_i$ , we use Theorem E to construct a non-degenerate simplex  $S$  that has these facet areas and has an arbitrarily small volume. Then, for the respective outer unit normals  $u_j$  of the facets of this simplex, we have  $\sum_{j=1}^{n+1} R_j u_j = 0$ .

**3.** We will now split each facet of the simplex into almost parallel facets to get the desired polytope with  $m$  facets. Let  $\varepsilon > 0$  be small. For each  $1 \leq j \leq n + 1$ , choose a linear 2-subspace  $X_j$  containing  $u_j$ . Choose a vector  $u_{ji} \in X_j$  for each  $i \in P_j$  such that the vectors  $-(1 - \varepsilon)(\sum_{i \in P_j} S_i)u_j$  and  $S_i u_{ji}$  (for  $i \in P_j$ ) are the side vectors of a

convex polygon in  $X_j$ . Since the length of the first vector is almost equal to the sum of the others, all  $u_{ji}$  are close to  $u_j$  for  $\varepsilon$  sufficiently small. Then all vectors  $S_i u_{ji}$  for  $1 \leq j \leq n + 1$  and  $i \in P_j$  linearly span  $\mathbb{R}^n$ , and their sum is

$$\sum_{1 \leq j \leq n+1} \left( \sum_{i \in P_j} S_i u_{ji} \right) = \sum_{1 \leq j \leq n+1} (1 - \varepsilon) \left( \sum_{i \in P_j} S_i \right) u_j = (1 - \varepsilon) \sum_{1 \leq j \leq n+1} R_j u_j = 0.$$

**4.** By Minkowski's Theorem F', there exists a non-degenerate convex polytope with facet outer unit normals  $u_{ji}$  and facet areas  $S_i$  (for all  $1 \leq j \leq n + 1$  and all  $i \in P_j$ ).

Observe that we have changed in the course of the proof the surface area measure only a little bit (in the weak\*-topology of  $C(\mathbb{S}^{n-1})^*$ ). Therefore, after a suitable translation, the obtained convex polytope is arbitrarily close to the original simplex  $S$  by Theorem G' (actually only by the continuity of the inverse of the bijection in that theorem). Since the simplex had an arbitrarily small volume, our convex polytope also has an arbitrarily small volume.  $\square$

**4.5. Third proof of Theorem 2.** The first two proofs of Theorem 2 did not give geometric information about the constructed polytopes. (The first proof used an argument by contradiction and the second proof used the examples of the simplices.) Now we give a third proof that is more quantitative and will give also geometric information. This proof constructs “needle-like” polytopes, as in Example 1. See also the paragraph following the statement of Theorem 2 (p. 5).

First we give the proof for  $n = 3$  dimensions. We begin with an elementary lemma. It shows that a convex polytope in  $\mathbb{R}^3$  that has steep (almost vertical) facets must have steep edges, as long as the angles between the normal vectors of different facets are bounded away from 0 and  $\pi$ .

**Lemma 1.** *Consider two planes in  $\mathbb{R}^3$  with unit normals  $u^+$  and  $u^-$ . Assume that  $u^+$  and  $u^-$  enclose an angle at most  $\varepsilon \in (0, \pi/2)$  with the  $xy$ -plane, and the angle between them lies in  $[\beta, \pi - \beta]$ , where  $0 < \beta \leq \pi/2$ . Then their intersection line encloses an angle at most*

$$\delta := \arcsin \frac{\sin \varepsilon}{\sin(\beta/2)}$$

with the  $z$ -axis, provided that  $\varepsilon \leq \beta/2$ . This inequality is sharp.

*Proof.* We choose a new coordinate system in the following way. The intersection line becomes the vertical axis, and the two normal vectors  $u^+, u^- \in \mathbb{S}^2$  lie in the horizontal plane, enclosing an angle  $\beta' \in [\beta, \pi - \beta]$  with each other. In the new coordinate system, the original North Pole becomes  $n = (n_1, n_2, n_3) \in \mathbb{S}^2$ .

By hypothesis,

$$\langle n, u^- \rangle, \langle n, u^+ \rangle \in [-\sin \varepsilon, \sin \varepsilon]. \quad (5)$$

We want to conclude that

$$|\langle (0, 0, 1), n \rangle| = |n_3| \geq \cos \delta, \quad (6)$$

i.e., that

$$\sqrt{n_1^2 + n_2^2} \leq \sin \delta. \quad (7)$$

The points  $(n_1, n_2) \in \mathbb{R}^2$  (projections of  $n$  to the  $xy$ -plane) for  $n$  satisfying (5) form a rhomb of height  $2 \sin \varepsilon$  and angles  $\beta', \pi - \beta'$ . A farthest point of this rhomb from  $(0, 0)$  is one of the vertices and its distance from  $(0, 0)$  is  $\max\{(\sin \varepsilon)/\sin(\beta'/2), (\sin \varepsilon)/\cos(\beta'/2)\} \leq (\sin \varepsilon)/\sin(\beta/2) = \sin \delta$ . That is, (7), or equivalently, (6) holds and both are sharp inequalities. Hence, the inequality of the lemma holds and it is sharp.  $\square$

**Lemma 2.** *Consider a convex polyhedron  $P \subset \mathbb{R}^3$  with facet areas  $S_1, \dots, S_m$ . Assume that its facet outer normals enclose an angle at most  $\varepsilon$  with the  $xy$ -plane and the angle between any two of them lies in  $[\beta, \pi - \beta]$ , where  $0 < \beta \leq \pi/2$ . Then its volume is bounded by*

$$V(P) \leq 2^{-1/4} \pi^{-1} \cdot \left( \sum_{i=1}^m S_i^{3/4} \right)^2 \cdot \left( \frac{\sin \varepsilon}{\sin(\beta/2)} \right)^{1/2},$$

if  $(\sin \varepsilon)/\sin(\beta/2) \leq 1/\sqrt{2}$ .

*Proof.* We denote by  $s_i(z)$  the length of the horizontal cross-section of the  $i$ -th facet at height  $z$ , and by  $s_i^{\max}$  the maximum length of such a horizontal cross-section. Let  $h_i$  be the “height” of the  $i$ -th face: the difference between the maximum and the minimum  $z$ -coordinates of its points. Let  $h'_i$  be the “tilted height” of this facet in its own plane, i.e., the height when the plane is rotated into vertical position about one of its horizontal cross-sections.

Since  $(\sin \varepsilon)/\sin(\beta/2) < 1$ , we have by Lemma 1 that  $P$  has no horizontal edges. Therefore, using the quantity  $\delta$  introduced in Lemma 1, we get

$$s_i^{\max} \leq h_i \cdot \tan \delta. \quad (8)$$

Namely, from the minimal  $z$ -coordinate — where  $s_i(z) = 0$  —  $s_i(z)$  can increase only with a speed at most  $2 \tan \delta$  ( $< \infty$ ) to reach its maximal value  $s_i^{\max}$ . This is clear for a vertical face, and for a nonvertical face the speed is even smaller. Observe that the  $i$ -th facet lies in an upwards circular cone with vertex the lowest point of the  $i$ -th facet and directrices enclosing an angle  $\delta$  with the  $z$ -axis. From the maximal value it must decrease again with speed at most  $2 \tan \delta$  till 0 at the maximal  $z$ -coordinate.

Therefore, using for (10) inequality (8),

$$S_i \geq s_i^{\max} h'_i / 2 \geq s_i^{\max} h_i / 2 \quad (9)$$

$$\geq (s_i^{\max})^2 / (2 \tan \delta). \quad (10)$$

This gives

$$s_i^{\max} \leq \sqrt{2S_i \tan \delta}. \quad (11)$$

These relations allow us to bound the volume  $V(P)$  as follows, by using the isoperimetric inequality on each horizontal slice.

$$\begin{aligned} V(P) &= \int_{-\infty}^{\infty} (\text{area of cross-section of } P \text{ at height } z) dz \\ &\leq \int_{-\infty}^{\infty} \left( \sum_{i=1}^m s_i(z) \right)^2 dz / (4\pi) \\ &= \sum_{i=1}^m \sum_{j=1}^m \int_{-\infty}^{\infty} s_i(z) s_j(z) dz / (4\pi) \\ &\leq \sum_{i=1}^m \sum_{j=1}^m s_i^{\max} s_j^{\max} \min\{h_i, h_j\} / (4\pi) \quad (12) \end{aligned}$$

$$\begin{aligned} &\leq \sum_{i=1}^m \sum_{j=1}^m s_i^{\max} s_j^{\max} \sqrt{h_i h_j} / (4\pi) \\ &= \left( \sum_{i=1}^m s_i^{\max} \sqrt{h_i} \right)^2 / (4\pi) \\ &= \left( \sum_{i=1}^m \sqrt{s_i^{\max}} \sqrt{s_i^{\max} h_i} \right)^2 / (4\pi) \\ &\leq \left( \sum_{i=1}^m (2S_i \tan \delta)^{1/4} \sqrt{2S_i} \right)^2 / (4\pi) \quad (13) \end{aligned}$$



$$\begin{aligned}
&= \frac{\left(\sum_{i=1}^m S_i^{3/4}\right)^2}{\sqrt{2}\pi} \cdot \sqrt{\frac{(\sin \varepsilon)/\sin(\beta/2)}{\sqrt{1 - (\sin^2 \varepsilon)/\sin^2(\beta/2)}}} \\
&\leq \frac{\left(\sum_{i=1}^m S_i^{3/4}\right)^2}{\sqrt{2}\pi} \cdot \sqrt{\frac{\sqrt{2} \sin \varepsilon}{\sin(\beta/2)}}.
\end{aligned} \tag{14}$$

The first inequality uses the isoperimetric inequality. The second inequality (12) bounds the integral by an upper bound of the non-negative integrand times the length of the interval where the integrand is positive. For (13), we have used (9) and (11). To obtain (14), we have used Lemma 1. The last inequality simplifies the denominator under the assumption  $(\sin \varepsilon)/\sin(\beta/2) \leq 1/\sqrt{2}$  of the lemma.  $\square$

**4.6. Third proof of Theorem 2 for  $n = 3$  dimensions.** As in the first proof, we use Minkowski's Theorem F'. We want to apply Lemma 2, making  $\varepsilon$  small. Thus, we must let the normal vectors with given lengths  $S_i$  converge to the  $xy$ -plane, keeping their sum to be 0. Moreover, the linear span of the outer unit facet normals should be  $\mathbb{R}^3$ . Then we apply Minkowski's Theorem F'. In the limiting configuration the normals will lie in the  $xy$ -plane. They must form angles in  $[\beta, \pi - \beta]$  (with  $\beta \in (0, \pi/2]$ ) with each other in order that Lemma 2 should work. Thus we must avoid parallel sides.

Will show that there is only one exceptional case in which parallel sides cannot be avoided. Consider a planar convex  $m$ -gon  $M$  with sides  $S_i$  that has the minimum number of parallel pairs of sides. Let us assume that  $M$  has a side such that the sum of the two incident angles is different from  $\pi$ . Then by a small length-preserving motion of this side and the neighbouring two sides, one can achieve the following. This side changes its direction while new parallel pairs of sides are not created. Therefore,  $M$  can have a parallel pair of sides only if, for each of these sides, the sum of the incident angles is  $\pi$ . That is, we have four vertices that determine two parallel sides and whose outer angles (i.e.,  $\pi$  minus the inner angles) have sum  $2\pi$ . Since the sum of all outer angles is  $2\pi$ , there are no more vertices and  $M$  must be a parallelogram. If its sides are not equal then we rearrange the side vectors so as to obtain a (convex) deltoid that is not a parallelogram. So the only remaining case is when  $m = 4$  and  $S_1 = S_2 = S_3 = S_4$ . However, this case has been treated in Example 1: a tetrahedron with four faces of equal areas and having an arbitrarily small positive volume. Suitable inflations provide examples for all values of  $S_i$ .

Disregarding this exceptional case, we have now a strictly convex polygon in the  $xy$ -plane without parallel sides. Assume that the angle between any two edges is in the range  $[\beta_1, \pi - \beta_1]$  for some  $\beta_1 > 0$ . We still need to perturb the sides so that the edge vectors span  $\mathbb{R}^3$ . Consider the first three consecutive vertices  $A_1, A_2, A_3$  of  $M$ . Let us fix  $A_1$  and  $A_3$ . Rotate the two sides  $[A_1, A_2]$  and  $[A_2, A_3]$  about the line through  $A_1$  and  $A_3$  through a small angle  $\alpha > 0$  while keeping their lengths fixed. The  $m - 2 \geq 2$  remaining side vectors span the  $xy$ -plane since they are not parallel. At the same time, the vector  $\overrightarrow{A_1 A_2}$  points out of the  $xy$ -plane and therefore the edge vectors span  $\mathbb{R}^3$ .

By making the angle of rotation  $\alpha$  small enough, we can ensure the following. The angle between all edge vectors of the perturbed polygon  $M(\alpha)$  is still in the range  $[\beta_2, \pi - \beta_2]$  for some fixed  $\beta_2 > 0$ . Moreover, the angle  $\varepsilon$  of the side vectors

with the  $xy$ -plane can be made arbitrarily small. We use the edge vectors  $\vec{S}_i$  of  $M(\alpha)$  as outer normals and construct the polytope  $P$  by Minkowski's Theorem F', with  $S_i := \|\vec{S}_i\|$  and  $u_i := \vec{S}_i/S_i$ . By Lemma 2, the volume can be made arbitrarily small.  $\square$

*Remark 6.* In the polytope that we have constructed, all facets except two are vertical. By going through the proof of Lemma 2, one can see the following. It would have been sufficient to assume the constraint  $[\beta, \pi - \beta]$  on the angles for those pairs of facet normals that involve at least one of the two nonvertical facets.

*Example 2.* For odd dimension  $n = 2k + 1$ , there is a higher-dimensional generalization of Example 1. Consider a large regular  $k$ -simplex of edge length  $a := 1/\varepsilon$  in the  $x_{k+2} \dots x_n$ -coordinate plane. It has  $k + 1$  vertices  $v_1, \dots, v_{k+1}$ . At each vertex  $v_i$ , we draw a short segment of length  $b := \varepsilon$  centred at  $v_i$  in the direction of the  $x_i$ -axis. The convex hull of the union of these segments is an  $n$ -simplex with congruent facets. The facet areas are  $\sim \text{const} \cdot a^k b^k = \text{const}$ , while the volume is  $\text{const} \cdot a^k b^{k+1} = \text{const} \cdot \varepsilon$ , which becomes arbitrarily small as  $\varepsilon \rightarrow 0$ .

#### 4.7. Third proof of Theorem 2 for $n > 3$ dimensions.

4.7.1. *Construction of an almost flat spatial polygon.* As for  $n = 3$ , we start with a planar convex  $m$ -gon  $M$  in the  $x_1 x_2$ -coordinate plane, where  $m \geq n + 1$ . It has side vectors  $\vec{S}_i$  (this notation will be preserved also after perturbations) with side lengths  $S_i$ , for  $1 \leq i \leq m$ .

$M$  is contained in the  $x_1 \dots x_{n-1}$ -coordinate hyperplane  $X$ . By small perturbations of the closed polygon  $M$  in  $X$  that preserve the side lengths  $S_i$ , we want to achieve that

- (\*) the perturbed (skew) closed polygon  $M \subset X$  has no  $n - 1$  side vectors lying in an  $(n - 2)$ -dimensional linear subspace of  $X$ .

Initially,  $M$  lies in a 2-dimensional plane. We will fulfill (\*) by following the proof of Proposition 6. Our desired conclusion is slightly stronger than in Proposition 6: there we excluded only the case that *all* vectors lie in a lower-dimensional subspace.

Assume that some  $i \leq n - 1$  side vectors lie in a linear subspace of dimension less than  $i$ , where  $i$  is the smallest number with this property. We will eliminate these linear dependencies iteratively. We have already seen how we can avoid parallel edges ( $i = 2$ ). The only case where parallel sides could not be avoided was  $m = 4$  and  $S_1 = S_2 = S_3 = S_4$  (a rhomb) and this happens only for  $n \leq m - 1 = 3$ . Therefore, we can assume  $i \geq 3$ . Observe also that by small perturbations the different vertices remain distinct.

We may any time rearrange the cyclic order of side vectors of  $M$  as we want. So we assume that the first  $i$  side vectors  $\vec{S}_1, \dots, \vec{S}_i$  are linearly dependent, and any  $i - 1$  side vectors are linearly independent. Number the vertices  $A_j$  so that  $\vec{S}_j$  goes from vertex  $A_j$  to  $A_{j+1}$  (indices taken modulo  $m$ ). We want remove this linear dependence by perturbing the vertex  $A_i$ . For a technical reason, we have to first refine the order of the side vectors even further. Let  $\sum_{j=1}^i \lambda_j \vec{S}_j = 0$  be the (unique, up to a scalar factor) linear dependence. The  $\lambda_j$ 's cannot be all equal, since  $\sum_{j=1}^i \vec{S}_j = \vec{A_1 A_{i+1}} = 0$  would imply that  $A_1$  and  $A_{i+1}$  are equal points. The

polygon has  $m > n > i$  sides, and hence this is excluded. Therefore, by permuting the side vectors if necessary, we can assume that  $\lambda_{i-1} \neq \lambda_i$ . Since all  $(i-1)$ -subsets are linearly independent, we have  $\lambda_j \neq 0$  for all  $j$ , and thus we can assume without loss of generality that  $\lambda_i = -1$ . In other words,  $\vec{S}_i = \sum_{j=1}^{i-1} \lambda_j \vec{S}_j$ , with  $\lambda_{i-1} \neq -1$ .

Now, fixing  $A_{i-1}$ ,  $A_{i+1}$ ,  $\|\vec{S}_{i-1}\|$ , and  $\|\vec{S}_i\|$ , we can perturb  $A_i$  as follows. The point  $A_i$  moves on an  $(n-3)$ -dimensional sphere in a hyperplane within  $X$  with affine hull orthogonal to the segment  $[A_{i-1}, A_{i+1}]$ . There is a small motion that moves  $A_i$  out of the subspace  $H := \text{aff}\{A_1, A_2, \dots, A_{i-1}, A_{i+1}\}$  of  $X$ .

Now we show that the dimension of this subspace  $H$  is in fact  $i-1$ , which implies that the dimension of  $\text{aff}\{A_1, A_2, \dots, A_i, A_{i+1}\}$  increases by 1 and  $\vec{S}_1, \dots, \vec{S}_i$  become linearly independent. Clearly,  $\dim H$  cannot be greater than  $i-1 = \dim \text{aff}\{A_1, A_2, \dots, A_{i-1}, A_{i+1}\}$ . To see that  $\dim H = i-1$ , we note that the vectors  $\vec{A_1A_2} = \vec{S}_1, \vec{A_2A_3} = \vec{S}_2, \dots, \vec{A_{i-2}A_{i-1}} = \vec{S}_{i-2}, \vec{A_{i-1}A_{i+1}} = \vec{S}_{i-1} + \sum_{j=1}^{i-1} \lambda_j \vec{S}_j$  are linearly independent, since  $\lambda_{i-1} \neq -1$ , and hence, their linear span has dimension  $i-1$ .

Thus, we have established that, by perturbing  $A_i$ , the vectors  $\vec{S}_1, \dots, \vec{S}_i$  become linearly independent. If the perturbation is small enough, then every set of side vectors that was linearly independent before the motion remains linearly independent. Therefore, the number of linearly dependent  $i$ -tuples of side vectors of  $M$  decreases. A finite number of iterations eliminates all linearly dependent  $i$ -tuples, and  $i$  can be increased (till  $n-1$ ), until  $(*)$  is eventually established. This concludes the construction of the polygone  $M$ .

Condition  $(*)$  can be rephrased in the following way. The determinant of any  $n-1$  normed side vectors  $\vec{S}_i/S_i$  of  $M$  (i.e., the signed volume of the parallelepiped spanned by them) is nonzero. We denote by  $b > 0$  the smallest absolute value of these determinants. This bound will play the role of the sine of the angle bound  $\beta$  in Lemmas 1 and 2.

4.7.2. *Steep facets imply steep edges.* We generalize Lemma 1 to higher dimensions:

**Lemma 3.** *Let  $n \geq 3$ , and consider  $n-1$  hyperplanes in  $\mathbb{R}^n$  making an angle at most  $\varepsilon < \pi/2$  with the vertical axis (the  $x_n$ -axis). If their unit normal vectors  $v_1, \dots, v_{n-1}$  span an  $(n-1)$ -parallelotope of volume at least  $b (> 0)$  then they intersect in a line. The angle between this line and the vertical direction is bounded by*

$$\delta := \arcsin \frac{(n-1)^{3/2} \sin \varepsilon}{b},$$

provided that  $(n-1)^{3/2} \sin \varepsilon \leq b$ .

For fixed  $n$ , the order of magnitude of this bound  $\delta$  as a function of  $\varepsilon$  and  $b$  is optimal. More precisely, for any  $\varepsilon$  and  $b$ , where  $0 < \varepsilon < \pi/2$  and  $0 < b \leq 1$ , there are instances with  $\sin \delta = \min\{1, (\sin \varepsilon) / \sin((\arcsin b)/2)\}$ .

*Proof.* Since the unit normal vectors  $v_1, \dots, v_{n-1}$  are linearly independent, the intersection of the hyperplanes is a line  $\ell$ . Let us choose a new orthonormal coordinate system where  $\ell$  is the last coordinate axis. Then the last coordinate of the vectors  $v_i$  is zero, and we may write these vectors as  $v_i = \begin{pmatrix} v'_i \\ 0 \end{pmatrix}$  with  $v'_i \in \mathbb{S}^{n-2} \subset \mathbb{R}^{n-1}$ . By assumption, the  $(n-1) \times (n-1)$  matrix  $V = (v'_1, \dots, v'_{n-1})$  has determinant of absolute value  $|\det V| \geq b$ .

Let  $p = \begin{pmatrix} p' \\ p_n \end{pmatrix}$ , with  $p' \in \mathbb{R}^{n-1}$ , be the unit vector of the original positive  $x_n$ -direction in the new coordinate system. Its angle  $\delta \in [0, \pi/2]$  with the line  $\ell$  satisfies  $\cos \delta = |p_n|$  and  $\sin \delta = \|p'\|$ , and thus our goal is to show that

$$\|p'\| \leq (n-1)^{3/2} \frac{\sin \varepsilon}{b}. \quad (15)$$

Let  $\alpha_i$  denote the angle between  $p$  and the normal  $v_i$ . By the angle assumption on the hyperplanes, we have  $\pi/2 - \varepsilon \leq \alpha_i \leq \pi/2 + \varepsilon$ . Therefore, with  $r_i := \cos \alpha_i = \langle p, v_i \rangle = \langle p', v'_i \rangle$ , we have  $|r_i| \leq \sin \varepsilon$ .

The  $n-1$  equations  $\langle p', v'_i \rangle = r_i$  form a linear system  $(p')^T V = (r_1, \dots, r_{n-1})$  (the column vectors of  $V$  being the  $v_i$ 's), i.e.,  $V^T p' = (r_1, \dots, r_{n-1})^T$ , which determines  $p'$  uniquely:

$$p' = (V^T)^{-1} (r_1, \dots, r_{n-1})^T. \quad (16)$$

We write  $\text{adj}(V^T)$  for the transpose of the matrix whose entries are the signed cofactors of the respective entries of  $V^T$ . By the formula  $(V^T)^{-1} = \text{adj}(V^T) / \det(V^T)$ , each entry of  $(V^T)^{-1}$  is an  $(n-2) \times (n-2)$  subdeterminant of  $V^T$  divided by  $\pm \det V^T$ . The rows of the submatrices of  $V^T$  are vectors of length at most 1 and therefore these subdeterminants are bounded in absolute value by 1. It follows that the entries of  $(V^T)^{-1}$  are bounded in absolute value by  $1/b$ . Since the  $|r_i|$ 's are at most  $\sin \varepsilon$ , we get from (16) that the  $n-1$  entries of  $p'$  are bounded by  $(n-1)(\sin \varepsilon)/b$  in absolute value. Hence we have proved (15).

To establish the lower bound, we can lift the tight three-dimensional example from Lemma 1 to  $n$  dimensions. For  $\varepsilon \leq \beta/2$ , the enclosed angle will be the same as in three dimensions, namely  $\arcsin((\sin \varepsilon) / \sin(\beta/2))$ , where  $\sin \beta = b$ . For  $\varepsilon > \beta/2$ , we use the example with  $\varepsilon = \beta/2$ . We embed the 3-dimensional example into  $\mathbb{R}^n$  by a linear isometry that maps the positive  $x, y, z$ -axes to the positive  $x_1, x_2, x_n$ -coordinate axes of  $\mathbb{R}^n$ . (The ‘‘vertical’’ direction is now the direction of the  $x_n$ -axis.) The two 2-planes of the three-dimensional example are turned into hyperplanes as follows. We replace them by their inverse images under the orthogonal projection of  $\mathbb{R}^n$  to the  $x_1 x_2 x_n$ -coordinate subspace. Simultaneously, we add the hyperplanes with equations  $x_3 = 0, \dots, x_{n-1} = 0$ .  $\square$

#### 4.7.3. Polytopes with steep facets have small volume.

**Lemma 4.** *Let  $n > 3$  be an integer. Assume that a convex polytope  $P \subset \mathbb{R}^n$  has facet areas  $S_1, \dots, S_m$ . Moreover, its outer unit facet normals enclose an angle at most  $\varepsilon \in (0, \pi/2)$  with the  $x_1 \dots x_{n-1}$ -plane. Also the volume of the  $(n-1)$ -parallelepiped spanned by any  $n-1$  unit facet normals of  $P$  is at least  $b > 0$ . Then its volume is bounded by*

$$V(P) \leq \text{const}_n \cdot \left( \sum_{i=1}^m S_i^{n/(2n-2)} \right)^2 \cdot \left( \frac{\sin \varepsilon}{b} \right)^{1/(n-1)},$$

if  $\sin^2 \varepsilon \leq b^2 / [2(n-1)^3]$ .

On the other hand, for  $n \geq 3$  and any  $m \geq 2n$ , there exists a suitable  $\varepsilon_0 \in (0, \pi/4)$ , such that the following holds. For any  $\varepsilon \in (0, \varepsilon_0)$ , there exists a convex polytope  $P(\varepsilon) \subset \mathbb{R}^n$ , with  $m$  facets, with the following properties. It satisfies the hypotheses of this lemma (except the one about the facet areas), with  $b$  depending

only on  $n$  and  $m$ , such that

$$\begin{aligned} V(P(\varepsilon)) &\geq \text{const}'_n \cdot S(P(\varepsilon))^{n/(n-1)} \cdot (\tan \varepsilon)^{1/(n-1)} \\ &\geq \text{const}'_n \cdot m^{-(n-2)/(n-1)} \cdot \left( \sum_{i=1}^m S_i(\varepsilon)^{n/(2n-2)} \right)^2 \cdot (\tan \varepsilon)^{1/(n-1)}. \end{aligned}$$

Here,  $S_1(\varepsilon), \dots, S_m(\varepsilon)$  are the areas of the facets of  $P(\varepsilon)$ . In particular, in the inequalities of Lemma 2 and this lemma, the order of magnitude as a function of  $\varepsilon$  is optimal.

*Proof.* We begin with the proof of the upper estimate. We denote by  $s_i(x_n)$  the  $(n-2)$ -volume of the horizontal cross-section of the  $i$ -th facet at height  $x_n$ . Moreover, we denote by  $s_i^{\max}$  the maximum  $(n-2)$ -volume of such a horizontal cross-section. Let  $h_i$  be the ‘‘height’’ of the  $i$ -th facet: the difference between the maximal and the minimal  $x_n$ -coordinates of its points. Let  $h'_i$  be the ‘‘tilted height’’ of this facet in its own hyperplane. That is, the height when the hyperplane is rotated into vertical position about one of its horizontal cross-sections.

Now, since  $\sin^2 \varepsilon \leq b^2/[2(n-1)^3]$ , the angle  $\delta$  from Lemma 3 lies in  $(0, \pi/2)$ . Hence, by Lemma 3,  $P$  has no horizontal edges, and thus, also no horizontal  $k$ -faces for any  $k \in \{1, \dots, n-2\}$ . Therefore, once more by Lemma 3, we know that every facet is contained in two rotationally symmetric cones with  $(n-1)$ -balls as bases. One cone has its apex at the unique lowest point of this facet and extends upwards from there. Its axis is vertical (parallel to the  $x_n$ -direction), and the directrices enclose an angle  $\delta$  with the  $x_n$ -axis. The other cone extends downwards from the highest point of the facet and has a vertical axis and directrices enclosing an angle  $\delta$  with the  $x_n$ -axis. We use the upwards cone from the minimal height till the arithmetic mean of the minimal and maximal heights. We use the downward cone for the other half of the vertical extent of the facet. By this argument, we can bound the maximum cross-section area  $s_i^{\max}$  of the  $i$ -th facet as follows.

$$s_i^{\max} \leq ((h_i/2) \cdot \tan \delta)^{n-2} \cdot \kappa_{n-2}. \quad (17)$$

(From the minimal height till the arithmetic mean of the minimal and maximal heights we have the following. Any horizontal cross-section of the cone is contained in some  $(n-1)$ -ball of radius at most  $R := (h_i/2) \cdot \tan \delta$ . Thus, any horizontal cross-section of the facet lies inside the intersection of its own affine hull with the upwards cone. That is, it lies in the intersection of an  $(n-2)$ -dimensional affine subspace with a cone whose base is an  $(n-1)$ -ball of radius at most  $R$ . Hence, this horizontal cross-section lies inside some  $(n-2)$ -ball of radius at most  $R$ . A similar argument holds for the downward cone.) Moreover, we also have

$$S_i \geq s_i^{\max} h'_i / (n-1) \geq s_i^{\max} h_i / (n-1). \quad (18)$$

Let us rewrite (17) and (18) as follows.

$$h_i^{-(n-2)} \cdot s_i^{\max} \leq ((\tan \delta)/2)^{n-2} \cdot \kappa_{n-2} \quad (19)$$

$$h_i \cdot s_i^{\max} \leq (n-1) S_i. \quad (20)$$

We multiply the  $1/[(2n-2)(n-2)]$ -th power of (19) with the  $n/(2n-2)$ -th power of (20) to get an inequality that we will need.

$$\begin{aligned}
& (s_i^{\max})^{(n-1)/(2n-4)} \sqrt{h_i} \\
& \leq ((\tan \delta)/2)^{1/(2n-2)} \cdot (\kappa_{n-2})^{1/[(2n-2)(n-2)]} \cdot ((n-1)S_i)^{n/(2n-2)}. \quad (21)
\end{aligned}$$

Let  $K := [(n-1)^{n-1} \kappa_{n-1}]^{-1/(n-2)}$  denote the constant of the isoperimetric inequality in  $n-1$  dimensions:

$$V_{n-1}(C) \leq K \cdot (V_{n-2}(\partial C))^{(n-1)/(n-2)} \quad (22)$$

(for  $C \subset \mathbb{R}^{n-1}$ ). Now we can bound the volume as follows.

$$\begin{aligned}
V(P) &= \int_{-\infty}^{\infty} [(n-1)\text{-volume of the cross-section of } P \text{ at height } x_n] dx_n \\
&\leq \int_{-\infty}^{\infty} \left[ \left( \sum_{i=1}^m s_i(x_n) \right)^{(n-1)/(2n-4)} \right]^2 dx_n \cdot K \\
&\leq \int_{-\infty}^{\infty} \left[ \sum_{i=1}^m s_i(x_n)^{(n-1)/(2n-4)} \right]^2 dx_n \cdot K \\
&= \int_{-\infty}^{\infty} \sum_{i=1}^m \sum_{j=1}^m s_i(x_n)^{(n-1)/(2n-4)} s_j(x_n)^{(n-1)/(2n-4)} dx_n \cdot K \\
&= \sum_{i=1}^m \sum_{j=1}^m \int_{-\infty}^{\infty} s_i(x_n)^{(n-1)/(2n-4)} s_j(x_n)^{(n-1)/(2n-4)} dx_n \cdot K \\
&\leq \sum_{i=1}^m \sum_{j=1}^m (s_i^{\max})^{(n-1)/(2n-4)} (s_j^{\max})^{(n-1)/(2n-4)} \min\{h_i, h_j\} \cdot K \quad (23)
\end{aligned}$$

$$\begin{aligned}
&\leq \sum_{i=1}^m \sum_{j=1}^m (s_i^{\max})^{(n-1)/(2n-4)} (s_j^{\max})^{(n-1)/(2n-4)} \sqrt{h_i h_j} \cdot K \\
&\leq (\tan \delta)^{1/(n-1)} \cdot 2^{-1/(n-1)} \cdot (\kappa_{n-2})^{1/[(n-1)(n-2)]} \cdot (n-1)^{n/(n-1)} \quad (24) \\
&\quad \times \left( \sum_{i=1}^m S_i^{n/(2n-2)} \right)^2 \cdot K
\end{aligned}$$

$$\begin{aligned}
&= \text{const}_n \cdot \left( \sum_{i=1}^m S_i^{n/(2n-2)} \right)^2 \left( \frac{(n-1)^{3/2}(\sin \varepsilon)/b}{\sqrt{1 - (n-1)^3(\sin^2 \varepsilon)/b^2}} \right)^{1/(n-1)} \quad (25) \\
&\leq \text{const}'_n \cdot \left( \sum_{i=1}^m S_i^{n/(2n-2)} \right)^2 \cdot \left( \frac{\sin \varepsilon}{b} \right)^{1/(n-1)}.
\end{aligned}$$

The first inequality uses the isoperimetric inequality (22). The second inequality uses the concavity of the function  $t^{(n-1)/(2n-4)}$  for  $t \in [0, \infty)$  and its vanishing at  $t = 0$ . (Observe that  $0 < (n-1)/(2n-4) \leq 1$ .) Inequality (23), as in (12), bounds the integral of a non-negative function by an upper bound of the integrand times the length of the interval where the integrand is positive. For (24), we have used the bound (21) that we derived above. Inequality (25) uses the bound  $\delta$  from Lemma 3. Finally, by hypothesis, the expression under the square root in the denominator of (25) is bounded below by  $1 - (n-1)^3(\sin^2 \varepsilon)/b^2 \geq 1/2$ . We have therefore established the claimed upper bound.

Now we give the example for the lower bound for  $n \geq 3$  and  $m \geq 2n$ . Let  $\varepsilon \in (0, \varepsilon_0)$ , where  $\varepsilon_0 \in (0, \pi/4)$  will be chosen later. Let us write  $\mathbb{R}^n = \mathbb{R}^{n-1} \oplus \mathbb{R}$ . Let  $T^+, T^- \subset \mathbb{R}^{n-1}$  be regular  $(n-1)$ -simplices circumscribed about the unit ball  $B^{n-1}$  of  $\mathbb{R}^{n-1}$ . Put them in such a general position w.r.t. each other so that any  $n-1$  of their altogether  $2n$  facet outer normals linearly span  $\mathbb{R}^{n-1}$ . Let  $n \leq m^+, m^-$  and  $m = m^+ + m^-$ . Let  $R^\pm$  be obtained from  $T^\pm$  by intersecting it still with  $m^\pm - n$  closed halfspaces in  $\mathbb{R}^{n-1}$ , all containing  $B^{n-1}$ , with their boundaries touching  $B^{n-1}$ . Then

$$B^{n-1} \subset R^\pm \subset T^\pm \subset (n-1)B^{n-1}.$$

Let the altogether  $m = m^+ + m^-$  facet outer unit normals of  $R^+$  and  $R^-$  satisfy the same condition of general position as above. Namely, any  $n-1$  of them linearly span  $\mathbb{R}^{n-1}$ . Let  $b > 0$  be the minimum of the  $(n-1)$ -volumes of the  $(n-1)$ -parallelotopes spanned by any  $n-1$  of these altogether  $m$  facet outer unit normals.

Observe that for  $n = 3$  and  $m \geq 2$ , the largest value of  $b$  is  $\sin(\pi/m)$  — if we do not begin the construction with two regular triangles but allow any  $m$  facet outer unit normals in  $\mathbb{S}^{n-2} = \mathbb{S}^1$ . For  $n > 3$ , the maximal value of  $b$  can be bounded from above as follows — again not beginning with two regular simplices, but allowing any  $m$  facet outer unit normals in  $\mathbb{S}^{n-2}$ . Let us choose altogether  $n-1$  outer unit normal vectors of  $R^+$  and  $R^-$ , say,  $u_1, \dots, u_{n-1} \in \mathbb{S}^{n-2}$ . We have

$$\begin{aligned} \frac{|\det(u_1, \dots, u_{n-1})|}{(n-1)!} &= V_{n-2}(\text{conv}\{u_1, \dots, u_{n-1}\}) \cdot \text{dist}(0, \text{aff}\{u_1, \dots, u_{n-1}\}) / (n-1) \\ &\leq V_{n-2}(\text{conv}\{u_1, \dots, u_{n-1}\}) / (n-1). \end{aligned}$$

Here,  $\text{dist}(\cdot, \cdot)$  denotes distance. Thus, it suffices to bound  $V_{n-2}(\text{conv}\{u_1, \dots, u_{n-1}\})$  from above. This is the spherical analogue — for  $\mathbb{S}^{n-2}$  — of the celebrated Heilbronn problem. This problem asks about the maximum of the minimal  $n$ -volume of  $n$ -simplices spanned by any  $m$  points in  $[0, 1]^n$ . This problem is poorly understood. For an extensive literature on this problem, see [13, Ch. 11.2]. Unfortunately, this spherical variant cannot be reduced to the case of  $[0, 1]^{n-2}$  by taking the projection of, say, the intersection of  $\mathbb{S}^{n-2}$  with each orthant to the tangent  $\mathbb{R}^{n-2}$  at its centre. Namely, the area of  $\text{conv}\{u_1, \dots, u_{n-1}\}$  can be large even if its projection has a small area. In one direction, we have an implication: large projection areas imply large areas — but large areas still do not imply large values of  $|\det(u_1, \dots, u_{n-1})|$ . However, this spherical variant is a special case of the  $(n-1)$ -dimensional Heilbronn problem for  $[0, 1]^{n-1}$ . Namely, we can just add to any set of  $(n-1)$ -dimensional vectors in  $\mathbb{S}^{n-2}$  the single vector  $0$  — but probably we lose an essential part of the information in this way.

Let  $P^\pm(\varepsilon)$  be the half-infinite pyramid with vertex  $(0, \dots, 0, \pm \tan \varepsilon)$  and base  $R^\pm$ . Then

$$C_i^\pm(\varepsilon) \subset P^\pm(\varepsilon) \subset C_o^\pm(\varepsilon),$$

where  $C_i^\pm(\varepsilon)$  and  $C_o^\pm(\varepsilon)$  is a half-infinite cone with vertex  $(0, \dots, 0, \pm \tan \varepsilon)$  and base  $B^{n-1}$  and  $(n-1)B^{n-1}$ , respectively. Therefore,

$$C_i(\varepsilon) := C_i^+(\varepsilon) \cap C_i^-(\varepsilon) \subset P(\varepsilon) := P^+(\varepsilon) \cap P^-(\varepsilon) \subset C_o(\varepsilon) := C_o^+(\varepsilon) \cap C_o^-(\varepsilon).$$

Here,  $C_i(\varepsilon)$  and  $C_o(\varepsilon)$  are double cones over  $B^{n-1}$  and  $(n-1)B^{n-1}$ , respectively, with vertices  $(0, \dots, 0, \pm \tan \varepsilon)$ . Moreover,  $P(\varepsilon)$  is a convex polytope with  $m$  facets, all facet outer unit normals enclosing an angle  $\varepsilon$  with the  $x_1 \dots x_{n-1}$ -hyperplane.

(Actually they enclose an angle  $\varepsilon$  with the respective facet outer unit normal of  $R^+$  or  $R^-$  in  $\mathbb{R}^{n-1}$ .) If  $\varepsilon_0$  and thus also  $\varepsilon$  is sufficiently small then still any  $n - 1$  facet outer unit normals of  $P(\varepsilon)$  span an  $(n - 1)$ -parallelotope of volume at least some  $b' \in (0, b)$ .

A routine calculation gives

$$\frac{V(P(\varepsilon))}{S(P(\varepsilon))^{n/(n-1)}} \geq \frac{V(C_i(\varepsilon))}{S(C_o(\varepsilon))^{n/(n-1)}} = \frac{(\tan \varepsilon)^{1/(n-1)}}{n(2\kappa_{n-1})^{1/(n-1)}[1 + (n-1)\tan^2 \varepsilon]^{n/(2n-2)}}.$$

Therefore,

$$\begin{aligned} V(P(\varepsilon)) &\geq \frac{S(P(\varepsilon))^{n/(n-1)} \cdot (\tan \varepsilon)^{1/(n-1)}}{n(2\kappa_{n-1})^{1/(n-1)}[1 + (n-1)\tan^2 \varepsilon_0]^{n/(2n-2)}} \\ &\geq \frac{m^{-(n-2)/(n-1)} \left(\sum_{i=1}^m S_i(\varepsilon)^{n/(2n-2)}\right)^2 \cdot (\tan \varepsilon)^{1/(n-1)}}{n(2\kappa_{n-1})^{1/(n-1)}[1 + (n-1)\tan^2 \varepsilon_0]^{n/(2n-2)}}. \end{aligned}$$

Here,  $S_1(\varepsilon), \dots, S_m(\varepsilon)$  are the areas of the facets of  $P(\varepsilon)$ . The second inequality is equivalent to Hölder's inequality for the numbers  $S_i(\varepsilon)$ , between their arithmetic mean and their power mean with exponent  $n/(2n-2) \in (0, 1)$ . Finally, observe that  $\tan \varepsilon_0 \in (0, 1)$ .  $\square$

**4.7.4. Conclusion of the proof.** Now we can finish the third proof of Theorem 2 for  $n > 3$ . We proceed as for  $n = 3$  but instead of Lemma 2 we use Lemma 4. In Section 4.7.1 (see its last paragraph), we have constructed a closed  $m$ -gon  $M$  in the  $(n-1)$ -dimensional subspace  $X$  with the following property. Any  $n-1$  normed side vectors  $\vec{S}_i/S_i$  span a parallelotope of volume at least  $b$ . We follow the third proof of Theorem 2 for  $n = 3$ . We take the first three consecutive vertices  $A_1, A_2, A_3$  of  $M$  and “rotate”  $A_2$  out of the subspace  $X$ , keeping  $A_1, A_3$  and the lengths  $|A_1A_2|$  and  $|A_2A_3|$  fixed. We have a whole  $(n-2)$ -dimensional sphere on which  $A_2$  can move, which intersects  $X$  orthogonally. By bounding the distance by which  $A_2$  moves by a suitable threshold we can ensure the following. Any  $n-1$  normed side vectors  $\vec{S}_i/S_i$  still span a parallelotope of volume at least  $b'$  with some weaker bound  $b' > 0$ . The angle between  $[A_1, A_2]$  or  $[A_2, A_3]$  and the “horizontal” hyperplane  $X$  can be made arbitrarily small. Thus, Lemma 4 guarantees that the volume tends to zero as well.  $\square$

## 5. PROOFS FOR THE HYPERBOLIC CASE

For general concepts in hyperbolic geometry, we refer to [5, 7, 18, 37, 40, 45]. In particular, a *Lambert quadrilateral in  $\mathbb{H}^2$*  is a quadrilateral that has three right angles.

**5.1. Proof of Proposition 1.** Let  $H$  be the hyperplane of the facet  $F_m$  of  $P$  of area  $S_m$ . Let  $p: \mathbb{H}^n \rightarrow H$  be the orthogonal projection of  $\mathbb{H}^n$  to  $H$ . The image by  $p$  of the union of the  $m-1$  facets different from  $F_m$  contains  $F_m$ .

Let  $dS$  be a surface element at a point  $x \in \mathbb{H}^n$ . Let its image by  $p$  be the surface element  $dS'$  at  $p(x)$ . Clearly, it suffices to show that  $dS' \leq dS$ . We may assume that  $dS$  is an (infinitesimal)  $(n-1)$ -ball of radius  $dr$  in the tangent space  $T_x(\mathbb{H}^n)$  of  $\mathbb{H}^n$  at  $x$ .



First we deal with the case when  $dS$  is orthogonal to the line  $\ell(x, p(x))$ . (For  $x \in H$ , we mean by  $\ell(x, p(x))$  the line containing  $x$  and orthogonal to  $H$ .) Then  $dS'$  is an infinitesimal  $(n-1)$ -ball in  $T_{p(x)}(\mathbb{H}^n)$  of some radius  $dr'$ . By the trigonometric formulas of Lambert quadrilaterals in  $\mathbb{H}^2$  (see [45, §29, (V)] or [17, Theorem 2.3.1]), we have  $1 \leq \cosh |xp(x)| = (\tanh(dr)) / \tanh(dr')$ . Hence,  $dr' \leq dr$  and therefore,  $dS' \leq dS$ .

Now we extend this analysis to the case when  $dS$  is not orthogonal to the line  $\ell(x, p(x))$ . Then the image by  $p$  of the infinitesimal  $(n-1)$ -ball  $dS$  in  $T_x(\mathbb{H}^n)$  is an infinitesimal  $(n-1)$ -ellipsoid in  $T_{p(x)}(\mathbb{H}^n)$ . It has  $n-2$  semiaxes equal to  $dr'$  and the  $(n-1)$ -st semiaxis smaller than  $dr'$ . Hence,  $dS' < dS$  in this case.

The case of equality is clear: the polytope must degenerate to the doubly counted facet  $F_m$ .  $\square$

**5.2. Proof of Proposition 2.** From maximality of  $S_m, S_{m-1}, \dots, S_3$ , it follows that all vertices lie at infinity. Namely, a vertex cannot be incident only to the facets of areas  $S_2, S_1$ . Hence, also  $S_2, S_1$  are maximal.  $\square$

**5.3. Proof of Proposition 3.** Let  $P$  be a convex polyhedron as in the proposition with respective facets  $F_1, \dots, F_m$ . Let us consider any vertex  $v$  of some facet  $F_i$ . In the facets incident to  $v$ , the angle of  $F_i$  at  $v$  is at most the sum of the angles of all other facets incident to  $v$ . To see this, we intersect  $P$  with an infinitesimally small sphere with centre at this vertex (in the conformal model). We obtain a convex spherical polygon whose side lengths are the (convex) angles of the facets incident to  $v$  at  $v$ , all these angles being in  $[0, \pi)$ .

Summing these inequalities over all vertices  $v$  of  $F_i$  we obtain the following. The sum  $t_1$  of the angles of  $F_i$  is at most the sum  $t_2$  of the angles of all other facets at the vertices of  $F_i$ . The sum  $t_2$  is bounded above by the sum  $t_3$  of *all angles of all facets different from  $F_i$* . The resulting inequality  $t_1 \leq t_3$  is equivalent to the inequality to be proved.

Clearly, if we have at least one finite vertex with incident edges not in a plane, then we have at least one strict inequality among the summed inequalities. So in this case, we have strict inequality in the proposition.  $\square$

The inequality  $t_1 \leq t_3$  from this proof is discussed for the spaces  $\mathbb{R}^3$  and  $\mathbb{S}^3$  in Remark 9 in Section 6.1.

#### 5.4. Proof of Theorem 3.

**Proposition 8** ([5, p. 127], [28, Theorem 1, Proposition 2]). *For  $n \geq 2$ , a simplex in  $\mathbb{H}^n$  (with vertices at infinity admitted) is of maximal volume if and only if all its vertices are at infinity and it is regular. It has a finite volume.*

Let  $v_n$  be the maximal volume of a simplex in  $\mathbb{H}^n$ . For instance,  $v_2 = \pi$  and  $v_3 = -3 \int_0^{\pi/3} \log |2 \sin u| du = 1.0149416 \dots$  ([40, p. 20], [5, p. 127]). Obviously, the facet areas  $S_i$  of a compact simplex in  $\mathbb{H}^n$  are smaller than  $v_{n-1}$ .

**Lemma 5.** *The area  $S$  of a right triangle  $\triangle ABC \subset \mathbb{H}^2$  with angle  $\angle ACB = \pi/2$  and side lengths  $|AC| = b$  and  $|BC| = a$  fulfills the equation*

$$\tan S = \frac{\sinh a \cdot \sinh b}{\cosh a + \cosh b}.$$

*Proof.* This is a routine consequence of the trigonometric formulas for a right triangle in  $\mathbb{H}^2$ . We use  $S = \pi/2 - \alpha - \beta$ ,  $\tan \angle CBA = (\tanh b)/\sinh a$ ,  $\tan \angle CAB = (\tanh a)/\sinh b$  [18, p. 238], and  $\tanh x = (\sinh x)/\cosh x$ .  $\square$

**Lemma 6.** *Let  $d > 0$ . Assume that  $\Delta ABC \subset \mathbb{H}^2$  is a triangle such that  $|AB| \leq d$  and  $|AC| \leq d$ . Then the area  $S$  of this triangle is bounded by the inequality*

$$S \leq 2 \arctan \frac{\cosh d - 1}{2\sqrt{\cosh d}}.$$

*Proof.* Without loss of generality we may assume that  $|AB| = |AC| = d$ . Let  $H$  be the orthogonal projection of  $A$  to the line  $\ell(B, C)$ . The segment  $AH$  cuts the triangle  $ABC$  into two congruent right triangles. With  $x = \cosh |AH|$  and  $y = \cosh |BH| = \cosh |CH|$ , we have  $x, y \geq 1$  and  $xy = \cosh d$ . Let  $S$  be the area of  $\Delta ABC$ . Lemma 5 gives

$$\tan^2(S/2) = \frac{(x^2 - 1)(y^2 - 1)}{(x + y)^2}.$$

Looking for the maximum of the numerator and the minimum of the denominator subject to the constraints  $x, y > 0$  and  $xy = \cosh d$ , we see that the maximal value of  $S$  is attained for  $x = y = \sqrt{\cosh d}$ . This proves the lemma.  $\square$

To show that a tetrahedron with given facet areas exists, we will use a topological argument, which is encapsulated in the following lemma. The lemma guarantees the existence of a zero of a function under certain conditions on the boundary.

**Lemma 7.** *Let  $F = (f_1, f_2): P \rightarrow \mathbb{R}^2$  be a continuous function defined on a rectangular domain  $P = [0, a] \times [0, b]$ , where  $a, b > 0$ . Assume that there are  $u_1, u_2, v_1, v_2 \in \mathbb{R}$  such that  $u_1 > u_2$  and  $v_1 < v_2$  and*

$$\begin{aligned} f_1(x, 0) + f_2(x, 0) &\leq 0, & f_1(x, b) + f_2(x, b) &\geq 0, & \text{and} \\ u_1 f_1(0, y) + u_2 f_2(0, y) &\leq 0, & v_1 f_1(a, y) + v_2 f_2(a, y) &\leq 0 \end{aligned}$$

*for every  $0 \leq x \leq a$  and every  $0 \leq y \leq b$ . Then there exists a point  $(c, d) \in P$  such that  $F(c, d) = (0, 0)$ .*

*Proof.* The conclusion clearly holds if  $F$  vanishes at some point of the boundary  $\partial P$  of  $P$ . If  $F$  has no zero on  $\partial P$ , then it is sufficient to establish that the index of the vector field  $F$  on the curve  $\partial P$  is 1. This implies that  $F$  has a zero in the interior of  $P$  [29, p. 98, proof of Theorem VI.12, sufficiency].

To determine the index of  $F$ , we define the auxiliary function  $F_0: \partial P \rightarrow \mathbb{S}^1$  as follows. On the vertical boundaries of  $P$ , we let  $F_0(0, y) = A_L := (-1/\sqrt{2}, 1/\sqrt{2})$  and  $F_0(a, y) = A_R := -A_L = (1/\sqrt{2}, -1/\sqrt{2})$  for  $0 \leq y \leq b$ . On the lower boundary,  $F_0(x, 0) = (\xi, \eta)$  turns counterclockwise in the half-plane  $\xi + \eta \leq 0$  with constant angular velocity from  $A_L$  to  $A_R$  as  $x$  varies from 0 to  $b$ . The upper boundary is similar, but there  $F_0(x, b)$  changes clockwise in the half-plane  $\xi + \eta \geq 0$ . Then it follows from the assumptions that, for  $(x, y) \in \partial P$ ,  $F(x, y)$  and  $F_0(x, y)$  never point to opposite directions. Hence,  $F(x, y)/\|F(x, y)\|, F_0(x, y): \partial P \rightarrow \mathbb{S}^1$  are homotopic. Therefore, the index of  $F$  equals the index of  $F_0$ , namely 1.  $\square$

We still need two lemmas that together form a sharpening of two lemmas from [11].

**Lemma 8** ([11, Lemmas 1 and 2]). *Consider a (possibly degenerate) triangle  $A$  in  $\mathbb{S}^2$ ,  $\mathbb{R}^2$  or  $\mathbb{H}^2$  with sides  $a, b, x$ , where  $a, b > 0$ . For the case of  $\mathbb{S}^2$ , we additionally assume  $a + b \leq \pi$ . Then, for  $a, b$  fixed and  $|a - b| \leq x \leq a + b$ , the area  $\overline{A}$  of this triangle is a concave function of  $x$ . (For  $x = a + b = \pi$  on  $\mathbb{S}^2$ , we define  $\overline{A}$  by a limit procedure: namely, fixing  $a, b$ , we let  $x \rightarrow a + b = \pi$ . Accordingly, we set  $\overline{A} = \pi$ . Observe that for  $a + b = \pi$ , the area  $\overline{A}$  is half the area of a digon with sides containing the sides  $a, b$  of  $A$ .) In addition, the area is strictly concave for  $\mathbb{R}^2$  and  $\mathbb{H}^2$  and, under the additional constraint  $a + b < \pi$ , also for  $\mathbb{S}^2$ .  $\square$*

We calculate more precise details about this concave function and the value of its maximum.

**Lemma 9.** *We use the notations and hypotheses of Lemma 8 and denote by  $\gamma$  the angle between the sides  $a, b$ . For  $\mathbb{S}^2$ , let us additionally assume  $a + b < \pi$ . Then  $\overline{A}$  equals 0 for  $x = |a - b|$  and  $x = a + b$ , and it has a unique maximum for some value  $x = x_{\max}$ , with corresponding angle  $\gamma = \gamma_{\max}$ .*

*For  $\mathbb{H}^2$ , we have*

$$\begin{aligned} \cosh(x_{\max}/2) &= \sqrt{(\cosh a + \cosh b)/2}, \\ \cos \gamma_{\max} &= \tanh(a/2) \cdot \tanh(b/2), \end{aligned}$$

*and the value of the maximal area is*

$$\begin{aligned} \pi - 2 \arcsin \frac{\sinh(a/2)}{\sinh r} - 2 \arccos \frac{\tanh(a/2)}{\tanh r} + \\ \pi - 2 \arcsin \frac{\sinh(b/2)}{\sinh r} - 2 \arccos \frac{\tanh(b/2)}{\tanh r}, \end{aligned}$$

*where  $\cosh r = \sqrt{(\cosh a + \cosh b)/2}$ .*

*For  $\mathbb{R}^2$ , we have*

$$x_{\max}^2 = a^2 + b^2, \quad \gamma_{\max} = \pi/2,$$

*and the maximal area is  $ab/2$ .*

*For  $\mathbb{S}^2$ , we have*

$$\begin{aligned} \cos(x_{\max}/2) &= \sqrt{(\cos a + \cos b)/2}, \\ \cos \gamma_{\max} &= -\tan(a/2) \cdot \tan(b/2), \end{aligned}$$

*and the maximal area is*

$$\begin{aligned} 2 \arcsin \frac{\sin(a/2)}{\sin r} + 2 \arccos \frac{\tan(a/2)}{\tan r} - \pi + \\ 2 \arcsin \frac{\sin(b/2)}{\sin r} + 2 \arccos \frac{\tan(b/2)}{\tan r} - \pi, \end{aligned}$$

*where  $\cos r = \sqrt{(\cos a + \cos b)/2}$ .*

*Moreover, letting  $y/2$  be the distance between the midpoint of the side  $x$  and the common vertex of the sides  $a$  and  $b$ , we have the following equivalences:*

$$\gamma \in [0, \gamma_{\max}) \iff x < y, \quad \text{and } \gamma = \gamma_{\max} \iff x = y, \quad \text{and } \gamma \in (\gamma_{\max}, \pi] \iff x > y.$$

*Proof.* For  $\mathbb{R}^2$ , the statement is elementary. Therefore, we investigate only the cases of  $\mathbb{H}^2$  and  $\mathbb{S}^2$ .

Denote the vertices of the triangle opposite to the sides  $a, b$  and  $x$ , by  $A, B$  and  $C$ , respectively. Let  $D$  be the mirror image of  $C$  with respect to the midpoint of the

side  $x$ . Then the quadrilateral  $ABCD$  is centrally symmetric with respect to the intersection  $O$  of its diagonals  $BC$  (of length  $x$ ) and  $AD$  (of length  $y$ ). Its area is  $2\overline{A}$ , so it suffices to investigate its area.

We recall the isoperimetric property of the circle in  $\mathbb{R}^2$ ,  $\mathbb{H}^2$ , and on  $\mathbb{S}^2$  — but in the last case of radius  $r < (a+b)/2 < \pi/2$  — among sets of equal perimeter. Namely, that the maximum area is attained for the circle. For  $\mathbb{S}^2$ , one must restrict the candidate to (closed) sets contained in some open half- $\mathbb{S}^2$  [52, Ch. 18, §6, 2, (18.39)]. Observe that a piecewise  $C^1$  closed curve on  $\mathbb{S}^2$  with length less than  $2\pi$  lies in some open half- $\mathbb{S}^2$ , by elementary integral-geometric considerations [52, Ch. 7, §2, (7.11) and Ch. 18, §6, 1, (18.37)]. (A very detailed exposition of the isoperimetric inequality in spaces of constant curvature, i.e., in  $\mathbb{R}^n$ ,  $\mathbb{H}^n$ , and  $\mathbb{S}^n$ , can be found in [53]. See [54] for further details.)

For  $\gamma = 0$ , we have  $x = |a - b| < a + b = y$ , while for  $\gamma = \pi$ , we have  $x = a + b > |a - b| = y$ . Therefore, for some  $\gamma \in (0, \pi)$ , we have  $x = y$ . This implies that, for this  $\gamma$ , i.e., for this  $x$ ,  $ABCD$  is inscribed in a circle of centre  $O$  and radius  $r := x/2 = y/2$ . By the isoperimetric property of the circle — on  $\mathbb{S}^2$  of radius  $r < (a+b)/2 < \pi/2$ , in the sense described above — this value of  $\gamma$  must therefore be  $\gamma_{\max}$ , and this  $x$  is  $x_{\max}$ , see [33, p. 63, Problem 21], [32, §5, Problem 63], [34, p. 52]. (These references deal with the case of  $\mathbb{R}^2$ . However, their well-known proof carries over to  $\mathbb{H}^2$  and  $\mathbb{S}^2$  if we use the isoperimetric property of the circle — on  $\mathbb{S}^2$  of radius  $r < (a+b)/2 < \pi/2$ , in the sense described above.)

We determine the radius of this circle. We use the law of cosines for the triangles  $\triangle AOC$ ,  $\triangle BOC$  and write  $\varphi := \angle BOC$ . For  $\mathbb{H}^2$ , we have

$$\cosh a = \cosh(x/2) \cdot \cosh(y/2) - \sinh(x/2) \cdot \sinh(y/2) \cdot \cos \varphi,$$

and

$$\cosh b = \cosh(x/2) \cdot \cosh(y/2) + \sinh(x/2) \cdot \sinh(y/2) \cdot \cos \varphi.$$

Adding these, we obtain

$$\cosh a + \cosh b = 2 \cosh(x/2) \cdot \cosh(y/2). \quad (26)$$

Analogously, for  $\mathbb{S}^2$ , we obtain

$$\cos a + \cos b = 2 \cos(x/2) \cdot \cos(y/2).$$

(These are the analogues of the parallelogram law in  $\mathbb{R}^2$ .) Thus, for  $\mathbb{H}^2$ , we have

$$\cosh a + \cosh b = 2 \cosh^2 r = 2 \cosh^2(x_{\max}/2),$$

and, for  $\mathbb{S}^2$ , we have

$$\cos a + \cos b = 2 \cos^2 r = 2 \cos^2(x_{\max}/2)$$

in the range  $0 < r < (a+b)/2 < \pi/2$ .

Furthermore, for  $\mathbb{H}^2$ ,  $x$  is a strictly increasing function of  $\gamma$  and, by (26),  $y$  is a strictly decreasing function of  $x$ . Hence, for  $x = 2r$  we have  $y = 2r$ , for  $|a-b| \leq x < 2r$  we have  $2r < y \leq a+b$ , and similarly, for  $2r < x \leq a+b$  we have  $|a-b| \leq y < 2r$ . These imply the last equivalences in the lemma for  $\mathbb{H}^2$ .

Next we determine  $\cos \gamma_{\max}$  for  $\mathbb{H}^2$ . The law of cosines for the triangle  $\triangle ABC$  gives

$$\cos \gamma_{\max} = \frac{\cosh a \cdot \cosh b - \cosh(2r)}{\sinh a \cdot \sinh b}.$$

In this equation, we have  $\cosh(2r) = 2\cosh^2 r - 1 = \cosh a + \cosh b - 1$ , and this implies the formula in the lemma. (Observe that  $0 < a, b$  implies  $\cos \gamma_{\max} = \tanh(a/2) \cdot \tanh(b/2) \in (0, 1)$ .)

For  $\mathbb{S}^2$ , the proof of the last equivalences in the lemma and the calculation of  $\cos \gamma_{\max}$  are analogous. (Observe that now  $0 < a, b$  and  $a/2 + b/2 < \pi/2$  imply  $\cos \gamma_{\max} = -\tan(a/2) \cdot \tan(b/2) \in (-1, 0)$ .)

Finally, the value of the maximum follows from the trigonometric formulas for a right triangle in  $\mathbb{H}^2$  and  $\mathbb{S}^2$ .  $\square$

We prove Theorem 3 with the following

**Construction 1.** Consider a number

$$S \in (0, \pi/2) \tag{27}$$

and a number  $t > 0$  such that

$$2\sinh(t/2) > \tan S.$$

(Later,  $S$  will be the area of a compact right triangle, which explains condition (27). At the same time, this explains the hypothesis  $0 < S_4 < \pi/2$  of Theorem 3, since in the proof,  $S$  will be chosen for example as  $S_4$ .)

Now we define a function

$$f_{t,S}: [0, t] \rightarrow \mathbb{R}$$

as follows. For any  $x \in [0, t]$ , consider the function

$$g_x(y) := \arctan \frac{\sinh x \cdot \sinh y}{\cosh x + \cosh y} + \arctan \frac{\sinh(t-x) \cdot \sinh y}{\cosh(t-x) + \cosh y},$$

where  $y \in [0, \infty)$ . It is easy to see that  $(d/dy)g_x(y) > 0$  for  $y \in [0, \infty)$ , and  $g_x(0) = 0$ , and

$$\begin{aligned} \lim_{y \rightarrow \infty} g_x(y) &= \arctan(\sinh x) + \arctan(\sinh(t-x)) \\ &\geq \arctan(\sinh x + \sinh(t-x)) \geq \arctan(2\sinh(t/2)) > S \end{aligned}$$

for all  $x \in [0, t]$ . Here, at the first inequality, we used concavity of the function  $\arctan y$  on  $[0, \infty)$  and  $\arctan 0 = 0$ . At the second inequality, we used convexity of the function  $\sinh x$  on the interval  $[0, t]$ . Therefore, there is a unique  $\tilde{y} \in (0, \infty)$  such that  $g_x(\tilde{y}) = S$ . We put

$$f_{t,S}(x) := \tilde{y} \in (0, \infty). \tag{28}$$

Now we investigate some properties of this function. Obviously,  $f_{t,S}$  is continuous on  $[0, t]$  (moreover, it is  $C^1$  on  $(0, t)$ ), and  $f_{t,S}(x) = f_{t,S}(t-x)$ .

Here is the geometric interpretation of  $f_{t,S}$ . Consider a triangle  $\Delta ABC \subset \mathbb{H}^2$  with the following properties:

- (1)  $|AB| = t$ ,
- (2) the area of  $\Delta ABC$  is  $S$ ,
- (3) if  $H$  is the orthogonal projection of  $C$  to the line  $\ell(A, B)$ , then  $H \in [A, B]$  and  $|AH| = x \in [0, t]$ .

Then it is easy to see (using Lemma 5) that  $|CH| = f_{t,S}(x)$ . It is also easy to see that for  $0 < \tilde{S} < S$  and for every  $x \in [0, t]$ , we have  $f_{t,\tilde{S}}(x) < f_{t,S}(x)$ .

In what follows we determine the number

$$h_{t,S} := f_{t,S}(0) = f_{t,S}(t). \quad (29)$$

By Lemma 5, we have

$$\tan S = \frac{\sinh t \cdot \sinh h_{t,S}}{\cosh t + \cosh h_{t,S}}.$$

Solving this equation for  $\cosh h_{t,S}$ , we get

$$\cosh h_{t,S} = \frac{\tan^2 S \cdot \cosh t + \sqrt{1 + \tan^2 S} \cdot \sinh^2 t}{\sinh^2 t - \tan^2 S}.$$

(Observe that  $\sinh t > 2 \sinh(t/2) > \tan S$  by the strict convexity of the function  $\sinh t$  on  $[0, \infty)$  and  $\sinh 0 = 0$ . Therefore, the denominator in this formula is positive.) From this, we see that  $\cosh h_{t,S} \rightarrow 1/\cos S$  for  $t \rightarrow \infty$ .

*Proof of Theorem 3. 1.* First we consider the case when hypothesis (1) of Theorem 3 holds.

Let us take a  $t > 0$  such that

$$2 \sinh(t/2) > \tan S_4, \quad (30)$$

and for the number  $h_{t,S_4} := f_{t,S_4}(0) = f_{t,S_4}(t)$  defined in (28) and (29), we have

$$\frac{\cosh h_{t,S_4} - 1}{2\sqrt{\cosh h_{t,S_4}}} < \tan(S_1/2). \quad (31)$$

Such a  $t$  exists since  $\cosh h_{t,S_4} \rightarrow 1/\cos S_4$  for  $t \rightarrow \infty$ , and

$$\lim_{t \rightarrow \infty} \frac{\cosh h_{t,S_4} - 1}{2\sqrt{\cosh h_{t,S_4}}} = \frac{1 - \cos S_4}{2\sqrt{\cos S_4}} < \tan(S_1/2),$$

by hypothesis (1) of the theorem.

Consider any plane  $\sigma$  in  $\mathbb{H}^3$ . Take points  $A_1, A_2 \in \sigma$  such that  $|A_1 A_2| = t$ , where  $t$  has been chosen above. Let  $\sigma^+$  and  $\sigma^-$  be the two half-planes bounded by the line  $\ell(A_1, A_2)$  in  $\sigma$ .

For a given  $x \in [0, t]$ , we take the point  $H = H(x)$  on the segment  $[A_1, A_2]$  satisfying  $|A_1 H| = x$ . Consider the half-line  $l_x$  from  $H$  in  $\sigma^+$  that is orthogonal to the line  $\ell(A_1, A_2)$ . Now let  $\tilde{A}_4 = \tilde{A}_4(x)$  be the point on  $l_x$  satisfying  $|\tilde{A}_4 H| = f_{t,S_3}(x)$ , and let  $\tilde{A}_3 = \tilde{A}_3(x)$  be the point on  $l_x$  satisfying  $|\tilde{A}_3 H| = f_{t,S_4}(x)$ . By  $S_4 \geq S_3$ , we have  $\tilde{A}_4 \in [\tilde{A}_3, H]$ , and thus also  $|H \tilde{A}_4| \leq |H \tilde{A}_3|$ .

Let  $\{\gamma(\varphi)\}$  be the one-parameter group of rotations about the line  $\ell(A_1, A_2)$  through the angles  $\varphi$ , in some definite sense of rotation, with  $\gamma(0)$  being the identity. For  $\varphi \in [0, \pi]$  we consider the point  $A_3 = A_3(x, \varphi) := \gamma(\varphi)(\tilde{A}_3(x))$ . Note that  $A_3(x, \pi) \in \sigma^-$ . Consider also  $A_4(x, \varphi) := \tilde{A}_4(x)$ ,  $A_1(x, \varphi) := A_1$  and  $A_2(x, \varphi) := A_2$ .

We are going to prove that there exists an  $(x, \varphi) \in [0, t] \times [0, \pi]$  such that the (possibly degenerate) tetrahedron  $T = T(x, \varphi) := A_1(x, \varphi)A_2(x, \varphi)A_3(x, \varphi)A_4(x, \varphi)$  has facet areas  $S_1, S_2, S_3, S_4$ , where  $S_i$  is the area of the facet opposite to  $A_i(x, \varphi)$ .

Let  $s_i(x, \varphi)$  be the area of the facet of  $T(x, \varphi)$  that is opposite to the vertex  $A_i = A_i(x, \varphi)$ . By our construction, we obviously have  $s_4(x, \varphi) = S_4$  and  $s_3(x, \varphi) = S_3$ .

Let us define functions  $f_1, f_2: [0, t] \times [0, \pi] \rightarrow \mathbb{R}$  as follows.

$$f_1(x, \varphi) = s_2(x, \varphi) - S_2, \quad f_2(x, \varphi) = s_1(x, \varphi) - S_1. \quad (32)$$

It is easy to see that

$$f_1(x, 0) + f_2(x, 0) = S_4 - S_1 - S_2 - S_3 < 0, \quad (33)$$

$$f_1(x, \pi) + f_2(x, \pi) = S_3 + S_4 - S_1 - S_2 \geq 0. \quad (34)$$

Now we check that

$$f_1(0, \varphi) < 0, \quad f_2(t, \varphi) < 0, \quad (35)$$

for all  $\varphi \in [0, \pi]$ .

For the first inequality in (35), we note that the point  $\tilde{A}_3(0)$  satisfies  $|\tilde{A}_3(0)A_1| = f_{t,S_4}(0) = h_{t,S_4}$ , and the point  $\tilde{A}_4(0)$  satisfies  $|\tilde{A}_4(0)A_1| = f_{t,S_3}(0) =: h_{t,S_3} \leq h_{t,S_4}$ . Therefore, for every  $\varphi \in [0, \pi]$ , the triangle  $\Delta A_1 A_3(0, \varphi) A_4(0, \varphi)$  satisfies  $|A_1 A_3(0, \varphi)| = h_{t,S_4}$  and  $|A_1 A_4(0, \varphi)| \leq h_{t,S_4}$ . By Lemma 6 and (31), we get

$$s_2(0, \varphi) \leq 2 \arctan \frac{\cosh h_{t,S_4} - 1}{2\sqrt{\cosh h_{t,S_4}}} < S_1 \leq S_2.$$

Therefore,  $f_1(0, \varphi) = s_2(0, \varphi) - S_2 \leq s_2(0, \varphi) - S_1 < 0$  for all  $\varphi \in [0, \pi]$ .

For the second inequality of (35), we replace in the above argument  $A_1, \tilde{A}_3(0)$ , and  $\tilde{A}_4(0)$  by  $A_2, \tilde{A}_3(t)$ , and  $\tilde{A}_4(t)$ , respectively. We get  $f_2(t, \varphi) = s_1(t, \varphi) - S_1 < 0$  for all  $\varphi \in [0, \pi]$ .

Taking into account the inequalities (33–35), by applying Lemma 7 with  $(u_1, u_2, v_1, v_2) := (1, 0, 0, 1)$ , we find an  $(x, \varphi) \in [0, t] \times [0, \pi]$  such that  $f_1(x, \varphi) = f_2(x, \varphi) = 0$ . This means that  $s_1(x, \varphi) = S_1$  and  $s_2(x, \varphi) = S_2$  for the corresponding (possibly degenerate) tetrahedron  $T$ .

**2.** Now we consider the case when hypothesis (2) of Theorem 3 holds. We use the same construction of the tetrahedron  $T$  as in the first case. For the functions  $f_1$  and  $f_2$  defined by (32), we get the inequalities (33) and (34). Now we check that

$$f_2(0, \varphi) \geq 0, \quad f_1(t, \varphi) \geq 0 \quad (36)$$

for all  $\varphi \in [0, \pi]$ . For this we note that

$$s_1(0, \varphi) \geq s_1(0, 0) = S_4 - S_3 \geq S_2 \geq S_1 \quad (37)$$

and

$$s_2(t, \varphi) \geq s_2(t, 0) = S_4 - S_3 \geq S_2 \quad (38)$$

for all  $\varphi \in [0, \pi]$  provided  $t$  is sufficiently large, as we will prove. Of course, we have to prove only the first inequalities in (37) and (38).

We will investigate  $s_1(0, \varphi)$ . (The case of  $s_2(t, \varphi)$  is analogous.) Recall that  $|H\tilde{A}_4| \leq |H\tilde{A}_3|$ , which implies  $h_{t,S_3} = |A_1(0, \varphi)A_4(0, \varphi)| \leq |A_1(0, \varphi)A_3(0, \varphi)| = h_{t,S_4}$ . For  $t$  fixed but  $\varphi \in [0, \pi]$  variable, the length of the third side of the triangle  $\Delta A_1(0, \varphi)A_3(0, \varphi)A_4(0, \varphi)$  lies in the range

$$|A_3(0, \varphi)A_4(0, \varphi)| \in [h_{t,S_4} - h_{t,S_3}, h_{t,S_4} + h_{t,S_3}].$$

Therefore, to show

$$s_1(0, \varphi) \geq s_1(0, 0), \quad (39)$$

we must show the following. Let  $a := |A_2(0, \varphi)A_4(0, \varphi)|$  and  $b := |A_2(0, \varphi)A_3(0, \varphi)|$ . Then  $t \leq a \leq b$ , since

$$\begin{cases} \cosh t \leq \cosh a = \cosh t \cdot \cosh |A_1(0, \varphi)A_4(0, \varphi)| \\ \leq \cosh t \cdot \cosh |A_1(0, \varphi)A_3(0, \varphi)| = \cosh b. \end{cases}$$

Let  $c := |A_1(0, \varphi)A_3(0, \varphi)| - |A_1(0, \varphi)A_4(0, \varphi)| = h_{t,S_4} - h_{t,S_3}$ . Then the area of the triangle with sides  $a, b, c$  is less than or equal to the area of the triangle with the same first two sides  $a, b$  and with third side in the interval

$$[h_{t,S_4} - h_{t,S_3}, h_{t,S_4} + h_{t,S_3}] \subset [h_{t,S_4} - h_{t,S_3}, 2h_{t,S_4}] \subset [h_{t,S_4} - h_{t,S_3}, \text{const}].$$

For the last inclusion observe the following. By the geometric interpretation, if  $S_4$  is fixed and  $t$  is above the bound  $2 \operatorname{arsinh}[(\tan S_4)/2]$  from (30) and increases, then  $h_{t,S_4}$  decreases. Therefore,  $h_{t,S_4}$  remains bounded for fixed  $S_4$  if  $t$  increases from its originally chosen value  $t_0$ , say, to infinity.

Inequality (39) is proved if we show the following monotonicity property. Fixing the first two sides  $a, b$  and varying the third side  $x$  in the interval  $[h_{t,S_4} - h_{t,S_3}, \text{const}]$ , the area is a monotonically increasing function of  $x$ .

Now we apply Lemmas 8 and 9 to the triangle with sides  $a, b, x$ . We need to show that its area is increasing for  $x \in [b - a, \text{const}]$ , where we know from the preceding considerations that  $0 \leq b - a \leq \text{const}$ . By these Lemmas, this area-increasing property is satisfied for  $x \in [b - a, x_{\max}]$ , where  $x_{\max}$  is defined by  $\cosh^2(x_{\max}/2) = (\cosh a + \cosh b)/2$ . Thus, to complete the argument, it suffices to show that  $x_{\max} \geq \text{const}$ , i.e., that  $x_{\max} \rightarrow \infty$  for  $t \rightarrow \infty$ .

We estimate  $x_{\max}$  from below. We have

$$\cosh^2(x_{\max}/2) = (\cosh a + \cosh b) / 2 \geq \cosh a > e^a / 2,$$

hence

$$(e^{x_{\max}/2})^2 > \cosh^2(x_{\max}/2) > e^a / 2,$$

and hence

$$x_{\max} > a - \log 2 \geq t - \log 2 \rightarrow \infty,$$

as we wanted to show. Thus, (36) is proved.

Taking in account inequalities (33), (34), and (36), we can apply Lemma 7 with  $(u_1, u_2, v_1, v_2) = (0, -1, -1, 0)$  to find a point  $(x, \varphi) \in [0, t] \times [0, \pi]$  such that  $f_1(x, \varphi) = f_2(x, \varphi) = 0$ . This means that  $s_1(x, \varphi) = S_1$  and  $s_2(x, \varphi) = S_2$  for the corresponding (possibly degenerate) tetrahedron  $T$ .

**3.** It remains to exclude degeneration of our tetrahedron. Our construction yields degenerate tetrahedra only for  $\varphi = 0$  and  $\varphi = \pi$ . In the first case,  $S_4 = S_1 + S_2 + S_3$ , which contradicts our hypotheses. In the second case,  $S_4 + S_3 = S_2 + S_1$ , which implies  $\pi > S_4 = S_3 = S_2 = S_1 > 0$ . (By the way, this can occur only for case (1) of the theorem.) Then a suitable regular tetrahedron satisfies the conclusion of the theorem.  $\square$

*Remark 7.* Let us apply the construction in the proof of Theorem 3 to the numbers  $S_i \varepsilon^2$  and  $t \varepsilon$  rather than  $S_i$  and  $t$ , where  $\varepsilon \rightarrow 0$ . Then, for sufficiently small  $\varepsilon > 0$ , hypothesis (1) from Theorem 3 holds, and as an analogue of (30) we have  $2 \sinh(t \varepsilon / 2) > \tan(S_4 \varepsilon^2)$ . In the limit, we obtain a Euclidean tetrahedron with facet areas  $S_i$  and one edge of length  $t$ . Letting  $t \rightarrow \infty$  gives another proof for the last statement of Theorem E for  $\mathbb{R}^3$  (existence of tetrahedra of arbitrarily small positive volume). Namely, the heights of the two facets meeting at the edge of length  $t$ , corresponding to this edge, are  $O(1/t)$ . Thus, the tetrahedron is included in a right circular cylinder of height  $t$  and radius  $O(1/t)$ . Hence, the volume of the tetrahedron is  $O(1/t)$ . Degeneration is excluded as in Step 3 of the above proof of Theorem 3.



## 6. PROOFS FOR THE SPHERICAL CASE

Recall our convention about the notion of *simplices in  $\mathbb{S}^n$*  at the beginning of Section 2.3.

**6.1. Proof of Proposition 4. 1.** We begin with the proof of the first inequality. Let the facets of  $P$  be  $F_i$ . Their areas satisfy the equation

$$S_i = \text{const}_n \cdot \int |F_i \cap \mathbb{S}^1| d\mathbb{S}^1. \quad (40)$$

Here  $|\cdot|$  denotes cardinality and  $\text{const}_n > 0$ . The integration is taken with respect to the unique  $O(n+1)$ -invariant probability Borel measure (for the standard embedding  $\mathbb{S}^n \subset \mathbb{R}^{n+1}$ ) on the manifold of all great- $\mathbb{S}^1$ 's in  $\mathbb{S}^n$  [52, Ch. 18, §6, 1].

In the integration, we may disregard those  $\mathbb{S}^1$ 's that lie in the great- $\mathbb{S}^{n-1}$ 's spanned by the facets  $F_1, \dots, F_m$ , since they have measure 0. By the same reason, we may disregard those  $\mathbb{S}^1$ 's that pass through the relative boundary of  $F_i$  (in the great- $\mathbb{S}^{n-1}$  spanned by it) for all  $i \in \{1, \dots, m\}$  simultaneously. If an  $\mathbb{S}^1$  does not lie in the above great  $\mathbb{S}^{n-1}$ 's and does not intersect the above relative boundaries then it cannot contain two opposite points of any  $F_i$ . Namely, since  $F_i$  lies in a closed half- $\mathbb{S}^{n-1}$ , both of these points would otherwise lie in the relative boundary of  $F_i$  taken with respect to the great- $\mathbb{S}^{n-1}$  spanned by it. If such an  $\mathbb{S}^1$  enters  $P$  at some point  $p \in F_m$  it must also leave  $P$ , through some other facet (since this  $\mathbb{S}^1$  does not contain two opposite points of  $F_m$ ) till it comes back to  $p$ . This holds even in the degenerate case, that is, when some portion of  $F_m$  is a doubly counted boundary of  $P$  either as a “flat” piece of  $P$  or as bounded from both sides by the interior of  $P$ .

These considerations imply that the integral (40) for  $i = m$  is at most the sum of the integrals for all  $1 \leq i \leq m - 1$ . Thus, (40) gives our inequality.

Now assume that  $P$  lies in an open half- $\mathbb{S}^n$  (in the open northern hemisphere, say) but does not degenerate to the doubly-counted facet  $F_m$ .

Let  $\mathbb{S}_m$  be the great- $\mathbb{S}^{n-1}$  spanned by  $F_m$ . If  $\cup_{i=1}^{m-1} F_i \not\subset \mathbb{S}_m$ , then there exists an  $x \in \cup_{i=1}^{m-1} \text{relint } F_i$  such that  $x \notin \mathbb{S}_m$ . Also, there exists a  $y \in \mathbb{S}_m \setminus F_m$  that also lies in the open northern hemisphere. Then  $x \neq y$ , and since both  $x$  and  $y$  lie in the open northern hemisphere, the great- $\mathbb{S}^1$   $xy$  through these two points exists.

The set of  $\mathbb{S}^1$ 's transversally intersecting  $\cup_{i=1}^{m-1} \text{relint } F_i$  but not intersecting  $F_m$  contains some neighbourhood of the great- $\mathbb{S}^1$   $xy$  in the set of all great- $\mathbb{S}^1$ 's and has therefore positive measure. This implies the strict inequality in this case.

If  $\cup_{i=1}^{m-1} F_i \subset \mathbb{S}_m$ , then in both cases  $\cup_{i=1}^{m-1} F_i \not\subset F_m$  and  $\cup_{i=1}^{m-1} F_i \subset F_m$ , we have strict inequality unless  $P$  degenerates to the doubly counted facet  $F_m$ . However, this degeneration was excluded.

**2.** We turn to the proof of the second inequality. We again use the formula (40). Again we disregard those  $\mathbb{S}^1$ 's that lie in the great- $\mathbb{S}^{n-1}$ 's spanned by any facet  $F_i$  of  $P$ , as well as those  $\mathbb{S}^1$ 's that pass through the relative boundary of any facet  $F_i$  of  $P$ . We compare the sum of the right hand sides of (40) for all  $1 \leq i \leq m$  with the analogous integral when in the right hand side of (40) we take a great- $\mathbb{S}^{n-1}$  rather than  $F_i$ .

Clearly, for a great- $\mathbb{S}^{n-1}$ , the cardinality of its intersection with a great- $\mathbb{S}^1$  is almost always 2. For the  $\mathbb{S}^1$ 's that were not disregarded and for any  $i$ , the cardinality  $|F_i \cap \mathbb{S}^1|$  is at most 1, since a great- $\mathbb{S}^1$  cannot contain two opposite points of  $F_i$  (see part 1 of this proof). If  $P$  is not degenerate, one great- $\mathbb{S}^1$  cannot transversally intersect the

interiors of three facets  $F_i$ . Namely, at each of these intersection points, it passes either into  $P$  or out of  $P$  (with some definite orientation of our  $\mathbb{S}^1$ ). Thus, there would be at least four points of intersection, and the intersection of  $P$  and this great- $\mathbb{S}^1$  would be the union of at least two disjoint non-trivial arcs. However, this contradicts convexity of  $P$ . Hence, the sum of the integrands in (40) over all facets  $i = 1, \dots, m$  is at most 2. For degenerate  $P$ , the same inequality holds. This implies the second inequality of the proposition.

If  $P$  lies in an open half- $\mathbb{S}^n$ , then the set of  $\mathbb{S}^1$ 's intersecting the boundary of the open half- $\mathbb{S}^n$  but not intersecting  $\cup_{i=1}^m F_i$  has a positive measure. (Namely, any great- $\mathbb{S}^1$  sufficiently close to the boundary of the open half- $\mathbb{S}^n$  has this property.) This implies the strict inequality in this case.  $\square$

*Remark 8.* Part 1 of the above proof of Proposition 4 extends also for  $\mathbb{H}^n$  and yields Proposition 1, however without the case of equality. We have to use also [52, Ch. 18, §6, 1], and instead of  $\mathbb{S}^1$ , we have to take a segment of a fixed positive length  $t$  and then let  $t$  tend to infinity. However, we preferred to give the elementary proof for Proposition 1.

*Remark 9.* Clearly, the argument for the inequality  $t_1 \leq t_3$  in the proof of Proposition 3 is valid also for  $\mathbb{R}^3$  and  $\mathbb{S}^3$ . However, for  $\mathbb{R}^3$ , this inequality is easy to show, see below. It is also easy to show for  $\mathbb{S}^3$ , provided that each facet is contained in a closed half- $\mathbb{S}^2$  and has at least three sides — in particular if the polyhedron is contained in an open half- $\mathbb{S}^3$  — see below. For  $\mathbb{R}^3$ , the sum of the angles of the  $k_i$ -gon  $F_i$  is  $t_1 = (k_i - 2)\pi$ . Also,  $F_i$  has  $k_i$  neighbouring faces, each of which has angle sum at least  $\pi$ , so  $t_1 = (k_i - 2)\pi < k_i\pi \leq t_3$ . Similarly, for  $\mathbb{S}^3$ , with the above hypotheses, the sum of the angles of  $F_i$  is  $t_1 = S_i + (k_i - 2)\pi$ , while every other facet  $F_j$  has an angle sum  $S_j + (k_j - 2)\pi \geq S_j + \pi$ , since  $k_j \geq 3$ . So the sum of the angles of the other facets  $F_j$  is at least  $\sum_{j \neq i} (S_j + \pi)$ , and hence,  $\sum_{j \neq i} S_j + k_i\pi \leq t_3$ . Therefore,  $t_1 = S_i + (k_i - 2)\pi < S_i + k_i\pi \leq (\sum_{j \neq i} S_j) + k_i\pi \leq t_3$ . Here, we used the first inequality of Proposition 4, which implies  $S_i \leq \sum_{j \neq i} S_j$ , provided every facet lies in some closed half- $\mathbb{S}^2$ .

**6.2. Proof of Theorem 4.** Now we give the spherical analogues of Lemmas 5 and 6.

**Lemma 10.** *The area  $S$  of a right triangle  $\Delta ABC \subset \mathbb{S}^2$  with angle  $\angle ACB = \pi/2$  and side lengths  $|AC| = b$  and  $|BC| = a$  (where  $0 < a, b \leq \pi$ ) fulfills the equation*

$$\tan S = \frac{\sin a \cdot \sin b}{\cos a + \cos b}$$

*if  $a + b \neq \pi$ . For  $a + b = \pi$ , the area is  $S = \pi/2$ .*

**Lemma 11.** *Let  $0 < d \leq \pi/2$ . Assume that  $\Delta ABC \subset \mathbb{S}^2$  is a triangle such that  $|AB| \leq d$  and  $|AC| \leq d$ . Then the area  $S$  of this triangle is bounded above by the inequality*

$$S \leq 2 \arctan \frac{1 - \cos d}{2\sqrt{\cos d}}$$

*if  $d \neq \pi/2$ . For  $d = \pi/2$  we have  $S \leq \pi$ .*

*Proof of Lemmas 10 and 11.* The case  $a + b = \pi$  of Lemma 10 and the case  $d = \pi/2$  of Lemma 11 is elementary. In the remaining cases  $\cos d > 0$  for Lemma 11, and we proceed analogously as in Lemmas 5 and 6.  $\square$

To prove Theorem 4, we use an analogous construction as for Theorem 3.

**Construction 2.** Let

$$S \in (0, \pi/2], \quad (41)$$

and choose

$$t = \pi/2.$$

This choice is motivated as follows. We will apply Lemma 8 to triangles with sides  $a, b \leq \pi/2$ , but Lemma 8 does not hold for  $a = b \in (t, \pi) = (\pi/2, \pi)$ . Thus,  $t = \pi/2$  is the largest value for which our proof applies. Also,  $S$  will be the area of a spherical triangle contained in a spherical triangle with three sides  $\pi/2$ , which explains the constraint (41). Now, we define a function

$$f_{t,S}: [0, t] \rightarrow \mathbb{R}$$

as follows. For any  $x \in [0, t]$ , consider the function

$$g_x(y) := \arctan \frac{\sin x \cdot \sin y}{\cos x + \cos y} + \arctan \frac{\sin(t-x) \cdot \sin y}{\cos(t-x) + \cos y},$$

defined for  $y \in [0, \pi/2]$ . It is easy to see that  $(d/dy)g_x(y) > 0$  for  $y \in [0, \pi/2)$ ,  $g_x(0) = 0$ , and

$$g_x(\pi/2) = \pi/2 \geq S,$$

for all  $x \in [0, t]$ . Therefore, there exists a unique  $\tilde{y} \in (0, \pi/2]$  such that  $g_x(\tilde{y}) = S$ . We put

$$f_{t,S}(x) := \tilde{y} \in (0, \pi/2].$$

Now we investigate some properties of this function. Obviously  $f_{t,S}$  is continuous on  $[0, t]$  (moreover, is  $C^1$  on  $(0, t)$ ), and  $f_{t,S}(x) = f_{t,S}(t-x)$ .

Here is the geometric interpretation of  $f_{t,S}$ . Consider a triangle  $\Delta ABC \subset \mathbb{S}^2$  with the following properties.

- (1)  $|AB| = t$ ,
- (2) the area of  $\Delta ABC$  is  $S$ ,
- (3)  $C$  has an orthogonal projection  $H$  to the line  $\ell(A, B)$  such that  $H$  lies in the segment  $[A, B]$  and  $|AH| = x \in [0, t]$ . (Observe that there are at least two orthogonal projections of  $C$  to  $\ell(A, B)$ .)

Then it is easy to see (using Lemma 10) that  $|CH| = f_{t,S}(x)$ . It is also easy to see that for  $0 < \tilde{S} < S$  and for every  $x \in [0, t]$ , we have  $f_{t,\tilde{S}}(x) < f_{t,S}(x)$ .

The boundary values

$$h_{t,S} := f_{t,S}(0) = f_{t,S}(t)$$

are easy to determine. By the geometric interpretation, this is the third side of a spherical triangle with two other sides of length  $\pi/2$  and area  $S$ , i.e.,

$$h_{t,S} = S.$$

*Proof of Theorem 4. 1.* First we consider the case when hypothesis (3) of Theorem 4 holds. We roughly follow the lines of the proof of Theorem 3 for the analogous case when hypothesis (1) holds.

We have

$$\frac{1 - \cos h_{t,S_4}}{2\sqrt{\cos h_{t,S_4}}} = \frac{1 - \cos S_4}{2\sqrt{\cos S_4}} \leq \tan(S_1/2), \quad (42)$$

by the hypothesis of the theorem. From this point onwards, the construction is the same as in Theorem 3.

We still have to show that our tetrahedron  $T$  satisfies our convention about simplices in  $\mathbb{S}^n$  (see the beginning of Section 2.3). A convex combinatorial simplex in an *open* half- $\mathbb{S}^n$  is always considered as a simplex in  $\mathbb{S}^n$ . Let  $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4$  denote the usual unit basis vectors, and let  $x_1, x_2, x_3, x_4$  denote the corresponding coordinates. We set  $A_1 := \mathbf{e}_1$  and  $A_2 := \mathbf{e}_2$ . The rotation about  $\ell(A_1, A_2)$  in  $\mathbb{S}^3$  maps  $A_3 := \mathbf{e}_3$  to  $\mathbf{e}_3 \cos \varphi + \mathbf{e}_4 \sin \varphi$ , say. Then  $T$  is in the closed half- $\mathbb{S}^3$  defined by the inequality  $x_1 + x_2 \geq 0$ . Moreover, if  $S_3 \leq S_4 < \pi/2$ , then  $T$  is contained in the open half- $\mathbb{S}^3$  defined by the inequality  $x_1 + x_2 > 0$ , and we are done. If  $S_3 \leq S_4 = \pi/2$ , a slight perturbation of the open half- $\mathbb{S}^3$  given by  $x_1 + x_2 > 0$  contains  $T$  for all  $\varphi \in [0, \pi]$ , and we are done. If  $S_3 = S_4 = \pi/2$ , and  $0 \leq \varphi < \pi/2$  is *fixed*, then also a slight perturbation of the open half- $\mathbb{S}^3$  given by  $x_1 + x_2 > 0$  contains  $T$  and then we are also done. The case  $S_3 = S_4 = \varphi = \pi/2$  will be treated in part 3 below.

We have to observe that from the construction, we have  $|A_3(x, \varphi)A_4(x, \varphi)| \leq f_{t,S_3}(x) + f_{t,S_4}(x) \leq 2f_{t,S_4}(x) \leq \pi$ . Thus, the edge  $[A_3(x, \varphi), A_4(x, \varphi)]$  of our tetrahedron is in the closed angular domain swept by  $\gamma(\varphi)\sigma^+$  for  $\varphi \in [0, \pi]$ , as in the hyperbolic case. (This explains the inequality  $S_4 \leq \pi/2$  of the theorem — and thus also the inequality  $S \leq \pi/2$  in the construction: without this inequality of the theorem, the last sentence would not be valid. Moreover, let  $S_3 = S_4 \in (\pi/2, \pi)$ . Then consider  $|A_3(x, \varphi)A_4(x, \varphi)|$  defined not as a distance but defined by analytic continuation from  $\varphi$ 's close to 0, i.e., by retaining the geometry of the figure. Then for  $\varphi = \pi$  we would have  $|A_3(x, \varphi)A_4(x, \varphi)| = 2f_{t,S_4}(x) \in (\pi, 2\pi)$ . This would imply that the tetrahedron  $T$  defined by the same analytic continuation, i.e., by retaining the geometry of the figure, would not be convex.)

We define  $s_i(x, \varphi)$  (for  $1 \leq i \leq 4$ ) and  $f_i(x, \varphi)$  (for  $i = 1, 2$ ) as in the proof of Theorem 3. The formulas

$$f_1(x, 0) + f_2(x, 0) = S_4 - S_1 - S_2 - S_3 < 0, \quad (43)$$

$$f_1(x, \pi) + f_2(x, \pi) = S_3 + S_4 - S_1 - S_2 \geq 0 \quad (44)$$

follow like (33) and (34) in the hyperbolic case. The formulas

$$f_1(0, \varphi) \leq 0, \quad f_2(t, \varphi) \leq 0,$$

for all  $\varphi \in [0, \pi]$  follow from Lemma 11 similarly as in the hyperbolic case from Lemma 6. (Observe that here we have non-strict inequalities. Namely, in hypothesis (3) of Theorem 4 and in (42), we have non-strict inequalities, whereas for the hyperbolic case, we had strict inequalities in hypothesis (1) of Theorem 3 and in (31).) Then, we choose  $(u_1, u_2, v_1, v_2) := (1, 0, 0, 1)$  and finish the proof of case 1 as in the hyperbolic case. Also here, the tetrahedron  $T$  is possibly degenerate. (Observe that allowing  $S_4 > \pi/2$ , we could have  $h_{t,S_4} > \pi/2$ . Then  $|A_1A_3(0, \varphi)|, |A_1A_4(0, \varphi)| \leq h_{t,S_4}$  makes it impossible to apply Lemma 11. This

explains once more the inequality  $S_4 \leq \pi/2$  of the theorem — and thus also the inequality  $S \leq \pi/2$  in the construction — for case **1**.)

**2.** Now we consider the case when hypothesis (4) of Theorem 4 holds. We roughly follow the lines of the proof of Theorem 3 when hypothesis (2) holds.

As in Step 1 of this proof, our tetrahedron satisfies our convention about the notion of a simplex in  $\mathbb{S}^n$  (defined the beginning of Section 2.3), unless  $S_3 = S_4 = \varphi = \pi/2$ . This last case will be handled below in part **3** of this proof.

We obtain the inequalities (43) and (44) similarly to (33) and (34) in the hyperbolic case.

Now we check that

$$f_2(0, \varphi) \geq 0, \quad f_1(t, \varphi) \geq 0,$$

for all  $\varphi \in [0, \pi]$ . As in the hyperbolic case, this reduces to showing that

$$s_1(0, \varphi) \geq s_1(0, 0), \quad s_2(t, \varphi) \geq s_2(t, 0). \quad (45)$$

We will investigate  $s_1(0, \varphi)$ . (The case of  $s_2(t, \varphi)$  is analogous.) Observe that the distance  $|A_3(x, \varphi)A_4(x, \varphi)|$  is a strictly increasing function of  $\varphi \in [0, \pi]$ , with

$$|A_3(x, 0)A_4(x, 0)| = h_{t, S_4} - h_{t, S_3} = S_4 - S_3,$$

and

$$|A_3(x, \pi)A_4(x, \pi)| = h_{t, S_4} + h_{t, S_3} = S_4 + S_3.$$

By the geometric interpretation (observe that  $|A_2(0, \varphi)A_1(0, \varphi)| = |A_2(0, \varphi)A_3(0, \varphi)| = |A_2(0, \varphi)A_4(0, \varphi)| = \pi/2$ ), we have

$$s_1(0, \varphi) = \angle A_3(0, \varphi)A_2(0, \varphi)A_4(0, \varphi) = |A_3(0, \varphi)A_4(0, \varphi)|,$$

where the last term is strictly increasing for  $\varphi \in [0, \pi]$ . This shows (45). Then, as in the hyperbolic case, we choose  $(u_1, u_2, v_1, v_2) := (0, -1, -1, 0)$  and finish the proof of **2**. Also here, the tetrahedron  $T$  can be degenerate.

**3.** It remains

- 1) to exclude degeneration of our tetrahedron, and
- 2) to verify that our tetrahedron satisfies our convention about the notion of a simplex in  $\mathbb{S}^n$  (see the beginning of Section 2.3).

1) is done exactly as in the hyperbolic case in Theorem 3. Here we can even have  $\pi \geq S_4 = S_3 = S_2 = S_1 > 0$ .

For 2) we have to handle the case  $S_3 = S_4 = \varphi = \pi/2$  only. From Construction 2 for any  $x \in [0, t] = [0, \pi/2]$  we have  $S \leq \pi/2$ , where equality can be attained for any  $x \in [0, t]$ . The case of equality is independent of  $x \in [0, t]$ : namely it is a regular spherical triangle with angles and sides  $\pi/2$ . Then the fact that the angle of the facets  $A_1(x, \varphi)A_2(x, \varphi)A_3(x, \varphi)$  and  $A_1(x, \varphi)A_2(x, \varphi)A_4(x, \varphi)$  is  $\varphi = \pi/2$  uniquely determines our simplex: it is a regular simplex in  $\mathbb{S}^3$  of edge  $\pi/2$  (thus we have also  $S_1 = S_2 = \pi/2$  — the vertices can be  $\mathbf{e}_1, \dots, \mathbf{e}_4$ ). This lies in some open half- $\mathbb{S}^3$ , therefore is among the simplices that we considered as simplices in  $\mathbb{S}^3$ .  $\square$

**6.3. Proof of Proposition 5.** For part (i), we start with  $n = 2$  dimensions. Here, one has a convex  $m$ -gon in a closed half- $\mathbb{S}^2$ , with sides  $S_1, \dots, S_m$ . In fact, it lies in an open half- $\mathbb{S}^2$ , has strictly convex angles, and is non-degenerate if  $S_m < S_1 + \dots + S_{m-1}$  and  $S_1 + \dots + S_m < 2\pi$ . For the degenerate cases, i.e., when  $S_m = S_1 + \dots + S_{m-1}$  or  $S_1 + \dots + S_m = 2\pi$ , we have a doubly counted segment or a great- $\mathbb{S}^1$ , respectively. If both equations hold, then we have also a digon, with one side subdivided to

$m - 1$  sides. If both inequalities are strict, we can copy the well-known proof in [34, pp. 53–54] — given there for the case of  $\mathbb{R}^2$ . Thus, we obtain the existence of such a convex  $m$ -gon. Actually, one gets such a convex  $m$ -gon that is inscribed in a circle of radius less than  $\pi/2$ .

Now we show how this construction can be lifted to higher dimensions. For  $\mathbb{S}^3$ , we embed the above  $m$ -gon in its equator, which is an  $\mathbb{S}^2$ . Each side of this polygon is then replaced by a facet that is the union of all meridians (whose lengths are  $\pi$ ) meeting that side. The vertices are replaced similarly by edges that are meridians meeting these vertices. Additionally, there are two new vertices at the North and South Poles. Then the ratio of the areas of the spherical digons and the lengths of the corresponding edges of our polygon is  $V_2(\mathbb{S}^2)/V_1(\mathbb{S}^1)$ , where  $V_i$  denotes  $i$ -volume. Moreover, the dihedral angles are the same as for the spherical  $m$ -gon in  $\mathbb{S}^2$ .

The inductive step is performed analogously for all  $n > 3$ . The other stated properties are obvious.

Part (ii) about simplices is proved by induction on  $n$ . For  $n = 2$ , we have a spherical triangle, and we have the same degenerate cases as in part (i) for  $m = n + 1 = 3$ . Let  $n \geq 3$ , and assume that the statement of the theorem holds for  $n - 1$ . With the factor  $\alpha := V_{n-2}(\mathbb{S}^{n-2})/V_{n-1}(\mathbb{S}^{n-1})$ , the numbers  $\alpha(S_1 + S_2), \alpha S_3, \dots, \alpha S_{n+1}$  satisfy the hypotheses of the proposition for  $n - 1$ . Arguing as in the second proof of Theorem 2 in Section 4.4 part 1, we establish that  $S_{\text{new}} := S_1 + S_2 \leq S_3 + \dots + S_{n+1}$ , since  $n + 1 \geq 4$ , and, for  $j \geq 3$ ,  $S_j \leq S_{\text{new}} + S_3 + \dots + S_{j-1} + S_{j+1} + \dots + S_{n+1}$ .

Therefore, we have on  $\mathbb{S}^{n-1}$  a polyhedral complex that is a combinatorial simplex with these facet areas. Again we consider  $\mathbb{S}^{n-1}$  as the equator of  $\mathbb{S}^n$ . We replace each facet and each lower-dimensional face of this polyhedral complex on  $\mathbb{S}^{n-1}$  by the union of all meridians (whose lengths are  $\pi$ ) meeting it. The resulting facets and also lower-dimensional faces are of one dimension higher than the original ones. Additionally, there are two new vertices at the North and South Poles. Thus, we have obtained a polyhedral complex on  $\mathbb{S}^n$  with facet areas  $S_1 + S_2, S_3, \dots, S_{n+1}$ .

This polyhedral complex has only two vertices at the two poles, and  $n$  edges joining them. Its  $n$  facets are obviously not simplices. The facet of area  $S_1 + S_2$  has  $n - 1$  edges. On each of these  $n - 1$  edges, we add an extra vertex at the same geographic latitude. Also we add an extra (convex) simplicial  $(n - 2)$ -face with these vertices, together with its faces of lower dimensions, that subdivides the facet of area  $S_1 + S_2$ . For a suitable choice of the latitude, the facet of area  $S_1 + S_2$  is subdivided into two new  $(n - 1)$ -dimensional simplicial facets of areas  $S_1$  and  $S_2$ . Omitting the facet of area  $S_1 + S_2$ , with all its faces of positive dimension, these two new facets with all their lower-dimensional faces are added as well. In each of the other facets (of areas  $S_3, \dots, S_{n+1}$ ), one  $(n - 2)$ -face has been subdivided into two new simplicial  $(n - 2)$ -faces. Thus, these other facets also become combinatorial  $(n - 1)$ -simplices, by induction with respect to  $n$ .

The other stated properties follow by the construction.  $\square$

**Added in proof**, 20. Oct. 2014. Observe that (i) of Theorem F' is preserved when passing to weak\* limits, so this will present no problem. However, (ii) of Theorem F' does not hold automatically, we have to prove it for  $K'$  (and for  $K''$ , but that follows from the considerations for  $K'$ ). For  $K$  we have (ii) of Theorem F', that can be rewritten as  $\int_{S^{n-1}} |\langle u, u_0 \rangle| d\mu_K > 0$ , for each  $u_0 \in S^{n-1}$ .

Since this integral is a continuous (actually Lipschitz) function of  $u_0$ , by compactness this integral even has a positive lower bound  $a$  independent of  $u_0$ . Now let  $\varepsilon > 0$  be sufficiently small. Let  $\{u_{01}, \dots, u_{0m}\} \subset S^{n-1}$  be an  $\varepsilon$ -net. Then for any  $u_{0k}$  we have  $\int_{S^{n-1}} |\langle u, u_{0k} \rangle| d\mu_K \geq a > 0$ . We may suppose that each  $K_{i_j}$  satisfies  $\int_{S^{n-1}} |\langle u, u_{0k} \rangle| d\mu_{K_{i_j}} > a/2 > 0$ . For any  $u_0 \in S^{n-1}$  we have  $\|u_0 - u_{0k}\| < \varepsilon$  for some  $1 \leq k \leq m$ . Then  $|\int_{S^{n-1}} |\langle u, u_0 \rangle| d\mu_{K_{i_j}} - \int_{S^{n-1}} |\langle u, u_{0k} \rangle| d\mu_{K_{i_j}}| \leq \int_{S^{n-1}} |\langle u, u_0 - u_{0k} \rangle| d\mu_{K_{i_j}} < \varepsilon \int_{S^{n-1}} 1 d\mu_{K_{i_j}}$ . We may also suppose that  $\int_{S^{n-1}} 1 d\mu_{K_{i_j}} < 2 \int_{S^{n-1}} 1 d\mu_K =: 2b$ . Then  $\int_{S^{n-1}} |\langle u, u_0 \rangle| d\mu_{K_{i_j}} > \int_{S^{n-1}} |\langle u, u_{0k} \rangle| d\mu_{K_{i_j}} - 2b\varepsilon > a/2 - 2b\varepsilon > 0$ . Hence passing to the limit  $K'$  we have  $\int_{S^{n-1}} |\langle u, u_0 \rangle| d\mu_{K'} \geq a/2 - 2b\varepsilon > 0$ . Thus the measure  $\mu_{K'}$  satisfies both (i) and (ii) of Theorem F', hence it is the surface area measure of a convex body  $K'$ .

**Acknowledgements.** The authors are indebted to R. Schneider for pointing out in his book [55] the proof of Theorem G, used in the second proof of our Theorem 2; to F. Morgan for pointing out the references [14] and [41]; and to G. Panina for pointing out the results about the orthocentric simplices [24, 20]. The second named author (E. M., Jr.) expresses his thanks to the Fields Institute, where part of this work was done during a Conference on Discrete Geometry, in September 2011. Research of the fifth author (G. R.) was initiated at the reunion conference of the special semester on discrete and computational geometry at the Bernoulli Center, EPFL Lausanne, February 27–March 2, 2012.

## REFERENCES

- [1] A. D. Aleksandrov, *Zur Theorie der gemischten Volumina von konvexen Körpern III, Die Erweiterung zweier Lehrsätze Minkowskis über die konvexen Polyeder auf die beliebigen konvexen Körper*, Mat. Sb. **3** (1938), 27–46, Russian, with German summary, Zbl.**18**.42402; English translation A. D. Alexandrov *Selected Works, Part I*, Selected Scientific papers, Edited by Yu. G. Reshetnyak and S. S. Kutateladze, transl. by P. S. V. Naidu, Classics of Soviet Math. **4**, Gordon and Breach, Amsterdam, 1996, MR**2000a**:01035.
- [2] A. D. Alexandrov, *Die innere Geometrie der konvexen Flächen*, Math. Lehrbücher und Monographien, II. Abteilung: Math. Monographien, **4**, Akademie Verlag, Berlin, 1955, MR**17**,74; English translation A. D. Alexandrov *Selected Works, Part II, Intrinsic Geometry of Convex Surfaces*, Edited by S. S. Kutateladze, transl. by S. Vakhrameyev, Chapman and Hall/CRC, Boca Raton, FL, 2006, MR**2006h**:01014.
- [3] A. D. Alexandrov, *Konvexe Polyeder*, Math. Lehrbücher und Monographien, II. Abteilung: Math. Monographien, **8**, Akademie Verlag, Berlin, 1958, MR**19**,1192; English translation: A. D. Alexandrov *Convex polyhedra*, transl. by N. S. Dairbekov, S. S. Kutateladze, A. B. Sossinsky, with comments and bibliography by V. A. Zalgaller and appendices by L. A. Shor, Yu. A. Volkov, Springer Monographs in Math., Springer, Berlin, 2005, MR**2005j**:52002.
- [4] V. A. Aleksandrov, *How to crumple a regular tetrahedral packet of milk, so that it could contain more* (Russian), Sorosovskij Obrazovatel'nyj Zhurnal **6** (2000), 121–127. Zbl. **947**.52012.
- [5] D. V. Alekseevskij, E. B. Vinberg, A. S. Solodovnikov, *Geometry of spaces of constant curvature*, In: Geometry II (Ed. E. B. Vinberg), Encyclopaedia Math. Sci. **29**, 1–138, Springer, Berlin, 1993, MR**95b**:53042.
- [6] F. C. Auluck, *The volume of a tetrahedron, the areas of the faces being given*, Proc. Indian Acad. Sci., Sect. A **7** (1938), 279–281, Zbl.**18**.37106.  
Online access: [http://www.ias.ac.in/j\\_archive/proca/7/4/279-281/viewpage.html](http://www.ias.ac.in/j_archive/proca/7/4/279-281/viewpage.html)
- [7] R. Baldus, *Nichteuklidische Geometrie, Hyperbolische Geometrie der Ebene*, Vierte Aufl., Bearb. und ergänzt von F. Löbell, Sammlung Göschen **970/970a**, de Gruyter, Berlin, 1964, MR**29**#3936.

- [8] D. D. Bleecker, *Volume increasing isometric deformations of convex polyhedra*, J. Diff. Geom. **43** (1996), 505–526, MR97g:52035.
- [9] C. W. Borchardt, *Über die Aufgabe des Maximums, welche der Bestimmung des Tetraeders von grösstem Volumen bei gegebenen Flächeninhalten der Seitenflächen für mehr als drei Dimensionen entspricht*, Math. Abh. Akad. Wiss. Berlin, 1866, 121–155 = C. W. Borchardts Gesammelte Werke, Verl. G. Leimer, Berlin, 1888, 201–232. Jahr. Fortschr. Math. **20**.15.01.
- [10] K. Böröczky, I. Bárány, E. Makai, Jr., J. Pach, *Maximal volume enclosed by plates and proof of the chessboard conjecture*, Discr. Math. **60** (1986), 101–120, MR87m:52016.
- [11] K. Böröczky, G. Kertész, E. Makai, Jr., *The minimum area of a simple polygon with given side lengths*, Periodica Mathematica Hungarica **39** (1–3) (1999), 33–49, MR2001e:51016.
- [12] K. Böröczky, Jr., *Finite packing and covering*, Cambridge Tracts in Math. **154**, Cambridge Univ. Press, Cambridge, 2004, MR2005g:52045.
- [13] P. Brass, W. Moser, J. Pach, *Research Problems in Discrete Geometry*, Springer, New York, NY, 2005, MR2006i:52001.
- [14] H. Brunn, *Über Ovale und Eiflächen*, Inauguraldissertation, München, 1887, Jahr. Fortschr. Math. **19**.0615.01.
- [15] K. Buchin and A. Schulz, *Inflating the cube by shrinking* (video), in: Proc. 23rd ACM Symp. Comput. Geom., Gyeongju, South Korea, June 6–8, 2007, pp. 125–126, ACM Press. Video: <http://www.computational-geometry.org/SoCG-videos/socg07video/>
- [16] Ju. D. Burago, V. A. Zalgaller, *Isometric piecewise-linear embeddings of two-dimensional manifolds with a polyhedral metric into  $\mathbb{R}^3$* , Algebra i Analiz **7** (1995) (3), 76–95 (in Russian); English translation in St. Petersburg Math. J. **7** (1996) (3), 369–385, MR96g:53091.
- [17] P. Buser, *Geometry and spectra of compact Riemann surfaces*, Progress in Mathematics **106**, Birkhäuser, Boston, Mass., 1992, MR93g:58149.
- [18] H. S. M. Coxeter, *Non-Euclidean Geometry. 6th ed.*, Spectrum Series, The Mathematical Association of America, Washington, DC, 1998, MR99c:51002.
- [19] N. Dunford, J. T. Schwartz, *Linear Operators, Vol. 1, General Theory*, Pure and Appl. Math. **7**, Interscience, New York-London, 1958, MR22#8302.
- [20] A. L. Edmonds, M. Hajja, H. Martini, *Orthocentric simplices and their centers*, Results Math. **47** (2005), 266–295, MR2006e:51021.
- [21] L. Fejes Tóth, *The isepiphany problem for  $n$ -hedra*, Amer. J. Math. **70** (1948), 174–180, MR9,460.
- [22] L. Fejes Tóth, *Reguläre Figuren*, Akadémiai Kiadó, Budapest, 1965, MR30#3408; in English *Regular Figures*, Pergamon Press, Macmillan, New York, 1964, MR29#2705.
- [23] W. Fenchel, B. Jessen, *Mengenfunktionen und konvexe Körper*, Danske Vid. Selsk., Mat.-Fys. Medd. **16** (1938), 1–31, Zbl.18.42401.
- [24] L. Gerber, *The orthocentric simplex as an extreme simplex*, Pacific J. Math. **56** (1975), 97–111, MR51#12717.
- [25] P. Gallagher, W. Ghang, D. Hu, Z. Martin, M. Miller, B. Perpetua, S. Waruhiu, *Surface-area-minimizing  $n$ -hedral tiles*, Rose-Hulman Undergrad. Math. J. **15** (2014), 209–236, MR3216229
- [26] M. Goldberg, *The isoperimetric problem for polyhedra*, Tôhoku Math. J. **40** (1935), 226–236, Zbl.10.41004.
- [27] P. Grizmann, J. M. Wills, D. Wrase, *A new isoperimetric inequality*, J. reine angew. Math. **379** (1987), 22–30, MR88h:52018.
- [28] U. Haagerup, H. J. Munkholm, *Simplices of maximal volume in hyperbolic  $n$ -space*, Acta Math. **147** (1981), 1–11, MR82j:53116.
- [29] W. Hurewicz, H. Wallman, *Dimension Theory*, Princeton Univ. Press, Princeton, N.J., 1941, MR3,312b.
- [30] K. S. K. Iyengar, *A note on Narasinga Rao’s problem relating to tetrahedra*, Proc. Indian Acad. Sci., Sect. A **7** (1938), 269–278, Zbl.18.37105.  
Online access: [http://www.ias.ac.in/j\\_archive/proca/7/4/269-278/viewpage.html](http://www.ias.ac.in/j_archive/proca/7/4/269-278/viewpage.html)
- [31] K. S. K. Iyengar, K. V. Iyengar, *On a problem relating to a tetrahedron*, Proc. Indian Acad. Sci., Sect. A **7** (1938), 305–311, Zbl.19.07506.  
Online access: [http://www.ias.ac.in/j\\_archive/proca/7/5/305-311/viewpage.html](http://www.ias.ac.in/j_archive/proca/7/5/305-311/viewpage.html)



- [32] I. M. Jaglom, W. G. Boltjanski, *Konvexe Figuren*, Hochschulbücher für Mathematik **24**, VEB Deutscher Verl. Wiss. Berlin, 1956, MR18,146f (MR14,197d); English translation I. M. Jaglom, V. G. Boltjanski, *Convex Figures*, transl. by P. J. Kelly and L. F. Walton, Rinehart and Winston, New York, 1960, MR23#A1283.
- [33] N. D. Kazarinoff, *Geometric inequalities*, Random House, New York-Toronto, 1961, New Mathematical Library **4**, MR24#A1.
- [34] D. A. Kryžanovskij, *Isoperimetry: maximum and minimum properties of geometrical figures*, 3-rd ed., with the redaction of I. M. Jaglom (in Russian), Fizmatgiz, Moskva, 1959.
- [35] J. L. Lagrange, *Solutions analytiques de quelques problèmes sur les pyramides triangulaires*, Nouv. Mém. Acad. Sci. Berlin, 1773, 149–176 = Œuvres complètes de Lagrange, Tome 3, Publié par J.-A. Serret, Gauthier-Villars, Paris, 1869, 659–692.
- [36] A. Lengyel, Zs. Gáspár, T. Tarnai, *The roundest polyhedra with symmetry constraint*, manuscript under preparation.
- [37] H. Liebmann, *Nichteuklidische Geometrie*, 3rd ed., de Gruyter, Berlin, 1923, Jahrb. Fortschr. Math. **49**.0390.01.
- [38] A. D. Milka, *Linear bendings of regular convex polyhedra* (Russian, with English, Russian, Ukrainian summaries), Mat. Fiz. Anal. Geom. **1** (1994), 116–130, MR98k:52056.
- [39] A. D. Milka, V. A. Gor'kavyj, *Volume increasing bendings of regular polyhedra* (Russian, with English summary), Zb. Pr. Inst. Mat. NAN Ukr. **6** (2009), 152–182, Zbl.1199,52007.
- [40] J. Milnor, *Hyperbolic geometry: the first 150 years*, Bull. Amer. Math. Soc., New Ser. **6** (1982), 9–24, MR82m:57005.
- [41] F. Morgan, *Riemannian Geometry. A Beginner's Guide*, 2nd ed., A. K. Peters, Wellesley, MA, 1998, MR98i:53001.
- [42] Yu. V. Nikonorova, *On an isoperimetric problem on the Euclidean plane*, Trudy Rubtsovsk. Ind. Inst. **9** (2001), 66–72 (Russian), Zbl.1006.51014.
- [43] S. P. Olovianishnikoff, *A generalization of Cauchy's theorem on convex polyhedra*, Rec. Math.=Mat. Sbornik N. S. **18 (60)** (3), 1946, 441–446 (Russian, with French summary) MR **8**,169.
- [44] I. Pak, *Inflating the cube without stretching*, Amer. Math. Monthly **115** (2008) (5), 443–445, MR2009c:52006.
- [45] O. Perron, *Nichteuklidische Elementargeometrie der Ebene*, Math. Leitfäden, Teubner, Stuttgart, 1962, MR25#2489.
- [46] A. V. Pogorelov, *Unique determination of convex surfaces (Russian)*, Trudy Mat. Inst. i. V. A. Steklova, **29**, Izdat. Akad. Nauk SSSR, Moskva-Leningrad, 1949, Zbl. **41**.508.
- [47] A. V. Pogorelov, *Geometric theory of stability of shells (Russian)*, Modern Problems of Math., Nauka, Moskva, 1966, MR34#3843.
- [48] A. V. Pogorelov, *Geometrical methods in the non-linear theory of elastic shells (Russian)*, Nauka, Moskva, 1967, MR36#3535.
- [49] A. V. Pogorelov, V. I. Babenko, *Geometrical methods in the theory of stability of thin shells (a survey)*, (in Russian, with English, Ukrainian summaries) Prikl. Mekh. **28** (1992) (1), 3–22; English translation: Internat. Appl. Mech. **28** (1992) (1), 1–17, MR93g:73089.
- [50] A. Narasinga Rao, *The Mathematics Student*, June 1937, **5** (2), 90.
- [51] I. S. Sabitov, *Algebraic methods for the solution of polyhedra*, Uspekhi Mat. Nauk **66** (2011) (3) (399), 3–66 (Russian); Transl. in: Russian Math. Surveys **66** (3) (2011), 445–505, MR2012k:52027.
- [52] L. A. Santaló, *Integral Geometry and Geometric Probability*, Encyclopedia of Mathematics and its Applications, **1**, Addison-Wesley, Reading, Mass. etc., 1976, MR55#6340.
- [53] E. Schmidt, *Die Brunn-Minkowskische Ungleichung und ihr Spiegelbild sowie die isoperimetrische Eigenschaft der Kugel in der euklidischen und nichteuklidischen Geometrie I*, Math. Nachr. **1** (1948), 81–157, MR10,471d.
- [54] E. Schmidt, *Die Brunn-Minkowskische Ungleichung und ihr Spiegelbild sowie die isoperimetrische Eigenschaft der Kugel in der euklidischen und nichteuklidischen Geometrie II*, Math. Nachr. **2** (1949), 171–244, MR11,534l.
- [55] R. Schneider, *Convex Bodies: the Brunn-Minkowski Theory*, Encyclopedia of Mathematics and its Applications, **44**, Cambridge University Press, Cambridge, 1993, MR94d:52007.

- [56] E. Steinitz, *Polyeder und Raumeinteilungen*, Enzyklopädie der Math. Wiss., mit Einschluss ihrer Anwendungen, 3. Band, Geometrie, Teil I, 2, Teubner, Leipzig, 1914–1931, Ch. 12, 1–139.
- [57] T. Tarnai, Z. Gáspár, A. Lengyel, *From spherical circle coverings to the roundest polyhedra*, *Phil. Mag.* **93** no-s 31–33 (2013), 3970–3982, DOI: <http://dx.doi.org/10.1080/14786435.2013.800652>
- [58] T. Tarnai, K. Hincz, A. Lengyel, *Volume increasing inextensional deformation of a cube*, Proc. Internat. Assoc. for Shell and Spatial Structures (IASS) Symp. 2013, “Beyond the limits of man”, Sept. 23–27, Wrocław Univ. of Techn., Poland, (eds. J. B. Odrębski and R. Tarczewski), CD-ROM, Paper ID 1253, 6pp., 2013.
- [59] T. Tarnai, A. Lengyel, K. Hincz, *Volume increasing isometric deformations of a cube*, manuscript under preparation.
- [60] A. Tarski, *A decision method for elementary algebra and geometry*, 2nd ed., Univ. Calif. Press, Berkeley, Los Angeles, Calif., 1951, MR13,423 (MR10,499).
- [61] K. Venkatachaliengar, *On a problem of the tetrahedron*, Proc. Indian Acad. Sci., Sect. A **7** (1938), 257–260, Zbl.18.37107.  
Online access: [http://www.ias.ac.in/j\\_archive/proca/7/4/257-260/viewpage.html](http://www.ias.ac.in/j_archive/proca/7/4/257-260/viewpage.html)
- [62] G. Wulff, *Zur Frage der Geschwindigkeit des Wachstums und der Auflösung der Kristallflächen*, *Zeitschr. für Kristallographie und Mineralogie* **34** (1901), 449–530.

N. V. ABROSIMOV, SOBOLEV INSTITUTE OF MATHEMATICS, NOVOSIBIRSK, ACAD. KOPTYUG AV. 4, 630090, RUSSIA;  
 NOVOSIBIRSK STATE UNIVERSITY, NOVOSIBIRSK, PIROGOV ST. 2, 630090, RUSSIA  
*E-mail address:* [abrosimov@math.nsc.ru](mailto:abrosimov@math.nsc.ru)

E. MAKAI, JR., A. RÉNYI INSTITUTE OF MATHEMATICS, HUNGARIAN ACADEMY OF SCIENCES, H-1364 BUDAPEST, PF. 127, HUNGARY  
*E-mail address:* [makai.endre@renyi.mta.hu](mailto:makai.endre@renyi.mta.hu), <http://www.renyi.mta.hu/~makai>

A. D. MEDNYKH, SOBOLEV INSTITUTE OF MATHEMATICS, NOVOSIBIRSK, ACAD. KOPTYUG AV. 4, 630090, RUSSIA;  
 NOVOSIBIRSK STATE UNIVERSITY, NOVOSIBIRSK, PIROGOV ST. 2, 630090, RUSSIA  
*E-mail address:* [mednykh@math.nsc.ru](mailto:mednykh@math.nsc.ru)

YU. G. NIKONOROV, SOUTH MATHEMATICAL INSTITUTE OF THE V. S. C. OF THE RUSSIAN ACADEMY OF SCIENCES, VLADIKAVKAZ, MARKUS STR. 22, 362027, RUSSIA  
*E-mail address:* [nikonorov2006@mail.ru](mailto:nikonorov2006@mail.ru)

G. ROTE, INSTITUT FÜR INFORMATIK, FREIE UNIVERSITÄT BERLIN, TAKUSTR. 9, 14159 BERLIN, GERMANY  
*E-mail address:* [rote@inf.fu-berlin.de](mailto:rote@inf.fu-berlin.de)