

The history of degenerate (bipartite) extremal graph problems

Zoltán Füredi
and Miklós Simonovits

May 15, 2013 / Final version

Contents

1	Introduction	1
1.1	Some central theorems of the field	2
1.2	The structure of this paper	3
1.3	Extremal problems	5
1.4	Other types of extremal graph problems	8
1.5	Historical remarks	9
2	The general theory, classification	10
2.1	The importance of the Degenerate Case	11
2.2	The asymmetric case of Excluded Bipartite graphs	13
2.3	Reductions: Host graphs	14
2.4	Excluding complete bipartite graphs	15
2.5	Probabilistic lower bound	17
2.6	Classification of extremal problems	20
2.7	General conjectures on bipartite graphs	22
3	Excluding complete bipartite graphs	22
3.1	Bipartite C_4 -free graphs and the Zarankiewicz problem	22
3.2	Finite Geometries and the C_4 -free graphs	25
3.3	Excluding C_4 : Exact results	27
3.4	Excluding $K(2, t + 1)$, $t > 1$	28
3.5	Excluding $K(3, 3)$, and improving the upper bound	31

3.6	Further applications of Algebraic Methods	33
3.7	The coefficient in the Kővári-T. Sós-Turán bound	36
3.8	Excluding large complete subgraphs	36
4	Excluding Cycles : C_{2k}	37
4.1	Girth and Turán numbers, upper bounds	38
4.2	Excluding a single C_{2k} , upper bounds	39
4.3	Eliminating short cycles, a promising attempt	41
4.4	A lower bound for C_6 : The Benson Construction	42
4.5	Girth 12 graphs by Benson and by Wenger	44
4.6	Short cycles, C_6 and C_{10}	45
4.7	Bipartite hosts with extreme sides	47
4.8	The effect of odd cycles	47
4.9	Large girth: Ramanujan graphs	49
4.10	The girth problem: the Lazebnik-Ustimenko approach	52
4.11	Cycle length distribution	53
5	Paths and long cycles	54
5.1	Excluding long cycles	54
5.2	Excluding P_k	55
5.3	Proof ideas	57
5.4	Generalizations	58
6	Excluding trees	58
6.1	Erdős-Sós conjecture	58
6.2	Sketch of the proof of Theorem 6.2	59
6.3	Komlós-Sós conjecture on median degree	61
7	More complex excluded subgraphs	61
7.1	The Erdős-Simonovits Reduction and the Cube theorem	62
7.2	Theta graphs and the Faudree-Simonovits reduction	65
7.3	A universal graph and dependent random choice	66
8	Eigenvalues and extremal problems	68
9	Excluding topological subdivisions	69
9.1	Large topological subgraphs	69
9.2	Turán numbers of subdivided graphs	71

10 Hypergraph Extremal Problems	71
10.1 Positive Density problems	71
10.2 Degenerate hypergraph problems	72
11 Supersaturated graphs	73
11.1 Erdős-Simonovits-Sidorenko conjecture	74
12 Ordered structures	76
12.1 Directed graphs, ordered graphs	76
12.2 Erdős-Moser conjecture on unit distances	76
12.3 Ordered submatrices	77
12.4 Ordered matrices and the Stanley-Wilf conjecture on subpermutations	78
13 Applications in Geometry	79
13.1 Applicability of the Kővári-T. Sós-Turán bound	79
13.2 Unit Distances	80
13.3 Cells in line arrangements	81
14 Further connections and problems	81
14.1 Connections of hypergraphs and critical graphs	81
14.2 A multiplicative Sidon problem and C_{2k} -free graphs	84
14.3 Cycle-free subgraphs of the d -dimensional hypercube	85
14.4 Two problems of Erdős	85
15 Acknowledgements	86

1. Introduction

This survey describes the theory of *Degenerate Extremal Graph Problems*, the main results of the field, and its connection to the surrounding areas.

Extremal graph problems we consider here are often called Turán type extremal problems, because Turán’s theorem and his questions were the most important roots of this huge area [242], [243].

Generally, we have a *Universe* of graphs, \mathbb{U} , where this universe may be the family of ordinary graphs, or digraphs, or hypergraphs, or ordered graphs, or bipartite graphs, etc and a property \mathcal{P} , saying, e.g., that $G \in \mathbb{U}$ does not contain some subgraphs $L \in \mathcal{L}$, or that it is Hamiltonian, or it is at most 3-chromatic, and we have some parameters on \mathbb{U} , say $v(G)$ and $e(G)$, the number of vertices and edges. Our aim is to maximize the second parameter under the condition that G has property \mathcal{P} and its first parameter is given.

We call such a problem *Turán type extremal problem* if we are given a family \mathcal{L} of graphs from our universe, G_n is a graph of n vertices, $e(G_n)$ denotes the number of edges of G_n and we try to maximize $e(G_n)$ under the condition that G_n contains no $L \in \mathcal{L}$, where “contains” means “not necessarily induced subgraph”. (Here graph may equally mean digraph, or multigraph, or hypergraph).

The maximum will be denoted by $\mathbf{ex}(n, \mathcal{L})$ and the graphs attaining this maximum without containing subgraphs from \mathcal{L} are called *extremal graphs*. The family of extremal graphs is denoted by $\mathbf{EX}(n, \mathcal{L})$ and $\mathbf{ex}(n, \mathcal{L})$ is called the *Turán number* of the family \mathcal{L} .

Speaking of $\mathbf{ex}(n, \mathcal{L})$ we shall always assume that $n \geq |V(L)|$, otherwise the problem is trivial.

Definition 1.1. If the Universe \mathbb{U} is the family of r -uniform hypergraphs¹, then we shall call the problem *degenerate* if the maximum,

$$\mathbf{ex}(n, \mathcal{L}) = o(n^r).$$

Otherwise we shall call it *non-degenerate*

¹ $r = 2$ included, moreover, mostly we think of $r = 2$.

Below we shall mention several open problems. Yet to get more problems, we refer the reader to the

Erdős homepage: www.renyi.hu/~p_erdos

where the papers of Erdős can be found [59]. Also, many open problems can be found in Chung-Graham [47].

1.1. Some central theorems of the field

We start with some typical theorems of the field and two conjectures. The aim of this “fast introduction” is to give a feeling for what are the crucial types of results here.

Theorem 1.2 (Kővári–T. Sós–Turán, [164]). *Let $K_{a,b}$ denote the complete bipartite graph with a and b vertices in its color-classes. Then*

$$\mathbf{ex}(n, K_{a,b}) \leq \frac{1}{2} \sqrt[a]{b-1} \cdot n^{2-(1/a)} + O(n).$$

We use this theorem with $a \leq b$, since that way we get a better estimate.

Theorem 1.3 (Kollár-Alon-Rónyai-Szabó [159], [11]). *If $b > (a-1)!$, then*

$$\mathbf{ex}(n, K_{a,b}) > c_a n^{2-(1/a)}.$$

Theorem 1.4 (Erdős, Bondy and Simonovits [32]).

$$\mathbf{ex}(n, C_{2k}) \leq 100kn^{1+(1/k)}.$$

Theorem 1.5 (Erdős–Simonovits, Cube Theorem [90]). *Let Q_8 denote the cube graph defined by the vertices and edges of a 3-dimensional cube. Then*

$$\mathbf{ex}(n, Q_8) = O(n^{8/5}).$$

Conjecture 1.6 (Erdős and Simonovits, Rational exponents). *For any finite family \mathcal{L} of graphs, if there is a bipartite $L \in \mathcal{L}$, then there exist a rational $\alpha \in [0, 1)$ and a $c > 0$ such that*

$$\frac{\mathbf{ex}(n, \mathcal{L})}{n^{1+\alpha}} \rightarrow c.$$

Theorem 1.7 (Füredi [111], [104]). *If $q \neq 1, 7, 9, 11, 13$, and $n = q^2 + q + 1$, then*

$$\mathbf{ex}(n, C_4) \leq \frac{1}{2}q(q+1)^2.$$

Moreover, if q is a power of a prime, then

$$\mathbf{ex}(n, C_4) = \frac{1}{2}q(q+1)^2.$$

Conjecture 1.8 (Erdős).²

$$\mathbf{ex}(n, \{C_3, C_4\}) = \frac{1}{2\sqrt{2}}n^{3/2} + o(n^{3/2}).$$

We close this part with a famous result of Ruzsa and Szemerédi:

Theorem 1.9 (Solution of the (6,3) problem, [210]). *If $\mathcal{H}_n^{(3)}$ is a 3-uniform hypergraph not containing 6 vertices determining (at least) 3 hyperedges, then this hypergraph has $o(n^2)$ hyperedges.*

The above theorems will be discussed in more details below.

1.2. The structure of this paper

The area is fairly involved. Figure 1 shows a complicated – but not complete – map of the interactions of some subfields of the field discussed here. We start with describing the Extremal problems in general, then move to the Degenerate problems, also describing why they are important. Among the most important degenerate extremal problems we mention are the extremal problem of $K_{a,b}$, and also C_{2k} , where – to classify the extremal problems – we shall need the Random Graph Method to get a lower bound in the problem of C_{2k} . These results are enough to give a good *classification* of degenerate extremal graph problems.

1. The two lowest boxes of Figure 1 show that this whole area has a strong connection to Geometry and Number Theory. This will be explained in Sections 13, 1.5, and 14.2.
2. The origins of this field are
 - (a) an early, singular result of Mantel,

²This conjecture is mentioned in [70] but it is definitely older, see e.g. Brown, [37].




Figure 1: Area Map

- (b) the multiplicative Sidon Problem (see Section 1.5)
- (c) Turán’s theorem and his systematic approach to the field.

So the origins come – in some sense – from Number Theory, are strongly connected to Finite Geometry, and in this way also to ordinary geometry (Turán’s theorem comes from the Erdős-Szekeres version of Ramsey Theorem, which they invented to solve the Esther Klein problem from Geometry.)

3. To understand the field we start with a very short description of the general theory, and then – skipping most of the hypergraph theory – we move to the main area of this paper: to the questions where we consider ordinary extremal graphs, and exclude some bipartite L : therefore, by Theorem 1.2, we have $\mathbf{ex}(n, \mathcal{L}) = O(n^{2-c})$.
4. One important phenomenon is that many extremal graph problems can be “reduced” to some *degenerate* extremal graph problems that we also call sometimes *bipartite extremal problems*.
5. The upper bounds in the simpler cases are obtained by some double counting, Jensen type inequalities, or applying some supersaturated graph theorems.³

³Lovász and Szegedy had a beautiful conjecture, which we formulate here only in a restricted form: Any (valid) extremal theorem can be proven by applying the Cauchy-Schwarz inequality finitely many times. This conjecture was killed in this strong form – by Hatami and Norine [142] – but proven in a weaker, “approximation-form”.

6. There are also much more complicated cases, where the above simple approach is not enough, we need some finer arguments. Perhaps the first such case was treated by Füredi: Section 7.3 and [107]. Also such an approach is the application of the general Dependent Random Choice Method, (see the survey of Fox and Sudakov [101]).
7. The lower bounds are sometimes provided by random graphs (see Section 2.5) but these are often too weak. So we often use some finite geometric constructions, (see Section 3.2) or their generalizations – coming from commutative algebras (see Sections 3.6, 4.9, 8), etc., and they occasionally provide matching upper and lower bounds. Again, there is an important general method with many important results, which we shall call the Lazebnik-Ustimenko method but will treat only very superficially in Section 3.6.

1.3. Extremal problems

We shall almost entirely restrict ourselves to Turán type extremal problems for ordinary simple graphs, i.e. loops, multiple edges are excluded.

To show the relation of the areas described here, we start with a list of some related areas.

1. Ramsey Theory
 - Problems not connected to density problems; in some sense these are the real Ramsey Problems
 - Problems connected to density problems, i.e. cases, where we do not really use the Ramsey Condition, only that some color class is large.
2. Ordinary extremal graph theory
 - Excluding bipartite graphs (degenerate problems)
 - Excluding topological subgraphs (very degenerate extremal problems)
 - Matrix problems, ordered and not ordered;
 - Non-degenerate case, and its relation to degenerate problems
3. Theory of extremal digraph problems
4. Ramsey-Turán Problems
5. Connection to Random Graphs

6. Hypergraph extremal problems
7. Connection of Number Theoretical problems to Extremal Graph Theory
8. Continuous problems
9. Applications

There are several surveys on these fields, see e.g., T. Sós [232], Füredi [108], [110], Simonovits [224], [228], [227], [222], Simonovits and Sós [230], [155]. Perhaps the first survey on this topic was Vera Sós' paper [232], discussing connections between extremal graph problems, finite geometries, block designs, etc. and, perhaps, the nearest to this survey is [225], Bollobás, [28] Sidorenko [217], [101], and also some books, e.g., Bollobás [26]. Of course, a lot of information is hidden in the papers of Erdős, among others, in [70], [73], [76].

So, here we shall concentrate on Case 2, but to position this area we shall start with some related fields, among others, with the general asymptotic in Case 2.

Problem 1.10 (General Host-graphs). In a more general setting we have a sequence (H_n) of “host” graphs and the question is, how many edges can a subgraph $G_n \subset H_n$ have under the condition that it does not contain any forbidden subgraph $L \in \mathcal{L}$. The maximum will be denoted by $\mathbf{ex}(H_n, \mathcal{L})$.

For $H_n = K_n$ we get back the ordinary extremal graph problems. There are several further important subcases of this question:

- (a) when $H_n = K_{a,b}$ for $a \approx n/2$;
- (b) when the host-graph is the k -dimensional cube, $n = 2^k$; see Section 14.3.
- (c) when H_n is a random graph on n vertices, see e.g. [209].

Notation. Given some graphs $G_n, T_{n,p}, T_k, H_\nu, \dots$ the (first) subscript will almost always denote the number of vertices.⁴ So K_p is the complete graph on p vertices, P_k the path on k vertices, C_k is the cycle of k vertices, while $\mathcal{C}_{\geq k}$ will denote the family of cycles of length at least k . $\delta(x)$ denotes the degree of the vertex x .

The complete bipartite graph $K_{a,b}$ with a vertices in its first class and b in its 2nd class will be crucial in this paper. Often, we shall denote it by

⁴One important exception is the complete bipartite graph $K(a,b) = K_{a,b}$, see below. Another exception is, when we list some excluded subgraphs, like L_1, \dots, L_ν .

$K(a, b)$, and its p -partite generalization by $K_p(a_1, \dots, a_p)$. If $\sum a_i = n$ and $|n_i - n_j| \leq 1$, then $K_p(a_1, \dots, a_p)$ is the Turán graph $T_{n,p}$ on n vertices and p classes.

Given two graphs U and W , their product graph is the one obtained from vertex-disjoint copies of these two graphs by joining each vertex of U to each vertex of W . This will be denoted by $U \otimes W$.⁵

Given a graph H , $v(H)$ is its number of vertices, $e(H)$ its number of edges and $\chi(H)$ its chromatic number, $d_{\min}(G)$ and $d_{\max}(G)$ denote the minimum and maximum degrees of G , respectively.

We shall write $f(x) \approx g(x)$ if $f(x)/g(x) \rightarrow 1$. Occasionally $[n]$ denote the set of first n integers, $[n] := \{1, 2, \dots, n\}$.

The Overlap. Some twenty years ago Simonovits wrote a survey [227] on the influence of Paul Erdős in the areas described above, Many-many features of these areas changed drastically since that. Jarik Nešetřil and Ron Graham were the editors of that survey-volume, and now they decided to republish it. Fortunately, the authors had the option to slightly rewrite their original papers. Simonovits has rewritten his original paper [228], basically keeping everything he could but *indicating* many new developments, and adding remarks and many new references to it.

To make *this* paper readable and self-contained, we shall touch on some basic areas also treated there, or in other survey papers of ours, Here, however, we shall explain many-many results and phenomena just mentioned in other survey papers.

Remark 1.11. There is also a third new survey to be mentioned here: Simonovits gave a lecture at the conference on Turán's 100th anniversary, in 2011. His lecture tried to cover the whole influence of Paul Turán in Discrete Mathematics. In the volume of this conference Simonovits wrote a survey [229] covering his lecture, except that

- the area called Statistical Group Theory is discussed in a survey of Pálffy and Szalay [200] and
- some parts of the applications of Extremal Graph Theory, primarily in Probability Theory are covered by Katona [153], in the same volume.

⁵This product is often called also the joint of the two graphs.

1.4. Other types of extremal graph problems

Above we still tried to maximize the number of edges, hyperedges, etc. More generally, instead of maximizing $e(G_n)$, we may maximize something else:

1. *Min-degree problems* (or Dirac type problems): How large min-degree can G_n have without containing subgraphs from \mathcal{L} .
2. *Median problems* which will be called here LoebL-Komlós-Sós type problems: Given a graph G_n , which m and d ensure that if G_n has at least m vertices of degree $\geq d$, then G_n contains some $L \in \mathcal{L}$.
3. *Eigenvalue-extremal problems*⁶: maximize the maximum eigenvalue $\lambda(G_n)$ under the condition that G_n does not contain any $L \in \mathcal{L}$. (These are sharper forms of some extremal results, since the maximum eigenvalue

$$\lambda(G_n) \geq \frac{2e(G_n)}{n},$$

see Section 8.)

4. *Subgraph count inequalities*, which assert that if G_n contains many copies of some subgraphs L_1, \dots, L_λ , then we have at least one (or maybe “many”) subgraphs L .
5. *Diameter-extremal problems*. Here we mention just a subcase: if

$$\text{diam}(G_n) \leq d \quad \text{and} \quad d_{\max}(G_n) < M,$$

at least how many edges must G_n have. The Erdős-Rényi paper [85] is of importance here and also some related papers, like Füredi [105], [109].

6. *Combined extremal problems*: There are many-many further types of extremal problems. Here we mention, as an example, the results of Balister, Bollobás, Riordan and Schelp [17], where an odd cycle is excluded, and at the same time an upper bound is fixed on the degrees and the number of edges are to be maximized.

The approach 4 is very popular in the theory of Graph Limits [177]. We mention a breakthrough in this area in connection with Erdős’ combinatorial problems, of type 4. A famous conjecture of Erdős was

⁶As usual, given a graph G_n , an $n \times n$ matrix is associated to it, having n eigenvalues. The largest and the second largest is what we are mostly interested in.

Conjecture 1.12 (Erdős [74]). *A K_3 -free G_n contains at most $(\frac{n}{5})^5$ copies of C_5 's.*

The motivation of this is that the blownup⁷ C_5 , i.e. $C_5[n/5]$ has no triangles and has $(\frac{n}{5})^5$ copies of C_5 . Erdős conjectured that no triangle-free G_n can have more C_5 's than this. The first “approximation” was due to Ervin Győri:

Theorem 1.13 (Győri [133]). *A K_3 -free G_n contains at most $1.03(\frac{n}{5})^5$ C_5 's.*

Next Füredi improved the constant to 1.001 (unpublished) [114], and finally independently Grzesik [124], and Hatami, Hladký, Král, Norine, and Razborov [141] proved the conjecture.

1.5. Historical remarks

Erdős in 1938 [60] considered the following “multiplicative Sidon Problem”⁸.

Problem 1.14. How many integers, $a_1, \dots, a_m \in [1, n]$ can we find so that $a_i a_j = a_k a_\ell$ does not hold for any i, j, k, ℓ , unless $\{i, j\} = \{k, \ell\}$.

To get an upper bound in Problem 1.14, Erdős proved

Theorem 1.15. *Let $G[n, n]$ be a bipartite graph with n vertices in both classes. If it does not contain C_4 , then $e(G[n, n]) < 3n\sqrt{n}$,*

Much later this problem was asked in a more general setting: find an upper bound on $e(G[n, n])$ if $K_{a,b} \not\subset G[n, n]$. Zarankiewicz [254] posed the following question:

Problem 1.16 (Zarankiewicz problem). Determine the largest integer $Z(m, n, a, b)$ for which there is an $m \times n$ 0-1 matrix containing $Z(m, n, a, b)$ 1's without an $a \times b$ submatrix consisting entirely of 1's.

⁷Given a graph H , its blownup version $H[t]$ is defined as follows: we replace each vertex x of H by t independent new vertices and we join two new vertices coming from distinct vertices x, y iff xy was an edge of H .

⁸For a longer description of the number theoretical parts see [228]. Erdős also refers to his “blindness” overlooking the general problem in [70].

Hartman, Mycielski and Ryll-Nardzewski [139] gave upper and lower bounds for the case $a = b = 2$, weaker than the Erdős-Klein⁹ result, and Kővári, T. Sós and Turán (see Theorem 1.2) provided a more general upper bound. We shall discuss these problems and results in details in Sections 2.4 and 3.2.

While exact values of $Z(m, n, a, b)$ are known for infinitely many parameter values, mostly only asymptotic bounds are known in the general case. Even $Z(m, n, 2, 2)$ is not sufficiently well known.

2. The general theory, classification

In many ordinary extremal problems the minimum chromatic number plays a decisive role. The **subchromatic number** $p(\mathcal{L})$ of \mathcal{L} is defined by

$$p(\mathcal{L}) = \min\{\chi(L) : L \in \mathcal{L}\} - 1. \quad (2.1)$$

Recall that the Turán graph $T_{n,p}$ is the largest graph on n vertices and p classes.

Claim 2.1.

$$\mathbf{ex}(n, \mathcal{L}) \geq e(T_{n,p}) = \left(1 - \frac{1}{p}\right) \binom{n}{2} + o(n^2). \quad (2.2)$$

Indeed, $T_{n,p}$ does not contain any $L \in \mathcal{L}$. An easy consequence of the Erdős-Stone theorem [95] provides the asymptotic value of $\mathbf{ex}(n, \mathcal{L})$, at least if $p(\mathcal{L}) > 1$.

Theorem 2.2 (Erdős-Simonovits [89]). *If \mathcal{L} is a family of graphs with subchromatic number $p > 0$, then*

$$\mathbf{ex}(n, \mathcal{L}) = \left(1 - \frac{1}{p}\right) \binom{n}{2} + o(n^2).$$

This means that $\mathbf{ex}(n, \mathcal{L})$ depends only very loosely on \mathcal{L} ; up to an error term of order $o(n^2)$; it is already determined by $p(\mathcal{L})$.¹⁰ The question is whether the structure of the extremal graphs is also almost determined by

⁹In [70] Erdős (again) attributes the finite geometric construction to Eszter (Esther) Klein.


¹⁰Better error terms are proved e.g. in [218], however, this will not be discussed here.

$p(\mathcal{L})$, and (therefore) it must be very similar to that of $T_{n,p}$ ¹¹. The answer is **YES**. This is expressed by the following results of Erdős and Simonovits [66], [67], [218]:

Theorem 2.3 (The Asymptotic Structure Theorem). *Let \mathcal{L} be a family of forbidden graphs with subchromatic number p . If $S_n \in \mathbf{EX}(n, \mathcal{L})$, (i.e., S_n is extremal for \mathcal{L}), then it can be obtained from $T_{n,p}$ by deleting and adding $o(n^2)$ edges. Furthermore, if \mathcal{L} is finite, then the minimum degree*

$$d_{\min}(S_n) = \left(1 - \frac{1}{p}\right)n + o(n).$$

Further, the almost-extremal graphs are similar to $T_{n,p}$.



Theorem 2.4 (The First Stability Theorem). *Let \mathcal{L} be a family of forbidden graphs with subchromatic number p . For every $\varepsilon > 0$, there exist a $\delta > 0$ and an n_ε such that, if G_n contains no $L \in \mathcal{L}$, and if, for $n > n_\varepsilon$,*

$$e(G_n) > \mathbf{ex}(n, \mathcal{L}) - \delta n^2, \quad (2.3)$$

then G_n can be obtained from $T_{n,p}$ by changing¹² at most εn^2 edges.

Remark 2.5. For ordinary graphs ($r = 2$) we often call the degenerate extremal graph problems *bipartite extremal problems*. This is the case when \mathcal{L} contains some bipartite graphs. There is a slight problem here: we shall also consider the case when not only some $L \in \mathcal{L}$ is bipartite but $\chi(G_n) = 2$ is as well.

2.1. The importance of the Degenerate Case

There are several results showing that if we know sufficiently well the extremal graphs for the degenerate cases, then we can also reduce the non-degenerate cases to these problems.

¹¹Actually, this was the original question; Theorem 2.2 was a partial answer to it.

¹²deleting and adding

Exact Turán numbers, product conjecture

We start with an illustration. Let $O_6 = K(2, 2, 2)$ be the octahedron graph. Erdős and Simonovits proved that

Theorem 2.6 (Octahedron Theorem [91]). *If S_n is an extremal graph for the octahedron O_6 for n sufficiently large, then there exist extremal graphs G_1 and G_2 for the circuit C_4 and the path P_3 such that $S_n = G_1 \otimes G_2$ and $|V(G_i)| = \frac{1}{2}n + o(n)$, $i = 1, 2$.*

If G_1 does not contain C_4 and G_2 does not contain P_3 , then $G_1 \otimes G_2$ does not contain O_6 . Thus, if we replace G_1 by any H_1 in $\mathbf{EX}(v(G_1), C_4)$ and G_2 by any H_2 in $\mathbf{EX}(v(G_2), P_3)$, then $H_1 \otimes H_2$ is also extremal for O_6 .

More generally,

Theorem 2.7 (Erdős–Simonovits [91]). *Let L be a complete $(p+1)$ -partite graph, $L := K(a, b, r_3, r_4, \dots, r_{p+1})$, where $r_{p+1} \geq r_p \geq \dots \geq r_3 \geq b \geq a$ and $a = 2, 3$. There exists an $n_0 = n_0(a, b, \dots, r_{p+1})$ such that if $n > n_0$ and $S_n \in \mathbf{EX}(n, L)$, then $S_n = U_1 \otimes U_2 \otimes \dots \otimes U_p$, where*

1. $v(U_i) = n/p + o(n)$, for $i = 1, \dots, p$.
2. U_1 is extremal for $K_{a,b}$
3. $U_2, U_3, \dots, U_p \in \mathbf{EX}(n, K(1, r_3))$.

It follows that this theorem is indeed a reduction theorem.

Conjecture 2.8 (The Product Conjecture, Simonovits). *Assume that $p(\mathcal{L}) = \min_{L \in \mathcal{L}} \chi(L) - 1 > 1$. If for some constants $c > 0$ and $\varepsilon \in (0, 1)$*

$$\mathbf{ex}(n, \mathcal{L}) > e(T_{n,p}) + cn^{1+\varepsilon}, \quad (2.4)$$

then there exist p forbidden families \mathcal{M}_i , with

$$p(\mathcal{M}_i) = 1 \quad \text{and} \quad \max_{M \in \mathcal{M}_i} v(M) \leq \max_{L \in \mathcal{L}} v(L),$$

such that for any $S_n \in \mathbf{EX}(n, \mathcal{L})$, $S_n = G_1 \otimes \dots \otimes G_p$, where G_i are extremal for \mathcal{M}_i .

This means that the extremal graphs S_n are products of extremal graphs for some degenerate extremal problems (for \mathcal{M}_i), and therefore we may reduce the general case to degenerate extremal problems.

Remarks 2.9. (a) If we allow infinite families \mathcal{L} , then one can easily find counterexamples to this conjecture.

(b) If we allow linear error-terms, i.e. do not assume (2.4), then one can also find counterexamples, using a general theorem of Simonovits [221]; however, this is not trivial at all, see [223]

(c) A weakening of the above conjecture would be the following: for arbitrary large n , in Conjecture 2.8 there are several extremal graphs, and for each $n > n_{\mathcal{L}}$, some of them are of product form, (but maybe not all of them) and the families \mathcal{M}_i also may depend on n a little.

Further sources to read: Griggs, Simonovits, and Thomas [127], Simonovits, [226].

2.2. The asymmetric case of Excluded Bipartite graphs

The degenerate extremal graph problems have three different forms:

Problem 2.10 (Three versions). (a) Ordinary extremal graph problems, where some bipartite or non-bipartite sample graphs are excluded, and we try to maximize $e(G_n)$ under this conditions.

(b) The *bipartite case*, where the host graph is $K(m, n)$ and we maximize $e(G_{n+m})$ under the conditions that $G_{n+m} \subseteq K(m, n)$ and G_{n+m} contains no $L \in \mathcal{L}$. (Here we may assume that all $L \in \mathcal{L}$ are bipartite.) In this case we often use the notation $\mathbf{ex}(n, m; \mathcal{L})$. If $m \leq n$ but $m > cn$ for some constant $c > 0$, then the answer to this problems and to the problem of $\mathbf{ex}(n, \mathcal{L})$ are the same, up to a constant. If, however, we assume that n is much larger than m , then some surprising new phenomena occur, see Section 14.2.

(c) The *asymmetric case*. Color the vertices of the sample graphs L in RED-BLUE and exclude only those $G_n \subseteq K(m, n)$ where the RED vertices of some $L \in \mathcal{L}$ are in the FIRST class of $K(m, n)$: maximize $e(G_{n+m})$ over the remaining graphs $G_{n+m} \subseteq K(m, n)$.

Denote the maximum number of edges in this third case by $\mathbf{ex}^*(m, n, \mathcal{L})$.

Remark 2.11. We have seen Zarankiewicz' problem (i.e. Problem 1.16). That corresponds to an asymmetric graph problem, (c). If we exclude in an $m \times n$ matrix both a $a \times b$ and an $b \times a$ submatrices of 1's, that will correspond to a bipartite graph problem, (b).

Conjecture 2.12 (Erdős, Simonovits [225]).

$$\mathbf{ex}^*(n, n, \mathcal{L}) = O(\mathbf{ex}(n, \mathcal{L})).$$

The simplest case when we cannot prove this is $L = K(4, 5)$.

Remark 2.13 (Matrix problems). Case (c) has also a popular matrix form where we consider 0-1 matrices and consider an $n_1 \times n_2$ matrix not containing a submatrix A . The question is: how many 1-s can be in such a matrix. This problem has (at least) two forms: the unordered and ordered one. We return to the Ordered Case in Subsection 12.4.

2.3. Reductions: Host graphs

The following simple but important observation shows that there is not much difference between considering any graph as a “host” graph or only bipartite graphs.

Lemma 2.14 (Erdős’ bipartite subgraph lemma). *Every graph G_n contains a bipartite subgraph H_n with $e(H_n) \geq \frac{1}{2}e(G_n)$.*

This lemma shows that there is not much difference between considering K_{2n} or $K_{n,n}$ as a host graph.

Corollary 2.15. *If $\mathbf{ex}_B(n, \mathcal{L})$ denotes the maximum number of edges in an \mathcal{L} -free bipartite graph, then*

$$\mathbf{ex}_B(n, \mathcal{L}) \leq \mathbf{ex}(n, \mathcal{L}) \leq 2 \mathbf{ex}_B(n, \mathcal{L}).$$

Assume now that we wish to have an upper bound on $\mathbf{ex}(m, n, \mathcal{L})$, where $n \gg m$. One way to get such an upper bound is to partition the n vertices into subsets of size $\approx m$. If, e.g, we know that $\mathbf{ex}(m, m, \mathcal{L}) \leq cm^{1+\gamma}$, then we obtain that

$$\mathbf{ex}(m, n, \mathcal{L}) \leq \frac{n}{m} \cdot \mathbf{ex}(m, m, \mathcal{L}) \leq cnm^\gamma. \quad (2.5)$$

This often helps, however, occasionally it is too weak. Erdős formulated

Conjecture 2.16. *If $n > m^2$ then $\mathbf{ex}(m, n, C_6) = O(n)$.*

Later this conjecture was made more precise, by Erdős, A. Sárközy and T. Sós, and proved by Gyóri, see Section 14.2 and [134].

We start with a trivial lemma.

Lemma 2.17. *Let d be the average degree in G_n , i.e. $d := 2e(G_n)/n$. Then G_n contains a G_m with $d_{\min}(G_m) > d/2$.*

To solve the cube-problem, Erdős and Simonovits used two reductions. The first one was a reduction to bipartite graphs, see Section 2.14. The other one eliminates the degrees are much higher than the average.

Definition 2.18 (Δ -almost-regularity). G is Δ -regular if $d_{\max}(G) < \Delta \cdot d_{\min}(G)$.

Theorem 2.19 (Δ -almost-regularization [90]). *Let $e(G_n) > n^{1+\alpha}$, and $\Delta = 20 \cdot 2^{(1/\alpha)^2}$. Then there is a Δ -almost-regular $G_m \subset G_n$ for which*

$$e(G_m) > \frac{2}{5}m^{1+\alpha}, \quad \text{where} \quad m > n^{\alpha \frac{1-\alpha}{1+\alpha}},$$

unless n is too small.

This means that whenever we wish to prove that $\mathbf{ex}(n, \mathcal{L}) = O(n^{1+\alpha})$, we may restrict ourselves to bipartite Δ -almost-regular graphs.

It would be interesting to understand the limitations of this lemma better. The next remark and problem are in this direction.

Remark 2.20. By the method of random graphs one can show [90] that for every Δ and n and $\varepsilon > 0$, there is G_n with $e(G_n) = \lfloor n^{3/2} \rfloor$ which does not have a Δ -almost-regular subgraph G_m with $e(G_m) > \varepsilon \sqrt{nm}$.

Problem 2.21 (Erdős-Simonovits [90]). Is it true that for every Δ there exists an $\varepsilon > 0$ such that every G_n , with $e(G_n) = \lfloor n \log n \rfloor$, contains a Δ -almost-regular subgraph G_m , with $e(G_m) > \varepsilon m \log m$ where $m \rightarrow \infty$ when $n \rightarrow \infty$?

2.4. Excluding complete bipartite graphs

Certain questions from topology (actually, Kuratowski theorem on planar graphs) led to Zarankiewicz problem [254]. After some weaker results Kővári, T. Sós and Turán proved the following theorem, already mentioned in Section 1.1.

Theorem 2.22 (Kővári–T. Sós–Turán, [164]). *Let $K_{a,b}$ denote the complete bipartite graph with a and b vertices in its color-classes. Then*

$$\mathbf{ex}(n, K_{a,b}) \leq \frac{1}{2} \sqrt[a]{b-1} \cdot n^{2-(1/a)} + \frac{a-1}{2}n. \quad (2.6)$$

Remarks 2.23. (a) If $a \neq b$ then (2.6) is better if we apply it with $a < b$.

(b) We know from Theorem 2.2 that $\mathbf{ex}(n, \mathcal{L}) = o(n^2)$ if and only if \mathcal{L} contains a bipartite L . Actually Claim 2.1 and Theorem 2.22 show that if $\mathbf{ex}(n, \mathcal{L}) = o(n^2)$ then $\mathbf{ex}(n, \mathcal{L}) = O(n^{2-c})$, for some constant $c > 0$.

Conjecture 2.24 ([164], see also e.g. [70]). *The upper bound in Theorem 1.2 is sharp:*

$$\mathbf{ex}(n, K_{a,b}) > c_{a,b}n^{2-(1/a)}.$$

Sketch of proof of Theorem 2.22. The number of a -stars $K_{a,1}$ in a graph G_n is $\sum \binom{d_i}{a}$ where d_1, \dots, d_n are the degrees in G_n . If G_n contains no $K_{a,b}$ then at most $b-1$ of these a -stars can share the same set of endpoints. We obtain

$$\sum \binom{d_i}{a} = \text{the number of } a\text{-stars} \leq (b-1) \binom{n}{a}. \quad (2.7)$$

Extending $\binom{n}{a}$ to all $x > 0$ by

$$\binom{x}{a} := \begin{cases} \frac{x(x-1)\dots(x-a+1)}{a!} & \text{for } x \geq a-1, \\ 0 & \text{otherwise} \end{cases}$$

we have a convex function. Then Jensen's Inequality implies that, the left hand side in (2.7) is at least $n \binom{2e(G)/n}{a}$, and the result follows by an easy calculation. \square

Remark 2.25. Slightly changing the above proof we get analogous upper bounds on $e(G_n)$ in all three cases of Problem 2.10.

We shall return to these questions in Sections 3.1, 3.4 where we shall discuss some improvements of the upper bound and also some lower bounds. **Further sources to read:** Guy [128], Znám: [257], [256], Guy–Znám [129].

2.5. Probabilistic lower bound

The theory of random graphs is an interesting, important, and rapidly developing subject. The reader wishing to learn more about it should either read the original papers of Erdős, e.g., [61], [62], Erdős and Rényi, e.g., [84], or some books, e.g., Bollobás, [27], Janson, Łuczak and Ruciński, [149], Molloy and Reed [192].

Theorem 2.26 (Erdős-Rényi First Moment method). *Let $\mathcal{L} = \{L_1, \dots, L_t\}$ be a family of graphs, and let*

$$c = \max_j \min_{H \subseteq L_j} \frac{v(H)}{e(H)}, \quad \gamma = \max_j \min_{H \subseteq L_j} \frac{v(H) - 2}{e(H) - 1}, \quad (2.8)$$

where the minimum is taken only for subgraphs H where the denominator is positive.

(a) Let G_n be a graph of order n chosen uniformly, at random, from graphs with E_n edges. For every $\varepsilon > 0$ there exists a $\delta > 0$ such that if $E_n < \delta n^{2-c}$, then the probability that G_n contains at least one $L \in \mathcal{L}$ is at most ε .

(b) If we know only $E_n < \varepsilon n^{2-\gamma}$, then the probability that G_n contains at least $\frac{1}{2}E_n$ copies of $L \in \mathcal{L}$ is at most ε .

This implies that

$$\mathbf{ex}(n, \mathcal{L}) > c_{\mathcal{L}} n^{2-\gamma} \geq c_{\mathcal{L}} n^{2-c} \quad (2.9)$$

with $c \geq \gamma > 0$ defined above.

Remarks 2.27. (a) A graph L is called balanced if the minimum in (2.8), for c , is achieved for $H = L$. Erdős and Rényi formulated their result containing Theorem 2.26(a) only for balanced graphs L . The part we use is trivial from their proof.

(b) Later Bollobás extended the Erdős-Rényi theorem to arbitrary \mathcal{L} .

(c) Győri, Rothschild and Ruciński achieved the generality by embedding any graph into a balanced graph [136].

Corollary 2.28. *If a finite \mathcal{L} contains no trees,¹³ then for some $c_{\mathcal{L}} > 0$, $\mathbf{ex}(n, \mathcal{L}) \geq c_{\mathcal{L}} n^{1+c}$.*

¹³neither forests

Mostly the weaker Theorem 2.26(a) implies Corollary 2.28: it does, whenever \mathcal{L} is finite and each $L \in \mathcal{L}$ contains at least two cycles in the same component. However, for cycles we need the stronger Theorem 2.26(b).

For example, for $L = K_{a,b}$ we have $c = a^{-1} + b^{-1}$. Then, for c_0 sufficiently small, the probability that a graph G_n with $c_0 n^{2-c}$ edges does not contain $K_{a,b}$ is positive. Hence

$$\mathbf{ex}(n, K_{a,b}) \geq c_0 n^{2-(1/a)-(1/b)}.$$

Comparing this with the Kővári-T. Sós-Turán theorem (Theorem 2.22), we see that the exponent is sharp there, in some sense, if a is fixed while $b \rightarrow \infty$.

Proof of Theorem 2.26. Consider the random graph G_n with n labeled vertices, in which each edge occurs independently, with the same probability p . For each L_j , choose a subgraph H_j which attains the inner minimum for γ , in (2.8). Let $h_j := v(H_j)$, $e_j := e(H_j)$, and let α_j denote the number of copies of H_j in K_{h_j} , and β_j denote the expected number of copies of H_j in G_n .

Clearly, K_n contains $\alpha_j \binom{n}{h_j}$ copies of H_j . For each copy H of H_j , define a random “indicator” variable $k_H = k_H(G_n) = 1$ if $H \subseteq G_n$, and 0 otherwise. Since the number of copies of H_j in G_n is just $\sum_{H \subseteq K_n} k_H$, therefore, if \mathbb{E} denote the expected value, then

$$\beta_j = \sum_{H \subseteq K_n} \mathbb{E}(k_H) = \alpha_j \binom{n}{h_j} p^{e_j}.$$

Summing over j and taking $p = c_1 n^{-c}$, (for some $c_1 \in (0, 1)$) we get

$$\sum_j \beta_j \leq t \max_j \alpha_j \binom{n}{h_j} p^{e_j} \leq t \max c_1 n^{h_j - c e_j} = t c_1 n^{2-c}.$$

Now let $\eta(G_n) = e(G_n) - \sum_j \beta_j$. Then, for c_1 sufficiently small, the expected value is

$$\mathbb{E}(\eta(G_n)) > \frac{1}{2} \binom{n}{2} p > \frac{1}{5} c_1 n^{2-c}.$$

Hence there exists a G_n with $\eta(G_n) > \frac{1}{5} c_1 n^{2-c}$. Delete an edge from each H_j in this G_n . The resulting graph contains no L_j , and has at least $\frac{1}{5} \binom{n}{2} p \geq \frac{1}{11} c_1 n^{2-c}$ edges, completing the proof. \square

Remarks 2.29 (How did the probabilistic methods start?). Mostly we write that applications of the Random Graphs (probabilistic method) started when Erdős (disproving a conjecture of Turán on the Ramsey Numbers) proved the existence of graphs G_n without complete subgraphs of order $2 \log n$ and without independent sets of size $2 \log n$.

1. Erdős himself remarks (e.g., in [64]) that perhaps Szele was the first who applied this method in Graph Theory. (Erdős – in his birthday volume [76] – also mentions an even earlier application of J. Erőd but we did not succeed in locating that source.)
2. Perhaps the earliest case of applying probabilistic methods was that of Paul Turán’s proof of the Hardy-Ramanujan Theorem [241], where – reading the paper – it is obvious that Turán gave a probabilistic proof of a beautiful and important theorem, using the Chebishev inequality. However, either Turán did not realize that this is an application of the probabilistic method or he did not wish to burden the reader with that.
3. An important application of the probabilistic methods was that of Claude Shannon, when he constructed random codes.

Applying Theorem 2.26 to some families of cycles we obtain

Corollary 2.30. *For some constant $c_m > 0$,*

$$\mathbf{ex}(n, \{C_3, \dots, C_m\}) \geq c_m n^{1 + \frac{1}{m-1}}.$$

Erdős’ even cycles theorem asserts that $\mathbf{ex}(n, C_{2t}) = O(n^{1+(1/t)})$, and this upper bound is probably sharp.¹⁴ The random method (that is, Theorem 2.26) yields a lower bound of $cn^{1+\frac{1}{2t-1}}$, a weaker result. Simonovits thinks that it is unlikely that Theorem 2.26 ever yields a sharp bound for a finite family.¹⁵

Corollary 2.30 is used in the next section to prove that $\mathbf{ex}(n, \mathcal{L}) = O(n)$ if and only if contains a tree or forest.

¹⁴The reference is missing here, since Erdős did formulate this theorem but never have published a proof of it, as far as we know.

¹⁵Some related results of G. Margulis, and A. Lubotzky, R. Phillips and P. Sarnak will be discussed in Section 4.9.

2.6. Classification of extremal problems

The extremal graph problems can be classified in several ways. Here we shall speak of (a) *non-degenerate*, (b) *degenerate* and (c) *linear* extremal problems.

For Case (a) Theorem 2.3 provides an appropriately good description of the situation. In Case (b) $p(\mathcal{L}) = 1$. Here the “main term” disappears, $(1 - \frac{1}{p}) = 0$; therefore “the error terms dominate”. Case (c) will be discussed here shortly and in Sections 6 and 9 in more details.

The classification immediately follows from the following theorems:

Theorem 2.31. $\mathbf{ex}(n, \mathcal{L}) = o(n^2)$ if and only if \mathcal{L} contains a bipartite graph. Actually, if \mathcal{L} contains a bipartite graph then $\mathbf{ex}(n, \mathcal{L}) = O(n^{2-c})$ for, e.g., $c = 2/v(L)$ for any bipartite $L \in \mathcal{L}$. If \mathcal{L} does not contain bipartite graphs, then $\mathbf{ex}(n, \mathcal{L}) \geq \left\lfloor \frac{n^2}{4} \right\rfloor$.

Theorem 2.32. For finite \mathcal{L} , $\mathbf{ex}(n, \mathcal{L}) = O(n)$ if and only if \mathcal{L} contains a tree, or a forest. If $L \in \mathcal{L}$ is a tree or a forest, then, for $v(L) \geq 3$,

$$\mathbf{ex}(n, \mathcal{L}) < (v(L) - 2)n. \quad (2.10)$$

Theorem 2.33 (Erdős [61, 62]). If \mathcal{L} is finite and no $L \in \mathcal{L}$ is a tree, then $\mathbf{ex}(n, \mathcal{L}) > n^{1+c_{\mathcal{L}}}$ for some $c_{\mathcal{L}} > 0$.

Theorem 2.34 (Erdős [62], Bondy and Simonovits [32]). Given an integer k , for some constants $c_k, \tilde{c}_k > 0$,

$$c_k n^{1+\frac{1}{2k-1}} < \mathbf{ex}(n, \{C_3, \dots, C_{2k}\}) \leq \mathbf{ex}(n, C_{2k}) \leq \tilde{c}_k n^{1+\frac{1}{k}}. \quad (2.11)$$

Proof of Theorems 2.31, 2.32, and 2.33. If there is a bipartite $L \in \mathcal{L}$, then Theorem 2.22 implies the sharper upper bound of Theorem 2.31. Indeed, for $v = v(L)$, by $L \subseteq K(\lfloor v/2 \rfloor, v)$, we have,

$$\mathbf{ex}(n, \mathcal{L}) \leq \mathbf{ex}(n, L) \leq \mathbf{ex}(n, K(\lfloor v/2 \rfloor, v)) < \frac{1}{2} \sqrt{2v} \cdot n^{2-(2/v(L))} = O(n^{2-c}).$$

If the minimum chromatic number $p = p(\mathcal{L}) \geq 3$, then $T_{n,p}$ contains no forbidden $L \in \mathcal{L}$. Therefore

$$\mathbf{ex}(n, \mathcal{L}) > e(T_{n,2}) \geq e(T_{n,p}) = \left(1 - \frac{1}{p}\right) \binom{n}{2} + O(n).$$

Actually, $e(T_{n,2}) = \left\lceil \frac{n^2}{4} \right\rceil$. This completes the proof of Theorem 2.31. \square

It is easy to show that if G_n has minimum degree at least $r - 1$, then it contains every tree T_r (by induction on r). An induction on n yields (2.10), implying half of Theorem 2.32, when \mathcal{L} contains a tree (or a forest). If \mathcal{L} is finite and contains no trees, i.e., all the forbidden graphs contain some cycles, then we use Theorem 2.34, or simply Corollary 2.28, proved by probabilistic methods.¹⁶

Remark 2.35 (Infinite families). For infinite families the situation is different: if e.g. \mathcal{C} is the family of all cycles, then $\mathbf{ex}(n, \mathcal{C}) = n - 1$: all graphs but the forests are excluded. There are many further families without trees where the extremal number is linear, see Section 9.

Proof of Theorem 2.34. The lower bound comes from a random graph argument of Erdős. Concentrate on the upper bound. If we are not interested in the value of the constant, then we can basically use the following argument: Take a graph G_n with $cn^{1+\alpha}$ edges. Delete its minimum degree vertex, then the minimum degree vertex in the remaining graph, etc. At the end we get a G_m with minimum degree at least c_1m^α . In the obtained graph G_m fix a vertex x and denote by S_j the set of vertices at distance j from x . If $\mathbf{girth}(G_n) > 2k$, – as we assumed – then *basically* $|S_j| > d_{\min}(G_m) \cdot |S_{j-1}|$. Hence $m > |S_k| > c_1^k m^{\alpha k}$. So $\alpha \leq 1/k$. \square

Assume for a second that G_n itself is asymptotically regular:

$$\frac{d_{\min}(G_n)}{d_{\max}(G_n)} \rightarrow 1.$$

Then the previous argument asserts that $d := d_{\min}(G_n) < n^{1/k}$. Therefore

$$e(G_n) \leq \left(\frac{1}{2} + o(1) \right) nd \approx \frac{1}{2} n^{1+\frac{1}{k}}.$$

We shall return to the case of excluded trees, namely, to the Erdős-Sós conjecture on the extremal number of trees, and to the related Komlós-Sós conjecture in Section 6. One final question could be if $\mathbf{ex}(n, \mathcal{L})$ can be sublinear. This is answered by the following trivial result.

Theorem 2.36. *If \mathcal{L} is finite and $\mathbf{ex}(n, \mathcal{L}) < [n/2]$, then $\mathbf{ex}(n, \mathcal{L}) = O(1)$.*

¹⁶There are also deterministic proofs of Corollary 2.28, e.g., via the Margulis–Lubotzky–Phillips–Sarnak construction of Ramanujan graphs, see Construction 4.43.

Proof. Consider $n/2$ independent edges: this must contain an $L_1 \in \mathcal{L}$. Hence, there is an $L_1 \in \mathcal{L}$ contained in the union of t independent edges, for some t . Also, there exists an $L_2 \subseteq K(1, n-1)$. Hence an extremal graph S_n has bounded degrees and bounded number of independent edges. This proves 2.36. \square

Theorem 2.36 easily extends to hypergraphs.

2.7. General conjectures on bipartite graphs

We have already formulated Conjecture 1.6 on the rational exponents. We have to remark that for hypergraphs this does not hold: the Behrend construction [21] is used to get lower bounds in the Ruzsa–Szemerédi Theorem, (Thm 1.9), showing that there is no rational exponent in that case. Yet, Erdős and Simonovits conjectured that for ordinary graphs there is. One could also conjecture the inverse extremal problem:

Conjecture 2.37. *For every rational $\alpha \in (0, 1)$ there is a finite \mathcal{L} for which $c_1 n^{1+\alpha} < \mathbf{ex}(n, \mathcal{L}) < c_2 n^{1+\alpha}$, for some constants $c_1, c_2 > 0$.*

The third conjecture to be mentioned here is on “compactness” [93]:

Conjecture 2.38. *For every finite \mathcal{L} there is an $L \in \mathcal{L}$ for which $\mathbf{ex}(n, \mathcal{L}) > c \cdot \mathbf{ex}(n, L)$, for some constants $c > 0$.*

3. Excluding complete bipartite graphs

3.1. Bipartite C_4 -free graphs and the Zarankiewicz problem

Turán type extremal results (and Ramsey results as well) can often be applied in Mathematics, even outside of Combinatorics. Turán himself explained this applicability by the fact that – in his opinion – the extremal graph results were generalizations of the Pigeon Hole Principle.

Recall that $Z(m, n, a, b)$ denotes the maximum number of 1’s in an $m \times n$ matrix not containing an $a \times b$ minor consisting exclusively of 1’s. In 1951 Zarankiewicz [254] posed the problem of determining $Z(n, n, 3, 3)$ for $n \leq 6$, and the general problem has also become known as *the problem*

of Zarankiewicz.¹⁷ Obviously, $Z(m, n, 1, b) = m(b - 1)$ (for $n \geq b - 1$). Observe that $Z(m, n, a, b) = \mathbf{ex}^*(m, n, K_{a,b})$ (where $\mathbf{ex}^*(m, n, \mathcal{L})$ was defined following Remark 2.10.) Considering the adjacency matrix of a $K_{a,b}$ -free graph on n vertices we get $2\mathbf{ex}(n, K_{a,b}) \leq Z(n, n, a, b)$. We will use this upper bound many times.

We will see that the easy upper bound in Theorem 2.22 is pretty close to the truth for $a \leq 2$. Actually, Kővári, T. Sós and Turán [164] proved an upper bound for the Zarankiewicz function

$$Z(m, n, a, b) \leq \sqrt[a]{b-1} \cdot mn^{1-(1/a)} + (a-1)n \tag{3.1}$$

which was slightly improved by Znám [257], [256]. (he halved the last term to $(a-1)n/2$ in the case of $m = n$) and Guy [128].

A bipartite graph $G[M, N]$ where $|M| = m$, $|N| = n$ is C_4 -free if its “bipartite” $m \times n$ adjacency matrix contains no 2×2 full 1 submatrix.¹⁸ In other terminology, the hypergraph defined by the rows of this matrix is linear, and their hyperedges pairwise meet in at most one element. There is an important class of such hypergraphs, the Steiner k -systems $S(n, k, 2)$. A family \mathcal{S} of k -subsets of an n -set N is a Steiner k -system if every pair of elements is covered exactly once. For such an \mathcal{S} , clearly, $|\mathcal{S}| = m = \binom{n}{2} / \binom{k}{2}$. Such families are known to exist for $(m, n, k) = (q^2 + q + 1, q^2 + q + 1, q + 1)$ (called finite projective planes of order q), and $(m, n, k) = (q^2 + q, q^2, q)$ (affine planes) whenever q is a power of a prime. Also for any given k there exists an $n_0(k)$ such that $S(n, k, 2)$ exists for all *admissible* $n > n_0(k)$, i.e., when $(n-1)/(k-1)$ and $n(n-1)/k(k-1)$ are integers (Wilson’s existence theorem [248]).

Kővári, T. Sós and Turán [164] proved that

Theorem 3.1. $Z(n, n, 2, 2) = (1 + o(1))n^{3/2}$, and

$$Z(n, n, 2, 2) < [n^{3/2}] + 2n. \tag{3.2}$$

Further, if p is a prime, then

$$Z(p^2 + p, p^2, 2, 2) = p^3 + p^2.$$

¹⁷In Graph Theory two problems are connected to Zarankiewicz’ name: the extremal problem for matrices that we shall discuss here and the Crossing Number conjecture which is not our topic. Actually, the crossing number problem comes from Paul Turán, see [244].

¹⁸Here the “bipartite adjacency matrix” $A = (a_{ij})_{m \times n}$ is defined for a bipartite graph $G[U, V]$ and $a_{ij} = 1$ if $u_j \in U$ is joined to $v_j \in V$, otherwise $a_{ij} = 0$.

Reiman [206] returned to this topic, (see also [207]), slightly improving (3.2)

Theorem 3.2 (Reiman [206]).

$$Z(m, n, 2, 2) \leq \frac{1}{2} \left(m + \sqrt{m^2 + 4mn(n-1)} \right). \quad (3.3)$$

For large $m, n \rightarrow \infty$, and $m = o(n^2)$, this yields

$$Z(m, n, 2, 2) \leq \left(\frac{1}{2} + o(1) \right) n\sqrt{m}.$$

Further, for $m = n$, we get

$$Z(n, n, 2, 2) \leq \frac{1}{2}n(1 + \sqrt{4n-3}) \approx n\sqrt{n}. \quad (3.4)$$

Reiman also provides infinitely many graphs, using Finite Geometries, showing the sharpness of (3.3) and (3.4). We have equality when $m = n(n-1)/k(k-1)$ and a Steiner system $S(n, k, 2)$ exists. Thus he determined the case

$$Z(n, n, 2, 2) = \frac{1}{2}n(1 + \sqrt{4n-3}) = (q^2 + q + 1)(q + 1) \quad (3.5)$$

for $m = n = q^2 + q + 1$ when a projective plane of order q exists. Actually, in [207], Reiman also speaks about Zarankiewicz-extremal graphs connected to incidence-graphs of higher dimensional finite geometries.

Since Reiman's theorem the theory of finite geometries developed tremendously. We cite here a recent result whose proof used the most modern tools and stability results.

Theorem 1 (Damásdi, Héger, and Szőnyi [57]). Let $q \geq 15$, and $c \leq q/2$. Then

$$Z(q^2 + q + 1 - c, q^2 + q + 1, 2, 2) \leq (q^2 + q + 1 - c)(q + 1).$$

Equality holds if and only if a projective plane of order q exists. Moreover, graphs giving equality are subgraphs of the bipartite incidence graph of a projective plane of order q obtained by omitting c rows of its incidence matrix.

They proved many more exact results when a projective plane of order q exists. The extremal configurations are submatrices of the incidence matrix of a projective plane.

$$\begin{aligned} Z(q^2 + c, q^2 + q, 2, 2) &= q^2(q + 1) + cq \quad (0 \leq c \leq q + 1), \\ Z(q^2 - q + c, q^2 + q - 1, 2, 2) &= (q^2 - q)(q + 1) + cq \quad (0 \leq c \leq 2q), \\ Z(q^2 - 2q + 1 + c, q^2 + q - 2, 2, 2) &= (q^2 - 2q + 1)(q + 1) + cq \quad (0 \leq c \leq 3(q - 1)). \end{aligned}$$

These refer to bipartite host graphs. As we will see later, such exact results are rare for the general (non-bipartite) case. To estimate $\mathbf{ex}(n, C_4)$ seems to be harder, because the corresponding 0-1 matrices, the incidence matrix of a graph, should be symmetric.

3.2. Finite Geometries and the C_4 -free graphs

The method of finite geometric constructions is very important and powerful in combinatorics. In particular, it is often the best way to obtain lower bounds. It is for this reason that we include this section.

We give several constructions: the first two show that the Kővári-T. Sós-Turán theorem (Theorem 2.22) is sharp for both $K_{2,2}$ and $K_{3,3}$.

Remark 3.3. When we write that an upper bound is sharp, mostly we mean that it is sharp up to a multiplicative constant: it yields the correct exponent. There are a few exceptions, where sharpness means that the ratio of the upper and lower bounds tends to 1. This is the case for $C_4 = K_{2,2}$ and we have this also for $K_{3,3}$. Here, however, the matching upper bound for Construction 3.20 below is given not by Theorem 2.22 but by the Füredi improvement [112].

Perhaps the application of finite geometries in Extremal Graph Theory started in the Erdős paper, with the construction of Eszter Klein [60], to prove the sharpness of Theorem 1.15. The expression “Finite Geometry” was not mentioned there. We skip the description of this whole story, since it was described in several places, e.g., [227], [228].

Much later, Erdős and Rényi [85] used finite geometry for a diameter-extremal problem. This is a very large area, connected to our problems, yet we have to skip it. The interested reader is referred to [85], (translated into English in [208]).

Sharp extremal graph results were obtained by Reiman [206] and a Polarity Graph was used in [86] and [36] to give asymptotically sharp lower bound on $\mathbf{ex}(n, C_4)$. This lower bound can also be found in [85], **implicitly**: Erdős and Rényi considered the diameter-extremal problem, and do mention the properties of this graph.

The real breakthrough came by the Erdős-Rényi-T. Sós paper [86], (sharp lower bound for C_4) and by the Brown paper [36], providing asymptotically sharp lower bounds for $\mathbf{ex}(n, C_4)$ and for $\mathbf{ex}(n, K_{3,3})$. (See Remark 3.3.)

We know from Theorem 2.22 that $\mathbf{ex}(n, C_4) \leq \frac{1}{2}n^{3/2} + o(n^{3/2})$, but is this result sharp? In analyzing the proof, we realize that if it is sharp (that is, if there are infinitely many graphs G_n not containing C_4 and having $\approx \frac{1}{2}n\sqrt{n}$ edges). Almost all degrees are $\approx \sqrt{n}$ and almost every pair of vertices must have a common neighbor (and no pair has two). This suggests that the neighborhoods $N(x)$ behave much like the lines in a projective plane, in that the following statement “almost” holds: any two vertices lie in a common set, and any two sets intersect in one vertex.

Theorem 3.4 (Erdős-Rényi-T. Sós [86], and Brown [36], see also [164]).

$$\mathbf{ex}(n, C_4) = \frac{1}{2}n^{3/2} + O(n^{3/2-c}).$$

For the lower bound for $\mathbf{ex}(n, C_4)$ we use the following

Construction 3.5. *Let p be a prime, $n = p^2 - 1$. Construct a graph as follows: the vertices are the $p^2 - 1$ non-zero pairs (x, y) of residues (modulo p), and (x, y) is joined to (a, b) by an edge if $ax + by = 1$. (This graph may contain loops, but we simply delete them.)*

With $n = p^2 - 1$, the resulting graph H_n has the necessary properties to show the sharpness of Theorem 2.22 for C_4 :

(a) for a given pair (a, b) , mostly there are p solutions to $ax + by = 1$, so that, even after the loops are deleted, there are at least $\frac{1}{2}(p^2 - 1)(p - 1)$ edges in H_n and hence $e(H_n) > \frac{1}{2}n^{3/2} - n$;

(b) if H_n had a 4-cycle with vertices (a, b) , (u, v) , (a', b') and (u', v') , then the two equations $ax + by = 1$ and $a'x + b'y = 1$ would have two solutions, which is impossible. Since the primes are “dense” among the integers, this completes the proof of the sharpness of Theorem 2.22 for $a = b = 2$.

Remark 3.6. An alternative possibility is to use the much more symmetric polarity graph of the projective plane (we explain this in the next section):

here we used the Affine Geometric Variant because *here* we did not wish to use anything from Projective Geometry.

3.3. Excluding C_4 : Exact results

The polarity graph¹⁹, used in [85], was also used in [86] and [36] to prove that

$$\mathbf{ex}(n, C_4) \geq \frac{1}{2}q(q+1)^2, \quad \text{for } n = q^2 + q + 1. \quad (3.6)$$

if q is a prime power.

Construction 3.7 (The Polarity Graph from the finite field). *Assume that q is a prime power. Consider the Finite Field $GF(q)$. The vertices of our graph are the equivalence classes of the non-zero triples $(a, b, c) \in GF(q)^3$ where two of them, (a, b, c) and (a', b', c') are considered the same if $(a', b', c') = \lambda(a, b, c)$ for some $\lambda \neq 0$. There are $(q^3 - 1)/(q - 1) = q^2 + q + 1$ such classes. Further, the equivalence class of (a, b, c) is connected by an edge to the class of (x, y, z) if $ax + by + cz = 0$. Finally, we delete the $q + 1$ loops, i.e. those edges, where $a^2 + b^2 + c^2 = 0$. This graph is C_4 -free and it has $\frac{1}{2}(n(q+1) - (q+1))$ edges.*

In general, a polarity corresponds to a *symmetric* incidence matrix of a finite plane of size $(q^2 + q + 1) \times (q^2 + q + 1)$. According to a theorem of Baer [13] such a matrix has at least $q + 1$ non-zero elements in its diagonal. Therefore using the polarity graph we cannot avoid loosing on loops. This way Erdős, Rényi and Sós [86] showed that indeed $\mathbf{ex}(q^2 + q + 1, C_4) < \frac{1}{2}n(q + 1)$. Yet one could hope to get a better construction. Erdős conjectured [66], [71] that there are no better constructions, that is, (3.6) is sharp if $n = q^2 + q + 1$, (q is a prime power).

Füredi settled this conjecture in the following sense: First he proved [103] that if $q = 2^k$, then Erdős' conjecture holds. Next he settled the case $q \geq q_0$. Later he found a much shorter proof of the weaker assertion that the Polarity graphs are extremal; however, this shorter version did not give the extremal structure. So Füredi published the shorter version, while the longer version can be found on his homepage.

¹⁹ These C_4 -free graphs were studied earlier in finite geometry. The bipartite point-line incidence graph appeared in Levi's book (1942) and polarity graphs (modulo loops) obtained from *Levi graph* had been described already by Artzy (1956). For more details and references see Bondy [30].

Theorem 3.8 (Füredi [111], [104]). *If $q \neq 1, 7, 9, 11, 13$ and $n = q^2 + q + 1$, then $\mathbf{ex}(n, C_4) \leq \frac{1}{2}q(q+1)^2$ and for $q > 13$ the extremal graphs are obtained from a polarity of a finite projective plane. Hence if $q > 13$ is a prime power, then $\mathbf{ex}(n, C_4) = \frac{1}{2}q(q+1)^2$.*

The second part of this result probably holds for $q \in \{7, 9, 11, 13\}$, too.
Recently a new sharp construction has been found for $n = q^2 + q$.

Theorem 3.9 (Firke, Kosek, Nash and Williford [102]). *Suppose that q is even, $q > q_0$. Then*

$$\mathbf{ex}(q^2 + q, C_4) \leq \frac{1}{2}q(q+1)^2 - q.$$

Consequently, if $q > q_0$, $q = 2^k$ and $n = q^2 + q$ then $\mathbf{ex}(n, C_4) = q(q+1)^2 - q$.

They also announced that in a forthcoming paper they show that for all but finitely many even q , any $S_n \in \mathbf{EX}(q^2 + q, C_4)$ is derived from an orthogonal polarity graph by removing a vertex of minimum degree (the 1-vertex-truncated Polarity graph, see Construction 3.7). This result shows a kind of stability of the Polarity graph. More generally, McCuaig (private communication, 1985) **conjectured** that each extremal graph is a subgraph of some polarity graph. So this is true for infinitely many cases, but one of the present authors strongly disagrees and he believes just the opposite that for e.g., $n = q^2 + q + 2$ maybe the extremal graphs are obtained by *adding* an extra vertex and some edges to a polarity graph.

Remark 3.10. W. McCuaig calculated $\mathbf{ex}(n, C_4)$ for $n \leq 21$ (unpublished letter, 1985). Clapham, Flockart and Sheehan determined the corresponding extremal graphs [48], and Yuansheng and Rowlinson [253], – using computers, – extended these results to $n \leq 31$. (They also determined the graphs in $\mathbf{EX}(n, C_6)$ for $n \leq 26$, [252].) Garnick, Kwong, Lazebnik, and Nieuwejaar [122], [123] determined the values of $\mathbf{ex}(n, \{C_3, C_4\})$ for all $n \leq 30$.

3.4. Excluding $K(2, t+1)$, $t > 1$

A slightly sharper form of the upper bound (3.1) was presented by Hyltén-Cavallius [146]

$$Z(m, n, 2, k) \leq \frac{1}{2}n + \left\{ (k-1)nm(m-1) + \frac{1}{4}n^2 \right\}^{1/2}. \quad (3.7)$$

Obviously, for fixed k and large values of n, m , if $n = o(m^2)$, then the right hand side of (3.7) is $\approx \sqrt{k-1}m\sqrt{n}$. Using again the observation $2\mathbf{ex}(n, K_{2,t+1}) \leq Z(n, n, 2, t+1)$ one obtains the upper bound

$$\mathbf{ex}(n, K_{2,t+1}) \leq \frac{1}{2}n\sqrt{tn-t+1/4} + (n/4). \quad (3.8)$$

The following theorem shows that the above (easy) upper bound is the best possible asymptotically.

Theorem 3.11 (Füredi [113]). *For any fixed $t \geq 1$*

$$\mathbf{ex}(n, K_{2,t+1}) = \frac{1}{2}\sqrt{t}n^{3/2} + O(n^{4/3}). \quad (3.9)$$

To prove this Theorem one needs an appropriate lower bound, a construction. Let q be a prime power such that $(q-1)/t$ is an integer. We will construct a $K_{2,t+1}$ -free graph G on $n = (q^2-1)/t$ vertices such that every vertex has degree q or $q-1$. We will explain this below (Construction 3.15). Then G has more than $(1/2)\sqrt{t}n^{3/2} - (n/2)$ edges. The gap between the lower and upper bounds is only $O(\sqrt{n})$ for $n = (q^2-1)/t$. The lower bound for the Turán number for all n then follows from the fact that such prime powers form a dense subsequence among the integers. This means that for every sufficiently large n there exists a prime q satisfying $q \equiv 1 \pmod{t}$ and $\sqrt{nt} - n^{1/3} < q < \sqrt{nt}$ (see [145]).

Construction 3.15 below is inspired by constructions of Hytén-Cavallius and Mörs given for Zarankiewicz's problem $Z(n, n, 2, t+1)$.

Theorem 3.12 (Hytén-Cavallius [146]). $Z(n, n, 2, 3) = \sqrt{2}n^{3/2} + o(n^{3/2})$.
Also

$$\sqrt{\lfloor k/2 \rfloor} \leq \liminf_{n \rightarrow \infty} \frac{Z(n, n, 2, k)}{n^{3/2}}.$$

Theorem 3.13 (Mörs [194]). *For all $t \geq 1$,*

$$\frac{Z(n, n, 2, t+1)}{n^{3/2}} \rightarrow \sqrt{t}, \quad \text{as } n \rightarrow \infty.$$

The topic was so short of constructions that, as a first step, P. Erdős [66, 69] even proposed the problem whether $\lim_t(\liminf_n \mathbf{ex}(n, K_{2,t+1})n^{-3/2})$ goes to ∞ as $t \rightarrow \infty$.

Remark 3.14. Here we see three distinct quantities, exactly as it is described in Problem 2.10. $Z(m, n, 2, t + 1) = \mathbf{ex}^*(m, n, K_{2,t+1})$, estimated from below by Mörs, by a construction, and $\mathbf{ex}(m, n, K_{2,t+1})$ estimated by Füredi by the same construction. Füredi showed that the matrix of Mörs contains neither a $t \times 2$ submatrix, nor a $2 \times (t + 1)$ submatrix of 1's; finally, Füredi, slightly changing the definitions in Mörs's construction extended this "asymmetric matrix" result to the symmetric case and provided a non-bipartite graph, proving (3.9).

Construction 3.15. Let $GF(q)$ be the q -element finite field, and let $h \in GF(q)$ be an element of order t . This means, that $h^t = 1$ and the set $H = \{1, h, h^2, \dots, h^{t-1}\}$ form a t -element subgroup of $GF(q) \setminus \{0\}$. For $q \equiv 1 \pmod{t}$ such an element $h \in GF(q)$ always exists.

We say that $(a, b) \in GF(q) \times GF(q)$, $(a, b) \neq (0, 0)$ is equivalent to (a', b') , in notation $(a, b) \sim (a', b')$, if there exists some $h^\alpha \in H$ such that $a' = h^\alpha a$ and $b' = h^\alpha b$. The elements of the vertex set V of G are the t -element equivalence classes of $GF(q) \times GF(q) \setminus (0, 0)$. The class represented by (a, b) is denoted by $\langle a, b \rangle$. Two (distinct) classes $\langle a, b \rangle$ and $\langle x, y \rangle$ are joined by an edge in G if $ax + by \in H$. This relation is symmetric, and $ax + by \in H$, $(a, b) \sim (a', b')$, and $(x, y) \sim (x', y')$ imply $a'x' + b'y' \in H$. So this definition is compatible with the equivalence classes.

For any given $(a, b) \in GF(q) \times GF(q) \setminus (0, 0)$ (say, $b \neq 0$) and for any given x and h^α , the equation $ax + by = h^\alpha$ has a unique solution in y . This implies that there are exactly tq solutions (x, y) with $ax + by \in H$. The solutions come in equivalence classes, so there are exactly q classes $\langle x, y \rangle$. One of these classes might coincide with $\langle a, b \rangle$ so the degree of the vertex $\langle a, b \rangle$ in G is either q or $q - 1$.

We claim that G is $K_{2,t+1}$ -free. First we show, that for $(a, b), (a', b') \in GF(q) \times GF(q) \setminus (0, 0)$, $(a, b) \not\sim (a', b')$ the equation system

$$ax + by = h^\alpha \quad \text{and} \quad a'x + b'y = h^\beta \quad (3.10)$$

has at most one solution $(x, y) \in GF(q) \times GF(q) \setminus (0, 0)$. Indeed, the solution is unique if the determinant $\det \begin{pmatrix} a & b \\ a' & b' \end{pmatrix}$ is not 0. Otherwise, there exists a c such that $a = a'c$ and $b = b'c$. If there exists a solution of (2) at all, then multiplying the second equation by c and subtracting it from the first one we get on the right hand side $h^\alpha - ch^\beta = 0$. Thus $c \in H$, contradicting the fact that (a, b) and (a', b') are not equivalent.

Finally, there are t^2 possibilities for $0 \leq \alpha, \beta < t$ in (3.10). The set of solutions again form t -element equivalence classes, so there are at most t equivalence classes $\langle x, y \rangle$ joint simultaneously to $\langle a, b \rangle$ and $\langle a', b' \rangle$. \square

Since then, there have been two additional almost optimal constructions, strongly related to the Construction 3.15 above.

Construction 3.16 (Lazebnik, Mubayi [167]). *Let $GF(q)^*$ be the finite field of order q without the zero element. Suppose $q \equiv 1 \pmod{t}$ and let H be the t -element multiplicative subgroup of $GF(q)^*$. Define the graph G^\times as follows. Let $V(G^\times) = (GF(q)^*/H) \times GF(q)$. For $\langle a \rangle, \langle b \rangle \in (GF(q)^*/H)$ and $x, y \in GF(q)$, make $(\langle a \rangle, x)$ adjacent to $(\langle b \rangle, y)$ if $x + y \in \langle ab \rangle$.*

This graph (after deleting the eventual loops) is $K_{2,t+1}$ -free and every vertex has degree $q - 1$ or $q - 2$. Actually, Construction 3.16 differs from Construction 3.15 only in that its vertex set is smaller and instead of using the rule that $\langle a, b \rangle$ is adjacent to $\langle x, y \rangle$ if $ax + by \in H$ they use the rule $ay + bx \in H$. This change allows them to generalize it to multipartite hypergraphs.

The following example works only if t is a power of a prime, and $t|q$.

Construction 3.17 (Lenz, Mubayi [173]). *Suppose that t divides q and let H be an additive subgroup of $GF(q)$ of order t . Define the graph G^+ as follows. Let $V(G^+) = (GF(q)/H) \times GF(q)^*$. We will write elements of $GF(q)/H$ as $\langle a \rangle$. It is the additive coset of H generated by a , $\langle a \rangle = \{h + a : h \in H\}$. For $\langle a \rangle, \langle b \rangle \in (GF(q)/H)$ and $x, y \in GF(q)^*$, make $(\langle a \rangle, x)$ adjacent to $(\langle b \rangle, y)$ if $xy \in \langle a + b \rangle$. (This, in fact, means that there exists an $h \in H$ such that $xy = a + b + h$).*

3.5. Excluding $K(3, 3)$, and improving the upper bound

The main result of this section is the description of the asymptotically sharp value of $\mathbf{ex}(n, K_{3,3})$.

Theorem 3.18 (Brown [36] and Füredi [112]).

$$\mathbf{ex}(n, K_{3,3}) = \frac{1}{2}n^{5/3} + O(n^{(5/3)-c}) \quad \text{for some } c > 0.$$

The lower bound can be obtained from Brown's example (discussed below as Construction 3.20) who gave a $(p^2 - p)$ -regular $K_{3,3}$ -free graph on p^3 vertices for each prime p of the form $4k - 1$.

Improving the upper bound in Theorem 2.22 Füredi showed that Brown's example is asymptotically optimal.

Theorem 3.19 (Füredi [112]). *For all $m \geq a$, $n \geq b$, $b \geq a \geq 2$ we have*

$$Z(m, n, a, b) \leq (b - a + 1)^{1/a} m n^{1-(1/a)} + (a - 1)n^{2-(2/a)} + (a - 2)m. \quad (3.11)$$

For fixed $a, b \geq 2$ and $n, m \rightarrow \infty$ the first term is the largest one for $n = O(m^{a/(a-1)})$. This upper bound is asymptotically optimal for $a = 2$ and for $a = b = 3$ ($m = n$). We obtain

$$\mathbf{ex}(n, K_{3,3}) \leq \frac{1}{2}Z(n, n, 3, 3) \leq \frac{1}{2}n^{5/3} + n^{4/3} + \frac{1}{2}n. \quad (3.12)$$

Alon, Rónyai and Szabó [11] gave an example (discussed as Construction 3.25) showing that

$$\mathbf{ex}(n, K_{3,3}) \geq \frac{1}{2}n^{5/3} + \frac{1}{3}n^{4/3} - C. \quad (3.13)$$

for some absolute constant $C > 0$ for every n of the form $n = p^3 - p^2$, p is a prime. Their example shows that the upper bound (3.12) (and (3.11)) is so tight that that we cannot leave out the second order term. It would be interesting to see whether (3.11) is tight for other values of a and b , too.

The first step of the proof of Theorem 3.19 is that given a $K_{a,b}$ -free graph G , we apply the original bound (3.1) to the bipartite subgraphs $G[N(x), V \setminus N(x)]$ generated by the neighborhood of a vertex x and its complement.

When Brown gave his construction, the matching upper bound of Theorem 3.19 was not known yet. He wrote that even the existence of $\lim_{n \rightarrow \infty} \mathbf{ex}(n, K_{3,3})/n^{5/3}$ was unknown.

Construction 3.20. *Let p be an odd prime $n = p^3$ and $d \in GF(p)$, $d \neq 0$ a quadratic residue if p is of the form $4k - 1$ and d be a non-residue otherwise. Construct a graph B_n whose vertices are the triples (x, y, z) of residue classes (modulo p) and whose edges join vertices (x, y, z) and (x', y', z') if*

$$(x - x')^2 + (y - y')^2 + (z - z')^2 = d. \quad (3.14)$$

It is easy to see that the graph B_n has $\frac{1}{2}n^{5/3} + O(n^{4/3})$ edges. Given a vertex (x', y', z') , the equation (3.14) has $p^2 - p$ solutions by a theorem of Lebesgue. Thus (x', y', z') has this many neighbors.

We claim that B_n does not contain $K_{3,3}$. The geometric idea behind Construction 3.5 (concerning C_4 -free graphs) was to join a point of the finite plane to the points of its “polar” (with respect to the unit circle), and then to use the fact that two lines intersect in at most one point. In contrast, the Brown construction uses the fact that, if points of the Euclidean space \mathbb{E}^3 at distance 1 are joined, then the resulting infinite graph G does not contain $K_{3,3}$. This is easily seen as follows: suppose G does contain $K_{3,3}$. Then the three points of one color class cannot be collinear since no point is equidistant from three collinear points. On the other hand, only two points are equidistant from three points on a circle, and so $K_{3,3}$ cannot occur. There is one problem with this “proof”: in finite fields $\sum_i x_i^2 = 0$ can occur even if not all x_i ’s are 0’s. Therefore in finite geometries, in some cases, a sphere can contain a whole line. So here the geometric language must be translated into the language of analytic geometry, and the right hand side of (3.14) (that is d) must be chosen appropriately.

Theorem 3.21 (Nikiforov, [198]). *For $b \geq a \geq 2$ let $k \in [0, a - 2]$ be an integer. Then*

$$Z(m, n, a, b) \leq (b - k - 1)^{1/a} m n^{1 - (1/a)} + (a - 1) n^{1 + (k/a)} + km.$$

For $k = 0$ we get back Theorem 2.22, and substituting $k = a - 2$ we obtain (3.11). Nikiforov remarks that letting k run from 0 to $a - 2$, we may get the best results for various values of k as the relation of m and n varies, but we still have no constructions to substantiate this. Nikiforov also proves results on the spectral radius.

3.6. Further applications of Algebraic Methods

Most of the constructions providing sufficiently good lower bounds for Bipartite Extremal Graph Problems are coming either from Geometry or from Algebra²⁰. In all these cases the vertices of the graph-construction are “coordinatized” and two vertices are joined if some (usually polynomial) equations are satisfied.²¹

Actually, this motivated Conjecture 1.6 or its weakening: If we use a typical finite geometric construction, then there is a d -dimensional space, where

²⁰The Random Graph Methods are very nice but mostly they are too weak to provide sufficiently sharp lower bounds.

²¹Some of the constructions may seem number theoretic.

each vertex is joined to a t -dimensional subspace. Hence $n = p^d$, the degrees are around $n^{t/d}$, so the construction has around $n^{1+(t/d)}$ edges. The conjecture suggests that there are always such almost extremal constructions.²²

The most important question in this part is if one can find constructions²³ to provide lower bounds where the exponents match the exponents in the upper bounds. Here we shall discuss when do we know the sharpness of the Kővári-T. Sós-Turán upper bound, $\mathbf{ex}(n, K_{a,b}) = O(n^{2-(1/a)})$.

As we have mentioned in Section 1.1, Kollár, Rónyai and T. Szabó [159] gave a construction which was improved by Alon, Rónyai and Szabó [11] (Constructions 3.23 and 3.25 below). The basic idea of their proofs was – at least in our interpretation – the same as that of William G. Brown; however, much more advanced. In the three dimensional Euclidean space \mathbb{E}^3 the Unit Distance Graph contains no $K_{3,3}$. If we change the underlying field to a finite field $GF(q)$ (as Brown did in Construction 3.20) then we obtain a finite graph having $n = q^3$ vertices. The neighborhood of each vertex will have $\approx q^2$ neighbors, and therefore $\approx \frac{1}{2}n^{5/3}$ edges. Now comes the crucial part: despite the fact, that this is highly nontrivial, we could say, that – because of the geometric reason, – this graph contains no $K_{3,3}$ proving the sharpness of (2.6).²⁴

If we wish to extend the above construction to get lower bounds for $\mathbf{ex}(n, K_{a,a})$ and we mechanically try to use unit balls in the a -dimensional space $GF(q)^a$ then several problems occur. We would need that any a of them intersect in at most $a - 1$ points. Then we would be home.

In \mathbb{E}^4 we can choose two orthogonal circles of radii $\frac{1}{\sqrt{2}}$, e.g.,

$$\{(x_1, x_2, 0, 0) : x_1^2 + x_2^2 = 1/2\} \text{ and } \{(0, 0, x_3, x_4) : x_3^2 + x_4^2 = 1/2\},$$

then each point on the first one has distance 1 from each point in the second one. Hence the “Unit Distance Graph” contains $K(\infty, \infty)$. (Similarly, the “Unit Distance Graph” of $GF(q)^4$ contains a $K_{q,q}$.) So everything seems (!) to break down? Not quite, by the Kollár-Rónyai-Szabó construction. Instead

²²Here we have to make some remarks about our “Conjectures”: Many of them have the feature that it is not that interesting if they are true or false: in proving any alternative, we get new, important knowledge about our topics. The first such “Conjecture” was that of Turán on “Diagonal” Ramsey Numbers, that lead to the Erdős Random Graph Approach, see Remark 2.29.

²³Or “random constructions”.

²⁴Actually, as we have already discussed this in Subsection 3.20, the Will Brown’s lower bound also proves this sharpness, only, the lower bound of [11] is a little better.

of the ‘Euclidean metric’ they use a so-called *norm* in the space $GF(q^a)$. Two vectors \mathbf{x} and \mathbf{y} are connected if the norm of their sum is 1; $N(\mathbf{x} + \mathbf{y}) = 1$. (In this context there is not much difference between connecting them this way or take a bipartite graph and connecting the vertices in it if $N(\mathbf{x} - \mathbf{y}) = 1$).

Theorem 3.22 (Kollár, Rónyai, and T. Szabó [159] for $b > a!$, Alon, Rónyai, and Szabó [11] for $b > (a-1)!$). *There exists a $c_a > 0$ such that for $b > (a-1)!$ we have*

$$\mathbf{ex}(n, K_{a,b}) > c_a n^{2-(1/a)}.$$

Below we provide the Kollár-Rónyai-Szabó construction and a short verification. The norm of an element $\mathbf{x} \in GF(q^a)$ is defined as

$$N(\mathbf{x}) := \mathbf{x} \cdot \mathbf{x}^q \cdots \mathbf{x}^{q^{a-1}}.$$

Construction 3.23 (Kollár-Rónyai-T. Szabó [159], the Norm Graph). *The vertices of $G(q, a)$ are the elements $\mathbf{x} \in GF(q^a)$. The elements \mathbf{x} and \mathbf{y} are joined if $N(\mathbf{x} + \mathbf{y}) = 1$.*

We claim that $G(q, a)$ is $K_{b,a}$ -free where $b = a! + 1$. If we have a $K_{b,a} \subseteq G(q, a)$, then fixing – as parameters – the a vertices $\mathbf{y}_1, \dots, \mathbf{y}_a$, we get a equations of the form $N(\mathbf{x} + \mathbf{y}_i) = 1$ with b solutions $\mathbf{x} \in \{\mathbf{x}_1, \dots, \mathbf{x}_b\}$. Then we can use the following result from Algebraic Geometry with $t = a$.

Lemma 3.24. *Let K be a field and $\alpha_{i,j}, \beta_i \in K$ for $1 \leq i, j \leq t$ such that $\alpha_{i_1,j} \neq \alpha_{i_2,j}$ if $i_1 \neq i_2$. Then the system of equation*

$$\begin{aligned} (x_1 - \alpha_{1,1})(x_2 - \alpha_{1,2}) \cdots (x_t - \alpha_{1,t}) &= \beta_1 \\ (x_2 - \alpha_{2,1})(x_2 - \alpha_{2,2}) \cdots (x_t - \alpha_{2,t}) &= \beta_2 \\ &\vdots \\ (x_t - \alpha_{t,1})(x_2 - \alpha_{t,2}) \cdots (x_t - \alpha_{t,t}) &= \beta_t \end{aligned}$$

has at most $t!$ solutions $(x_1, x_2, \dots, x_t) \in K^t$. □

Construction 3.25 (Alon-Rónyai-T. Szabó [11]). *The vertices of the graph $H(q, a)$ are the elements $(x, X) \in GF(q)^* \times GF(q^{a-1})$ and (x, X) and (y, Y) are joined if $N(X + Y) = xy$.*

Here the norm $N(X)$ is defined in $GF(q^{a-1})$ and so it is $X \cdot X^q \cdots X^{q^{a-2}}$. The graph $H(q, a)$ has $(q-1)q^{a-1}$ vertices, it is $q^{a-1} - 1$ regular, and contains no $K_{a,b}$ with $b = (a-1)! + 1$. To show this we use Lemma 3.24 with $t = a-1$ only.

Theorem 3.26 (Ball and Pepe [19]). *The Alon-Rónyai-T. Szabó graph $H(q, 4)$ does not contain $K_{5,5}$. Hence $\mathbf{ex}(n, K_{5,5}) \geq (\frac{1}{2} + o(1))n^{7/4}$.*

This is better than the earlier lower bounds of $\mathbf{ex}(n, K_{a,b})$ for $a = 5$, $5 \leq b \leq 12$, and $a = 6$, $6 \leq b \leq 8$.

Recently, Blagojević, Bukh, and Karasev [24] gave a new algebraic construction to provide lower bounds on $Z(m, n, a, b)$ matching the (3.1) upper bound. Their example is weaker than the Kollár-Rónyai-Szabó in the sense that it only works for $b > (a^2(a + 1))^a$. On the other hand, they give new insight about the limits of the Algebraic Geometric method on which constructions may and which may not work.

We close this section mentioning that Noga Alon has a survey paper in the Handbook of Combinatorics [7] providing ample information on the topics treated here (i.e., applications of algebra in combinatorics).

3.7. The coefficient in the Kővári-T. Sós-Turán bound

Alon, Rónyai and Szabó [11] observed that their Construction 3.25 can be factored with a t -element subgroup $H \subset GF(q)^*$ (when t divides $q - 1$) in the same way as it was done in Construction 3.15. Namely, the vertex set of the new graph $H^t(q, a)$ are the elements $(x, X) \in GF(q)^*/H \times GF(q^{a-1})$ and (x, X) and (y, Y) are joined if $N(X + Y)x^{-1}y^{-1} \in H$. Then the graph $H^t(q, a)$ has $n = (q - 1)q^{a-1}/t$ vertices, its degrees are about q^{a-1} , and it contains no $K_{a,b}$ for $b = (a - 1)!t^{a-1} + 1$. Let $q \rightarrow \infty$. Then also $n \rightarrow \infty$, and we get that for these fixed values of a and b one gets

$$\mathbf{ex}(n, K_{a,b}) \geq (1 - o(1)) \frac{\sqrt[a]{b-1}}{2^{\sqrt[a]{a-1}} (a-1)!} n^{2-(1/a)}.$$

This clearly shows that the order of magnitude of the coefficient in the KST bound (3.1) should be indeed $\sqrt[a]{b-1}$.

Montágh [193] found a clever factorization of the Brown graph (using the spherical symmetry of the balls) thus proving the same result with even a slightly better bound than the bound of Alon, Rónyai and Szabó, for $a = 3$.

3.8. Excluding large complete subgraphs

The following theorem was discovered many times because its connections with Computer Science problems: given a graph G with n vertices, there

exists a decomposition of its edges into complete balanced bipartite graphs K_{a_i, a_i} having altogether $O(n^2/\log n)$ vertices, $\sum_i a_i = O(n^2/\log n)$. Lately, Mubayi and Gy. Turán [195] gave a polynomial algorithms finding such a subgraph partition efficiently. Strictly speaking, this is not a Turán type problem but their result implies, e.g., that there exists a polynomial algorithm to find a $K_{a,a}$ in a graph of $n^2/4$ edges of size $a = \Theta(\log n)$. The bound $O(\log n)$ is the best possible (shown by the random graph).

It is also not very difficult to show that usually the random graph gives the correct order of the Turán number $\mathbf{ex}(n, K_{a,a})$ for $n, a \rightarrow \infty$ simultaneously.

The case when a, b are very large i.e. $a + b = \Omega(n)$ was considered by Griggs, Quyang, and Ho [126], [125]. In this case $Z(m, n, a, b)$ is almost mn so they considered the dual question. Let us mention only one result of this type by Balbuena, García-Vázquez, Marcote, and Valenzuela, who have more papers on this topic.

Theorem 3.27 (see [14], [16] and the references there). $Z(m, n; a, b) = mn - (m + n - a - b + 1)$ if $\max\{m, n\} \leq a + b - 1$.

There is another direction of research, when the ratio of m and n is extreme. Here we only mention a classical result, that it is easy to solve in the case when n is very large compared to m .

Theorem 3.28 (Čulík [55]).

$$Z(m, n, a, b) = (a - 1)n + (b - 1) \binom{m}{a} \quad \text{for} \quad n \geq (b - 1) \binom{m}{a}.$$

4. Excluding Cycles : C_{2k}

To start with, Bondy wrote a long chapter in the Handbook of Combinatorics [30] and also a very nice survey on Erdős and the cycles of graphs [31].

Let \mathcal{C} be a (finite or infinite) set of cycles. The study of $\mathbf{ex}(n, \mathcal{C})$ is especially interesting if \mathcal{C} has a member of even length. However, constructions of dense graphs without some given even cycles is usually very difficult; the examples use polarities of finite geometries (generalized polygons [171]), or Ramanujan graphs [190], [181] or some other families of polynomials [170].

An odd cycle, C_{2k+1} is chromatically critical. Hence a theorem of Simonovits [221] implies that $\mathbf{ex}(n, C_{2k+1}) = \lfloor \frac{1}{4}n^2 \rfloor$ for $n > n_k$ and the only extremal graph is $K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$.

In this Section we concentrate on even cycles C_{2k} .

4.1. Girth and Turán numbers, upper bounds

What is $\mathbf{ex}(n, \{C_3, C_4, \dots, C_{g-1}\})$, the maximum number of edges in a graph with n vertices and girth g ? This problem can be considered in a dual form, what is the least number of vertices $n = n(d, g)$ in a graph of girth g and an average degree at least d ? If we replace ‘average’ with ‘minimum’ δ then a simple argument gives the so-called *Moore bound* for odd girth:

$$|V(G)| = n \geq n_0(\delta, 2k+1) := 1 + \delta \sum_{0 \leq i \leq k-1} (\delta-1)^i. \quad (4.1)$$

Alon, Hoory and Linial [8] showed that (4.1) holds for the average degree, too. Rearranging we have $d_{\text{ave}} < n^{1/k} + 1$, in other words

Theorem 4.1 (Upper bound when the girth is odd).

$$\mathbf{ex}(n, \{C_3, C_4, \dots, C_{2k}\}) < \frac{1}{2}n^{1+(1/k)} + \frac{1}{2}n. \quad (4.2)$$

To prove an upper bound $n^{1+(1/k)}$ is trivial by induction on n . Then (4.2) was improved but with a larger linear additive term.

Theorem 4.2 (Lam and Verstraëte [166], Excluding only even cycles).

$$\mathbf{ex}(n, \{C_4, C_6, \dots, C_{2k}\}) < \frac{1}{2}n^{1+(1/k)} + 2^{k^2}n. \quad (4.3)$$

They also note that for $k = 2, 3, 5$ the n -vertex polarity graphs of generalized $(k+1)$ -gons (defined by Lazebnik, Ustimenko and Woldar [171] described below as Construction 4.27) have $\frac{1}{2}n^{1+(1/k)} + O(n)$ edges and have no even cycles of length at most $2k$.

Corollary 4.3 ([166] and [171] Even girth is 6, 8 or 12). *In case of $2k \in \{4, 6, 10\}$ we have*

$$\mathbf{ex}(n, \{C_4, C_6, \dots, C_{2k}\}) = (1 + o(1))\frac{1}{2}n^{1+(1/k)}. \quad (4.4)$$

On the other hand, Füredi, Naor and Verstraëte [117] showed that if we exclude only C_{2k} , then $\mathbf{ex}(n, C_6) > 0.534n^{4/3}$ (see below as Construction 4.29) and Lazebnik, Ustimenko, and Woldar [169], showed that $\mathbf{ex}(n, C_{10}) > 0.5798n^{6/5}$ (see below as Construction 4.28).

Concerning the Moore bound for even girth we have

$$|V(G)| = n \geq n_0(\delta, 2k+2) := 2 \sum_{0 \leq i \leq k} (\delta - 1)^i. \quad (4.5)$$

Alon, Hoory and Linial [8] showed that (4.5) holds for the average degree, too. Rearranging, we have $d_{\text{ave}} < (n/2)^{1/k} + 1$, in other words

Theorem 4.4 (Upper bound when the girth is even).

$$\text{ex}(n, \{C_3, C_4, \dots, C_{2k+1}\}) < \frac{1}{2^{1+(1/k)}} n^{1+(1/k)} + \frac{1}{2}n. \quad (4.6)$$

This upper bound with a weaker error term was also proved earlier by Erdős and Simonovits [93].

Note that because of Theorems 3.2, 4.21, and 4.23 one can easily show that asymptotic bounds holds in (4.6) for $2k = 4, 6, 10$. The other cases are unsolved.

Theorem 4.5. *For $2k = 4, 6$ and 10 as $n \rightarrow \infty$ we have*

$$\text{ex}(n, \{C_3, C_4, \dots, C_{2k+1}\}) = (1 + o(1)) \frac{1}{2^{1+(1/k)}} n^{1+(1/k)}. \quad (4.7)$$

Moreover, infinitely many exact values are obtained for $2k = 4, 6, 10$: for $n = 2(q^k + q^{k-1} + \dots + q + 1)$,

$$\text{ex}(n, \{C_3, C_4, \dots, C_{2k+1}\}) = (q + 1)(q^k + q^{k-1} + \dots + q + 1) \quad (4.8)$$

whenever q is a power of a prime.

4.2. Excluding a single C_{2k} , upper bounds

Concerning our central problem, Erdős showed that excluding just one even cycle has essentially the same effect as excluding all smaller cycles as well. This is far from trivial! Erdős never published a proof of his result.

Theorem 4.6 (Erdős, The Even Cycle Theorem).

$$\text{ex}(n, C_{2k}) = O(n^{1+(1/k)}). \quad (4.9)$$

The first proof was published by Bondy and Simonovits in the following stronger form.

Theorem 4.7 (Bondy and Simonovits [32]). *Let G_n be a graph with m edges, and let t satisfy $2 \leq t \leq m/100n$ and $tn^{1/t} \leq m/10n$. Then G_n contains a C_{2t} .*

In some sense, this is a “pancyclic theorem”: there is a meta-principle, that if some reasonable conditions ensure the existence of a Hamiltonian cycle, then they ensure the existence of all shorter cycles. Here we go the other direction: if we ensure the existence of a C_{2k} , then we ensure the existence of all longer cycles, up to a natural limit, with the natural parity.

Corollary 4.8. *If G_n has at least $100kn^{n+(1/k)}$ edges, then it contains a C_{2t} , for every $t \in [k, kn^{1/k}]$.*

The Erdős-Bondy-Simonovits upper bound together with earlier known constructions imply that the exponent $1 + (1/k)$ is sharp for C_4 (see, e.g., Theorem 3.4), C_6 , and C_{10} (Theorems 4.22 and 4.24 below).

Corollary 4.9 (The only exact exponents for single cycles).
 $\mathbf{ex}(n, C_4) = \Theta(n^{3/2})$, $\mathbf{ex}(n, C_6) = \Theta(n^{4/3})$, $\mathbf{ex}(n, C_{10}) = \Theta(n^{6/5})$.

The upper end of the interval in Corollary 4.8 is also sharp, apart from the constant 100 take the disjoint union of complete graphs of order $200kn^{1/k}$. We made the following conjecture:

Conjecture 4.10 (Erdős–Simonovits). $\mathbf{ex}(n, C_{2k}) \geq c_k n^{1+(1/k)}$. Moreover,

$$\frac{\mathbf{ex}(n, C_{2k})}{n^{1+(1/k)}}$$

converges to a positive limit.

It is only known for C_4 . A weakening of this conjecture would be the following: Let $\Theta_{k,\ell}$ denote the graph of order $2 + (k-1)\ell$ in which two vertices are joined by ℓ paths of length k .

Conjecture 4.11 (Simonovits). *For each k there is an $\ell = \ell(k)$ for which $\mathbf{ex}(n, \Theta_{k,\ell}) \geq c_k n^{1+(1/k)}$.*

Perhaps the first very annoying unsolved problem on this area is

Conjecture 4.12. $\mathbf{ex}(n, C_8) \geq c_4 n^{5/4}$.

Returning to the Turán number of C_{2k} , the multiplicative constant of the upper bound in the Bondy-Simonovits theorem was improved by Verstraëte [246] from 100 to 8. The best known upper bound today is that of Oleg Pikhurko:

Theorem 4.13 (Pikhurko, [203]).

$$\mathbf{ex}(n, C_{2k}) \leq (k-1)n^{1+(1/k)} + 16(k-1)n.$$

Historical Remark 1.

(a) Pikhurko, in his very nice paper [203] gives a short description of the whole story.

(b) Pikhurko mentions that the Bondy-Simonovits proof gives a constant 20: originally it was stated as 100. It would be, extremely interesting if the upper bound $k-1+o(1)$ for $\mathbf{ex}(n, C_{2k})/n^{1+(1/k)}$ could be improved to $o(k)$.

4.3. Eliminating short cycles, a promising attempt

It was relatively easy to prove the upper bound (4.2) for the number of edges for a graph G_n with girth exceeding $2k$, $e(G_n) = O(n^{1+(1/k)})$. Suppose that G has no C_{2k} . Erdős bipartite subgraph lemma 2.14 states that there is a bipartite subgraph H with $e(H) \geq \frac{1}{2}e(G)$. This way we have eliminated all the odd cycles $C_3, C_5, \dots, C_{2k-1}$ from G . It is a natural to ask whether one can eliminate other short cycles, thus obtaining an easy proof for the Erdős-Bondy-Simonovits upper bound, (4.9).

Problem 4.14. Is it true that there exists a constant $\alpha_{2k} > 0$ such that each C_{2k} -free G_n contains an H_n with $\mathbf{girth}(H_n) > 2k$ and $e(H_n) > \alpha_{2k}e(G_n)$?

The answer is still unknown. The first step was done by E. Györi. The following lemma implies that α_6 exists and it is at least $1/4$.

Lemma 4.15 (Györi [134]). *If G_n is bipartite and it does not contain any C_6 , then it contains an H_n with*

$$e(H_n) \geq \frac{1}{2}e(G_n) + 1,$$

not containing C_4 's either (for $e(G) \geq 2$). This is sharp only for $G_n = K_{2,n-2}$.

We mention two generalizations.

Theorem 4.16 (Füredi, Naor and Verstraete [117]). *Let G be a hexagon-free graph. Then there exists a subgraph of G of girth at least five, containing at least half the edges of G .*

Furthermore, equality holds if and only if G is a union of edge-disjoint complete graphs of order four or five. We got $\alpha_6 = 1/2$.

Theorem 4.17 (Getting rid of C_4 's, Kühn and Osthus [165]). *Every bipartite C_{2k} -free graph G contains a C_4 -free subgraph H with $e(H) \geq e(G)/(k-1)$.*

The factor $1/(k-1)$ is best possible, as the example $K_{k-1, n-k+1}$ shows.

These theorems settle some special cases (namely $\mathcal{L} = \{C_4, C_{2k}\}$) of the following compactness conjecture of Erdős and Simonovits.

Conjecture 4.18 (Compactness. Erdős-Simonovits [93]). *For every finite family of graphs \mathcal{L} (containing bipartite members as well) there exists an $L_0 \in \mathcal{L}$ for which $\mathbf{ex}(n, \mathcal{L}) = O(\mathbf{ex}(n, L_0))$.*

The following result of Kühn and Osthus makes a little step toward solving Problem 4.14 and Conjecture 4.18.

Theorem 4.19 ([165]). *Let $g \geq 4$ be an even integer and let $\ell(g) =: \prod_{1 \leq i \leq g/2} i$. Suppose that $k-1$ is divisible by $\ell(g)$ and G_n is a C_{2k} -free graph. Then G_n contains an H_n with $\mathbf{girth}(H_n) > g$ such that $e(H_n) \geq e(G_n)/2(4k)^{(g-2)/2}$.*

In other words, for some very special values of k 's a C_{2k} -free graph contains a subgraph having a positive fraction of the edges and of girth at least $\Omega(\log k / \log \log k)$.

4.4. A lower bound for C_6 : The Benson Construction

In the preceding section, we asserted that the Erdős theorem on even circuits is sharp for C_4 , C_6 and C_{10} (and is conjectured to be sharp in all cases). For C_4 , the sharpness follows from Construction 3.5. For C_6 , it can be deduced from the Benson construction [22] which we explain below. Note that (about the same time) Singleton [231] described the same graph but his definition was much more complicated.

The points of the d -dimensional finite projective geometry $PG(d, q)$ are the equivalence classes of the nonzero vectors of $GF(q)^{d+1}$ where \mathbf{x} and \mathbf{y} are equivalent if there is a $\gamma \in GF(q)^*$ such that $\mathbf{x} = \gamma\mathbf{y}$. There are $(q^{d+1} - 1)/(q - 1)$ such classes. Then the i -dimensional subplanes are generated by the $(i + 1)$ -dimensional subspaces of the vector space $GF(q)^{d+1}$.

Let

$$A = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

Clearly, A is non-singular. Define the surface S by the equation $\mathbf{x}A\mathbf{x}^T = 0$,

$$S := \{\mathbf{x} \in PG(4, q) : \mathbf{x}A\mathbf{x}^T = 0\}$$

Construction 4.20 (Benson's C_6 -free bipartite graph). *Let \mathcal{L} be the set of lines of $PG(4, q)$ contained entirely in S . The vertex set of the bipartite graph B_q is $S \cup \mathcal{L}$, and $\mathbf{x} \in S$ is joined to $L \in \mathcal{L}$ if $\mathbf{x} \in L$.*

Theorem 4.21 (Benson [22], Singleton [231]). *B_q is a $(q + 1)$ -regular, bipartite, girth 8 graph with $2(q^3 + q^2 + q + 1)$ vertices.*

Corollary 4.22. $\text{ex}(n, C_6) \geq (1 + o(1))(n/2)^{4/3}$.

First, we can see that S does not contain a full 2-dimensional projective plane. We can use the fact that for \mathbf{x} and \mathbf{y} on S , the line \mathbf{xy} consists of the points $\mathbf{z} = a\mathbf{x} + (1 - a)\mathbf{y}$, and lies entirely in S if both $\mathbf{y}A\mathbf{y}^T = 0$ and $\mathbf{x}A\mathbf{y}^T = 0$.

Second, the number of lines from \mathcal{L} containing a given point $\mathbf{x} \in S$ is $q + 1$. Since the number of points on a line is $q + 1$ we immediately get that $|S| = |\mathcal{L}|$.

Furthermore, B_q contains no cycles of length 3, 4, 5 or 7. (For the odd cases this is because it is bipartite, and the existence of a 4-cycle would imply that two points of S are on two distinct lines.) Now suppose that B_q contains a 6-cycle $v_1w_1v_2w_2v_3w_3v_1$. Then S must contain the three lines v_1v_2 , v_2v_3 , and v_3v_1 , and so it must contain the plane $\langle v_1v_2v_3 \rangle$. But this is impossible. If we apply a coordinate transformation T with v_1 , v_2 and v_3 as the first three

base vectors, we get the matrix

$$\begin{pmatrix} 0 & 0 & 0 & ? & ? \\ 0 & 0 & 0 & ? & ? \\ 0 & 0 & 0 & ? & ? \\ ? & ? & ? & ? & ? \\ ? & ? & ? & ? & ? \end{pmatrix}$$

since $v_i A v_j^T = 0$. But then A' cannot be regular, contradicting the regularity of A . Hence B_q cannot contain C_6 either.

All these imply that $|S| = q^3 + q^2 + q + 1$ and that every $\mathbf{x} \in S$, $L \in \mathcal{L}$ if $\mathbf{x} \notin L$ then there exists a unique line $L' \in \mathcal{L}$ such that $\mathbf{x} \in L'$ and $L \cap L' \neq \emptyset$.

In concluding this section, we note that finite geometry constructions can also be used in hypergraph extremal problems (see [39], [40] and [220]).

4.5. Girth 12 graphs by Benson and by Wenger

Theorem 4.23 (Benson [22]). *Let q be an odd prime power. There is a $(q+1)$ -regular, bipartite, girth 12 graph B_q^* with $2(q^5 + q^4 + q^3 + q^2 + q + 1)$ vertices.*

Corollary 4.24. $\text{ex}(n, C_{10}) \geq (1 + o(1))(n/2)^{6/5}$.

One half of the vertex set of B_q^* are the points of the quadric Q_6 in $PG(6, q)$ defined by $x_0^2 + x_1x_{-1} + x_2x_{-2} + x_3x_{-3} = 0$. Its size is exactly $(q^6 - 1)/(q - 1)$. Then we select a set of lines \mathcal{L} contained entirely in Q_6 and covering each point of Q_6 exactly $q + 1$ times. The family \mathcal{L} is selected as follows: If $\mathbf{x} \in Q_6$ and $\mathbf{x}, \mathbf{y} \in L \in \mathcal{L}$ then \mathbf{x} and \mathbf{y} must satisfy the following six bilinear equations:

$$x_0y_i - x_iy_0 + x_{-j}y_{-k} - x_{-k}y_{-j} = 0$$

where (i, j, k) is a cyclic permutation of $(1, 2, 3)$ or $(-1, -2, -3)$.

Construction 4.25. *The bipartite graph B_q^* is defined, as before, by the incidences $\mathbf{x} \in L$.*

Now consider the much simpler example of Wenger.

Construction 4.26 (Wenger [247]). Let p be a prime, $k = 2, 3$ or 5 . $H_k(p)$ is defined as a bipartite graph with two vertex classes \mathbf{A} and \mathbf{B} , where $|\mathbf{A}| = |\mathbf{B}| = p^k$ and the vertices of \mathbf{A} are k -tuples $\mathbf{a} = (a_0, a_1, \dots, a_{k-1}) \in GF(p)^k$ and same for $\mathbf{b} = (b_0, b_1, \dots, b_{k-1}) \in \mathbf{B}$. The vertices \mathbf{a} and \mathbf{b} are joined if

$$b_j \equiv a_j + a_{j+1} \cdot b_{k-1} \pmod{p} \text{ for } j = 0, 1, \dots, k-2.$$

One can see that for every $\mathbf{a} \in \mathbf{A}$ each b_{k-1} determines exactly one $\mathbf{b} \in \mathbf{B}$ joined to it. This easily implies that $G[\mathbf{A}, \mathbf{B}]$ is p -regular, with $n = 2p^k$ vertices and $p^{k+1} = (n/2)^{1+(1/k)}$ edges.

Wenger gives an elegant proof of that $H_2(p)$ has no C_4 , $H_3(p)$ has no C_4 , nor C_6 . Finally, $H_5(p)$ contains no C_4 , C_6 or C_{10} , however, it has many C_8 's.

4.6. Short cycles, C_6 and C_{10}

The densest constructions of $2k$ -cycle-free graphs for certain small values of k arise from the existence of rank two geometries called *generalized d -gons*. These may be defined as rank two geometries whose bipartite incidence graphs are regular graphs of diameter d and girth $2d$. These are known to exist when d is three, four or six. This is the background of the above Constructions 4.20 and 4.25.

Construction 4.27 (Lazebnik, Ustimenko and Woldar [171]). One can use the existence of polarities of the generalized $(k+1)$ -gons to obtain dense $2k$ -cycle-free graphs when $k \in \{2, 3, 5\}$. In particular, for these k 's

$$\mathbf{ex}(n, C_{2k}) \geq \frac{1}{2}n^{1+(1/k)} + O(n) \quad (4.10)$$

for infinitely many n .

In [93], Erdős and Simonovits formulated the following conjecture. For fixed k and $n \rightarrow \infty$, $\mathbf{ex}(n, C_{2k}) = \frac{1}{2}n^{1+(1/k)} + o(n^{1+(1/k)})$. This holds for C_4 (Theorem 3.4), but was disproved first for C_{10} , then for C_6 by the following two examples.

Construction 4.28 (Lazebnik, Ustimenko and Woldar [171]). Consider a bipartite graph $G[A, B]$ of girth exceeding $2k$. Replace each vertex of A by $k-1$ new vertices with the same neighborhood. Then the new graph $G[(k-1)A, B]$ is still C_{2k} -free. In particular, starting with the girth 12 bipartite graph of

Theorem 4.23 (here $k = 5$) one gets a graph of about $5q^5$ vertices and about $4q^6$ edges, implying

$$\mathbf{ex}(n, C_{10}) \geq 4(n/5)^{6/5} > 0.5798n^{6/5} \quad (4.11)$$

for infinitely many n .

Since the C_6 -free graph of Construction 4.27 does not have C_3 and C_4 either, doubling a random subset appropriately, one obtains a denser C_6 -free graph:

Theorem 4.29 (Füredi, Naor and Verstraëte [117]). *For infinitely many n ,*

$$\mathbf{ex}(n, C_6) > \frac{3(\sqrt{5} - 2)}{(\sqrt{5} - 1)^{4/3}} n^{4/3} + O(n) > 0.5338n^{4/3}.$$

They also showed that

Theorem 4.30 (Füredi, Naor and Verstraëte [117]). $\mathbf{ex}(n, C_6) \leq \lambda n^{4/3} + O(n)$, where $\lambda \approx 0.6271$ is the real root of $16\lambda^3 - 4\lambda^2 + \lambda - 3 = 0$.

These theorems give the best known lower and upper bounds for $\mathbf{ex}(n, C_6)$. The proof of Theorem 4.30 requires a statement about hexagon-free bipartite graphs, which is interesting in its own right (see de Caen and Székely [44]). Let $\mathbf{ex}(m, n, C_6)$ be the maximum number of edges amongst all m by n bipartite hexagon-free graphs. Then

Theorem 4.31 (Füredi, Naor and Verstraëte [117]). *Let m, n be positive integers with $n \geq m$. Then*

$$\mathbf{ex}(m, n, C_6) < 2^{1/3}(mn)^{2/3} + 10n.$$

Furthermore, if $n = 2m$ then as n tends to infinity,

$$\mathbf{ex}(m, n, C_6) = \begin{cases} 2^{1/3}(mn)^{2/3} + O(n) & \text{for infinitely many } m \\ 2^{1/3}(mn)^{2/3} - o(n^{4/3}) & \text{for all } m. \end{cases}$$

The lower bound is given by the graph defined in Construction 4.28 starting with the Benson graph (Theorem 4.21, $k = 3$).

4.7. Bipartite hosts with extreme sides

We have already seen two such results concerning the Zarankiewicz number, by Reiman (Theorem 3.2) and Čulík (Theorem 3.28). András Sárközy and Vera Sós formulated the following conjecture ²⁵

Conjecture 4.32.

$$\mathbf{ex}(m, n, C_6) < 2n + c(nm)^{2/3}.$$

A weaker version of this was proved by Gábor N. Sárközy, [212] and later Győri [134] proved a stronger

Theorem 4.33. *There exists a constant $c_k > 0$ for which if $G[A, B]$ is a bipartite graph with color classes A, B , and $|A| = m, |B| = n \geq m^2$, and*

$$e(G[A, B]) \geq (k-1)n + c_k m^2,$$

then $G[A, B] \supset C_{2k}$.

This means that for $n > m^2$ the extremal number becomes linear. For more recent results see, e.g., Balbuena, García-Vázquez, Marcote, and Valenzuela [15]. Later Győri [135] showed that $c_3 = 1/8$, proving

$$\mathbf{ex}(m, n, C_6) \leq 2n + \frac{1}{8}m^2,$$

for $n, m > 100, n \geq m^2/16$ and here equality holds if m is a multiple of 4.

4.8. The effect of odd cycles

Let \mathcal{L} be a set of graphs and let $\mathbf{ex}_{\text{bip}}(n, \mathcal{L})$ denote the *bipartite Turán number* of \mathcal{L} , the size of the largest \mathcal{L} -free bipartite graph on n vertices.

Theorem 4.34 (Erdős and Simonovits [93]).

$$\mathbf{ex}(n, \{C_4, C_5\}) = (1 + o(1))\mathbf{ex}_{\text{bip}}(n, C_4) = (1 + o(1))(n/2)^{3/2}.$$

They also **conjecture** that the same holds for $\{C_3, C_4\}$ (i.e., for the girth problem) but this is still unsolved. Then, they make the following bold conjecture.

²⁵A weaker version of this conjecture was formulated by Erdős several years earlier.

Conjecture 4.35 (Erdős and Simonovits [93] on the effect of odd cycles).
 Let $\mathcal{C}_{2\ell+1}^{\text{odd}}$ denote the set of odd cycles $\{C_3, C_5, \dots, C_{2\ell+1}\}$. For any family \mathcal{L} consisting of bipartite graphs there exists an odd integer $2\ell + 1$ such that $\mathbf{ex}(n, \mathcal{L} \cup \mathcal{C}_{2\ell+1}^{\text{odd}}) \approx \mathbf{ex}_{\text{bip}}(n, \mathcal{L})$.

This conjecture was verified in a few cases by extending and sharpening Theorem 4.34 as follows.

Theorem 4.36 (Keevash, Sudakov and Verstraëte [157]). Let $\mathcal{C}_{2k}^{\text{even}}$ denote the set of even cycles $\{C_4, C_6, \dots, C_{2k}\}$. Suppose that $2k \in \{4, 6, 10\}$ and suppose that $2\ell + 1 > 2k$. Then

$$\mathbf{ex}(n, \mathcal{C}_{2k}^{\text{even}}, C_{2\ell+1}) = (1 + o(1))\mathbf{ex}_{\text{bip}}(n, \mathcal{C}_{2k}^{\text{even}}) \sim (n/2)^{1+(1/k)}.$$

They even proved a stability result (when $n \rightarrow \infty$) and, using it, an exact version: If $2k \in \{4, 6, 10\}$ and $2\ell + 1 \geq 5, 15,$ or 23 , respectively, and $n = 2(q^k + q^{k-1} + \dots + q + 1)$ then for $n > n_{2\ell+1}$ we have

$$\mathbf{ex}(n, \mathcal{C}_{2k}^{\text{even}} \cup C_{2\ell+1}) \leq (q + 1)n$$

and here equality holds only if there is a generalized $(k + 1)$ -gon of order q .

In a more recent work Allen, Keevash, Sudakov and Verstraëte [5] verified the stronger form of the Erdős-Simonovits conjecture proving that for any fixed $2\ell + 1 \geq 5$ one has $\mathbf{ex}(n, \{K_{2,t}, C_{2\ell+1}\}) \sim \mathbf{ex}_{\text{bip}}(n, K_{2,t})$ and $\mathbf{ex}(n, \{K_{3,3}, C_{2\ell+1}\}) \sim \mathbf{ex}_{\text{bip}}(n, K_{3,3})$. They also show

$$\mathbf{ex}(n, \{K_{2,t}, B_t, C_{2\ell+1}\}) \sim \mathbf{ex}_{\text{bip}}(n, \{K_{2,t}, B_t\}) \sim (n/2)^{3/2}$$

for any fixed $t \geq 2$ and $2\ell + 1 \geq 9$, where B_t is a “book” of t C_4 ’s sharing an edge: it has $2t + 2$ vertices and $3t + 1$ edges. Their main tool is the smoothness of the corresponding Turán number’s and the sparse regularity lemma of A. Scott [213].

On the other hand, for any $t \geq 1$ and prime $q > 2^{t^4}$, they construct $(t + 2)$ -partite graphs $G_{q,t}$ with no triangle or $K_{2,2t+1}$ having $n = (t + 2)q^2$ vertices and $\binom{t+2}{2}q^2(q - 1)$ edges. This implies

$$\mathbf{ex}(n, \{K_{2,2t+1}, C_3\}) \geq (1 + o(1)) \frac{t + 1}{\sqrt{t + 2}} n^{3/2}. \quad (4.12)$$

So, using $\mathbf{ex}_{\text{bip}}(n, K_{2t+1}) \sim \sqrt{tn}^{3/2}$, which follows easily from (3.7) and (3.9), they obtain

$$\liminf_{n \rightarrow \infty} \frac{\mathbf{ex}(n, \{K_{2,2t+1}, C_3\})}{\mathbf{ex}_{\text{bip}}(n, K_{2,2t+1})} \geq \frac{t + 1}{\sqrt{t(t + 2)}} > 1. \quad (4.13)$$

In particular the ratio is $2/\sqrt{3} + o(1)$ for $K_{2,3}$. We explain their construction yielding (4.12) only for $t = 1$.

Construction 4.37 (Allen, Keevash, Sudakov and Verstraëte [5]). *Let $q \equiv 2 \pmod{3}$ be a prime. Let G^q be a three-partite graph with parts A_1, A_2 and A_3 which are copies of $GF(q) \times GF(q)$. Join $(x_1, x_2) \in A_i$ to $(y_1, y_2) \in A_{i+1}$ if*

$$(y_1, y_2) = (x_1, x_2) + (a, a^2)$$

for some $a \in GF(q)$, $a \neq 0$.

The obtained graph is $K_{2,3}$ and C_3 -free, and has $n = 3q^2$ vertices and $n^{3/2}/\sqrt{3}$ edges. This yields the ratio $2/\sqrt{3} + o(1)$ for $K_{2,3}$ in (4.13). They believe that Erdős' Conjecture 1.8 is false:

Conjecture 4.38 ([5]).

$$\liminf_{n \rightarrow \infty} \frac{\mathbf{ex}(n, \{C_3, C_4\})}{\mathbf{ex}_{\text{bip}}(n, C_4)} > 1.$$

4.9. Large girth: Ramanujan graphs

Until this point we were fixing the excluded subgraphs. However, there is a subcategory of extremal graph problems, which we could also call “Parametrized Extremal Graph Problems”. Instead of defining them we give an almost trivial but important example: Horst Sachs and Erdős [87] reformulated the Moore bounds (4.1) (4.5), in a slightly different form.

Theorem 4.39. *If the minimum degree of G_n , $d := d_{\min}(G_n) > 2$ then G_n contains a C_ℓ with*

$$\ell < \frac{2 \log n}{\log(d-1)}. \quad (4.14)$$

Here we arrived at an area where some constructions (for lower bounds) were needed, and the lower bounds were easily obtained by probabilistic arguments; however they were very difficult to obtain them in a constructive way. Instead of going into details, we mention a result of Margulis [188] that (4.14) is sharp up to a constant: there are – not too complicated – Cayley graphs of constant (even) degrees d and girth at least $c \log_{d-1} n$. Here – surprisingly, Margulis' construction is better than the random graph and a construction of Imrich yields an even better constant c :

Theorem 4.40 (Imrich [147]). *For every integer $d > 2$ one can (effectively) construct infinitely many d -regular Cayley graphs X_n with*

$$\mathbf{girth}(X_n) > 0.4801 \frac{\log n}{\log(d-1)} - 2.$$

The next step in this area was a much deeper and more important results of Margulis [190, 189], Lubotzky, Phillips and Sarnak, [181] on the Expander graphs, that are eigenvalue-extremal. In this sense the Margulis-Lubotzky-Phillips-Sarnak graph is very nice. There is only one problem with it. While defining these graphs is non-trivial, but not extremely complicated, to verify their extremal properties requires deep mathematical tools. Below we give a very compressed description of it.

Definition 4.41. Given a connected k -regular graph X , we denote by $\lambda(X)$ the largest of the absolute values of eigenvalues of the adjacency matrix of X , different from k . An n -vertex k -regular graph $X_{n,k}$ is a *Ramanujan graph* if $\lambda(X_{n,k}) \leq 2\sqrt{k-1}$.

Remark 4.42. In case of k -regular graphs, the largest absolute values of the eigenvalues is k . The bipartite graphs have the property that if λ_i is eigenvalue, then $-\lambda_i$ is also an eigenvalue. By the Alon-Boppana inequality, (see Proposition 4.2 of [181])

$$\liminf_{n \rightarrow \infty} \lambda(X_{n,k}) = 2\sqrt{k-1}.$$

Ramanujan graphs are important because they are expander graphs, which are extremely important in Theoretical Computer Science.

There are quite a few cases, where – instead of using “random graph constructions” one tries to use Cayley Graphs. *Cayley graphs* are graphs whose vertices are the elements of some group \mathcal{G} and the edges are the pairs $(g, \alpha_i g)$, where $g \in \mathcal{G}$ and $\alpha_1, \dots, \alpha_k$ come from a generator system of \mathcal{G} . If we look for a digraph, then this is a correct definition. However, if we are looking for ordinary graphs, then we have to assume that $S := \{\alpha_1, \dots, \alpha_k\}$ is closed under taking the inverse: if $\alpha \in S$ then $\alpha^{-1} \in S$ as well. Now, if we choose \mathcal{G} and S appropriately, then the obtained graph will provide us with nice constructions; mainly, because it behaves as if it were a random graph, or, occasionally, even better.

Construction 4.43 ([181]). *Let p and q be unequal primes congruent to 1 mod 4. The Ramanujan graphs $X^{p,q}$ of [181] are $p+1$ -regular Cayley graphs²⁶ of the group $\mathbf{PSL}(2, \mathbb{Z}/q\mathbb{Z})$: $p+1$ generators of the group are fixed, which are obtained from the solutions of*

$$p = a^2 + b^2 + c^2 + d^2, \text{ where } a > 0 \text{ is odd and } b, c, d \text{ are even.} \quad (4.15)$$

The number of solutions of (4.15) is connected to the famous Ramanujan conjecture, which is still open. However, good approximations are known, by Eichler and Igusa, enough for the purposes of [181]. Originally most of the authors were interested in the eigenvalue properties (spectral gap) of these graphs, that are also strongly connected to them being expander graphs (see Alon, [6], Alon–Milman [10]).

From here on, $X_{n,k} = X^{p,q}$ is a special sequence of Ramanujan graphs, which is non-bipartite if the Jacobi symbol $\left(\frac{q}{p}\right) = 1$; then it has $n = (q^3 - q)/2$ vertices.

Theorem 4.44. *For $k = p+1$, $X^{p,q}$ is k -regular, its eigenvalues are $\lambda = \pm k$ or $|\lambda| \leq 2\sqrt{k-1}$.*

This property is optimal and leads to the best known explicit expander graphs. Alon turned the attention of the authors to that these graphs satisfy a number of extremal combinatorial properties.

Theorem 4.45 (Observation of Alon). *The girth of $X_{n,k}$ is asymptotically $\geq \frac{4}{3} \frac{\log n}{\log(k-1)}$.*

This gives larger girth than what was previously known by explicit or non-explicit constructions. Also, it is one of the “cleanest” way to define graphs with large girth and high chromatic number:

Theorem 4.46 ([181]). *If $X_{n,k}$ is a non-bipartite Ramanujan graph, then its independence number and chromatic number satisfy*

$$\alpha(X_{n,k}) \leq \frac{2\sqrt{k-1}}{k}n \quad \text{and} \quad \chi(X_{n,k}) \geq \frac{k}{2\sqrt{k-1}}.$$

For a more informative description of these and many other related areas see the survey of Alon in the Handbook [7].

²⁶There are two of them, a bipartite and a non-bipartite, we forget the bipartite one.

4.10. The girth problem: the Lazebnik-Ustimenko approach

After 20 years Theorem 4.47 still yields the best known lower bound for the girth problem: Lazebnik, Ustimenko and Woldar's work [170] gives a slight improvement (an $O(1)$ in the denominator of the exponent) to what we can get from the Ramanujan' graphs.

Theorem 4.47 ([170]). $\text{ex}(n, \{C_3, C_4, \dots, C_{2k+1}\}) = \Omega(n \cdot n^{2/(3k-3+\varepsilon)})$ where $k \geq 2$ is fixed, $\varepsilon = 0$ if k is odd, $\varepsilon = 1$ if k is even and $n \rightarrow \infty$.

We have seen basically two approaches on how to *construct* graphs with high girth. One was the use of Finite Geometries, and the other the use of Cayley Graphs of some matrix groups (Ramanujan graphs). There is (at least) one further important approach to this question which we find in the works of Lazebnik and Ustimenko and later Lazebnik, Ustimenko and Woldar.

Remark 4.48 (History). In this survey many important areas had to be skipped. One of them is the family of Lazebnik-Ustimenko type algebraic constructions. This family of constructions is much more flexible than many earlier ones, and provides a lot of new constructions in extremal graph theory, in Ramsey type problems, for graphs and hypergraphs as well. The first results were achieved by Lazebnik and Ustimenko [168]. Lazebnik and his coworkers created a school in this area. The reader is referred here to [167].

The main feature of this approach can be described (perhaps slightly cheating) as follows. We take a set R (a finite or infinite ring or field), its d^{th} power, and a sequence of polynomials f_2, \dots, f_d . Define a bipartite graph, where the colour classes A and B consist of vectors (a_1, \dots, a_d) and (b_1, \dots, b_d) that are joined if

$$\begin{aligned} a_2 + b_2 &= f_2(a_1, b_1) \\ a_3 + b_3 &= f_3(a_1, b_1, a_2, b_2) \\ &\dots \\ a_d + b_d &= f_d(a_1, b_1, \dots, a_d, b_d). \end{aligned}$$

We may also identify A and B to get non-bipartite graphs as well. In general, either we get digraphs, or some symmetry conditions are assumed on

the functions f_i , ensuring that if (a_1, \dots, a_d) is joined to (b_1, \dots, b_d) , then (b_1, \dots, b_d) and (a_1, \dots, a_d) are joined as well. Yet, it is not an easy area to describe it on a few pages: this is why we basically skip it. Perhaps the more interested reader should look at [172].

4.11. Cycle length distribution

As a measure of the density of the cycle lengths in a graph G , Erdős introduced the number $L(G)$, the sum of the reciprocals of the distinct cycle lengths of G . The following beautiful theorem, due to Gyárfás, Komlós and Szemerédi, proves a conjecture of Erdős and Hajnal, asserting that in some sense the complete graph or the complete bipartite graph are the densest concerning cycle lengths:

Theorem 4.49 ([131]). *There exists a positive constant $c > 0$ such that if $d_{\min}(G) \geq k$, then for the sum of the reciprocals of the cycle lengths ℓ_i of G we have*

$$L(G) = \sum \frac{1}{\ell_i} > c \log k.$$

The union of complete graphs K_{k+1} or bipartite graphs $K_{k,m}$ (where $m \geq k$) show that this lower bound is sharp.

Generalizing a theorem of Bondy and Vince [33], Gengua Fan proved several nice results on the distribution of cycle lengths. We mention only one of them.

Theorem 4.50 (G. Fan [96]). *Let xy be an edge in a 2-connected graph G , k be a positive integer and suppose that all the vertices of G but x and y have degrees at least $3k$. Then xy is contained in $k+1$ cycles C^0, C^1, \dots, C^k , such that $k+1 < |E(C^0)| < |E(C^1)| < \dots < |E(C^k)|$, $|E(C^i)| - |E(C^{i-1})| = 2$ for $i = 1, \dots, k-1$ and $1 \leq |E(C^k)| - |E(C^{k-1})| \leq 2$.*

A related result concerning k odd cycle lengths can be found in Gyárfás [130].

Next we recall a conjecture of Burr and Erdős.

Conjecture 4.51 (Burr and Erdős). *For every odd integer $k > 0$, and every integer ℓ , there exists a c_k such that if $e(G_n) > c_k n$, then some $m \equiv \ell \pmod{k}$, we have $C_m \subseteq G_n$.*

This was proved by Bollobás [25] with $c_k \leq ((k+1)^k - 1)/k$. Häggkvist and Scott ([137], [138]) decreased c_k and extended the Bollobás result, proving that every graph G_n with minimum degree at least $300k^2$ contains k cycles of consecutive even lengths. Soon after, the right order of magnitude of c_k was established.

Theorem 4.52 (Verstraëte [246]). *Let G_n be a graph with $e(G_n) \geq 4kn$. Then there are cycles of k consecutive even lengths in G_n .*

We close this part with the following theorem:

Theorem 4.53 (Sudakov, Verstraëte [234]). *Let $\text{girth}(G_n) = g$ be fixed and $d = 2e(G_n)/n$. Let $\mathcal{C}(G)$ denote the set of cycle-lengths in G . Then $\mathcal{C}(G_n)$ contains at least $\Omega(d^{\lfloor (g-1)/2 \rfloor})$ consecutive even integers, as $d \rightarrow \infty$.*

5. Paths and long cycles

In this section we shall describe results connected with $\text{ex}(n, P_k)$, $\text{ex}(n, \mathcal{C}_{\geq k})$, (where the cycles of at least k vertices are excluded). This problem was proposed by Turán and the (asymptotic) answer were given by Erdős–Gallai.

5.1. Excluding long cycles

Theorem 5.1 (Erdős and Gallai [80]). *Let G_n be a graph with more than $\frac{1}{2}(k-1)(n-1)$ edges, $k \geq 3$. Then G_n contains a cycle of length at least k . This bound is the best possible if $n-1$ is divisible by $k-2$.*

A matching lower bound $\frac{1}{2}(k-1)n - O(k^2)$ can be obtained gluing together complete graphs of sizes at most $k-1$. If k is odd, then there are nearly extremal graphs having a completely different structure. Namely, one can take a complete bipartite graph with partite sets A and B of sizes $|A| = \frac{k-1}{2}$ and $|B| = n - \frac{k-1}{2}$ and add all edges in A , too.

The exact value was determined by Woodall [249] and independently and at the same time by Kopylov [162].

Theorem 5.2 ([162], [249]). *Let $n = m(k-2) + r$, where $1 \leq r \leq k-2$, $k \geq 3$, $m \geq 1$ integers. If*

$$e(G_n) > m \binom{k-1}{2} + \binom{r}{2},$$

then G_n contains a cycle of length at least k , and this bound is the best possible:

$$\mathbf{ex}(n, \mathcal{C}_{\geq k}) = \frac{1}{2}(k-1)n - \frac{1}{2}r(k-r). \quad (5.1)$$

Caccetta and Vijayan [43] gave an alternative proof of the result. We need a definition.

Construction 5.3. Let $H_{n,k,s}$ be an n -vertex graph consisting of a complete graph K_{k-s} on the set $A \cup B$, $|A| = k - 2s$, $|B| = s$ and a complete bipartite graph $K_{s, n-(k-s)}$ with parts B and C where A, B and C form a partition of $V(H)$ (hence $|C| = n - (k - s)$ and $n \geq k$, $(k - 1)/2 \geq s \geq 1$).

The graph H contains no cycle of size k or larger and for $s \geq 2$ it is 2-connected. Denote its size by $h(n, k, s)$.

They all ([162], [249], [43]) characterized the structure of the extremal graphs in Theorem 5.2. Namely either

- the blocks of G_n are m complete graphs K_{k-1} and a K_r , or
- k is odd, $r = (k + 1)/2$ or $(k - 1)/2$ and q of the blocks of G_n are K_{k-1} 's and a copy of a $H_{n-q(k-2), k, (k-1)/2}$.

The strongest result on the field is due to Kopylov who also investigated 2-connected graphs.

Theorem 5.4 (Kopylov [162]). *Suppose that $n \geq k \geq 5$ and the 2-connected graph G_n contains no cycles of length of k or larger. Then*

$$e(G_n) \leq \max\{h(n, k, 2), h(n, k, \lfloor \frac{1}{2}(k-1) \rfloor)\}$$

and this bound is the best possible.

Moreover, only the graphs $H_{n,k,s}$ could be extremal, $s \in \{2, \lfloor (k-1)/2 \rfloor\}$.

This theorem was also conjectured by Woodall [249] and he also proved it for $n \geq (3k - 5)/2$. It was also reproved much later in [97].

5.2. Excluding P_k

One of the oldest problems is the question of determining $\mathbf{ex}(n, P_k)$.

Theorem 5.5 (Erdős and Gallai [80]). *If G_n is a graph containing no P_k , ($k \geq 2$), then*

$$e(G_n) \leq \frac{k-2}{2}n$$

with equality if and only if $k - 1$ divides n and all connected components of G are complete graphs on $k - 1$ vertices.

Consider the n -vertex graph G_n which is the union of $\lfloor n/(k - 1) \rfloor$ vertex-disjoint K_{k-1} and a K_r ($0 \leq r \leq k - 2$). If T_k is any connected k -vertex graph, then $T_k \not\subseteq G_n$. Hence

$$\mathbf{ex}(n, T_k) \geq \frac{k - 2}{2}n - \frac{1}{8}k^2. \tag{5.2}$$

In particular,

$$\mathbf{ex}(n, P_k) \geq \frac{k - 2}{2}n - \frac{1}{8}k^2. \tag{5.3}$$




Figure 2: Potential extremal graphs

If k is even, then there are nearly extremal graphs having a completely different structure. Namely, one can take a complete bipartite graph with partite sets A and B of sizes $|A| = \frac{k-2}{2}$ and $|B| = n - \frac{k-2}{2}$ and add all edges in A , too (Fig. 2). Faudree and Schelp [98] proved that the extremal graph for P_k can indeed be obtained in this way for all n and k . They needed this to prove some Ramsey theorems on paths. The variety of extremal graphs makes the solution difficult.

Theorem 5.6 (Faudree and Schelp [98] and independently Kopylov [162]).

Let $n \equiv r \pmod{k - 1}$, $0 \leq r < k - 1$, $k \geq 2$. Then

$$\mathbf{ex}(n, P_k) = \frac{1}{2}(k - 2)n - \frac{1}{2}r(k - 1 - r). \tag{5.4}$$

Faudree and Schelp also described the extremal graphs which are either
 — vertex disjoint unions of m complete graphs K_{k-1} and a K_r , or
 — k is even and $r = k/2$ or $k/2 - 1$ and another extremal graphs can be obtained by taking a vertex disjoint union of t copies of K_{k-1} ($0 \leq t < m$) and a copy of $H(n - t(k - 1), k/2, k/2 - 1)$.

Theorem 5.7 (Kopylov [162]). *Let G_n be a connected graph containing no P_k , ($k \geq 4$) and $n \geq k$. Then*

$$e(G) \leq \max\{h(n, k-1, 1), h(n, k-1, \lfloor \frac{1}{2}(k-2) \rfloor)\}$$

and this bound is the best possible.

Moreover, only the graphs $H_{n, k-1, s}$ could be extremal, $s \in \{1, \lfloor (k-2)/2 \rfloor\}$.

Balister, Györi, Lehel and Schelp [18] also provided the extremal structures.

5.3. Proof ideas

Let $\mathbf{ex}_{\text{con}}(n, P_k)$ be the maximum number of edges in connected, n -vertex, P_k -free graphs, and let $\mathbf{ex}_{2\text{-con}}(n, C_{\geq k})$ denote the maximum number of edges in 2-connected, n -vertex, $C_{\geq k}$ -free graphs. Determining these functions give upper bounds for $\mathbf{ex}(n, P_k)$ and $\mathbf{ex}(n, C_{\geq k})$.

Indeed, every P_k -free graph is a vertex disjoint union of P_k -free components, we have

$$\mathbf{ex}(n, P_k) = \max_{\sum n_i = n, n_i \geq 1} \sum \mathbf{ex}_{\text{con}}(n_i, P_k).$$

Similarly, a maximal $C_{\geq k}$ -free graph is connected and every connected graph is a cactus-like union of 2-connected blocks (and edges) so we have

$$\mathbf{ex}(n, C_{\geq k}) = \max_{\sum (n_i - 1) = n - 1, n_i \geq 2} \sum \mathbf{ex}_{2\text{-con}}(n_i, C_{\geq k}), \quad (5.5)$$

where we define $\mathbf{ex}_{2\text{-con}}(2, C_{\geq k}) = 1$.

Let G be a connected, n -vertex, P_k -free graph. Add a new vertex to it and join to all other vertices. We obtain G_{n+1} with $e(G_{n+1}) = e(G) + n$. This new graph has no cycle of length exceeding k and its connectivity is one larger than that of G_n . We obtain

$$\mathbf{ex}(n, P_k) + n \leq \mathbf{ex}(n+1, C_{\geq k+1}) \quad (5.6)$$

and

$$\mathbf{ex}_{\text{con}}(n, P_k) + n \leq \mathbf{ex}_{2\text{-con}}(n+1, C_{\geq k+1}). \quad (5.7)$$

So Theorem 5.1 and (5.6) imply Theorem 5.5. Similarly, Theorem 5.2 and (5.6) imply Theorem 5.6.

The upper bounds for $\mathbf{ex}_{2\text{-con}}(n, C_{\geq k+1})$ yield upper bounds for $\mathbf{ex}_{\text{con}}(n, P_k)$. (Actually, (5.7) and Theorem 5.4 lead to the solution of $\mathbf{ex}_{\text{con}}(n, P_k)$, Theorem 5.7).

Again Theorem 5.4 and (5.5) lead to Theorem 5.2 which is obviously stronger than Theorem 5.1.

Finally, the proof of Theorem 5.4 uses induction on n and k , by deleting small degree vertices, contracting edges, and finally applying Pósa's theorem on Hamiltonian graphs.

5.4. Generalizations

In a recent work Lidický, Hong Liu and Cory Palmer [174] determined the exact Turán number (and the unique extremal graph) when the forbidden graph L is a *linear forest*, each component is a path. They also considered *star-forests*.

Gyárfás, Rousseau, and Schelp [132] determined $\mathbf{ex}(K(m, n), P_k)$ for all m, n, k . Their formula and proof are rather involved, they distinguish 10 subcases.

6. Excluding trees

Here we shall discuss two extremal problems on trees: the Erdős-Sós conjecture and the Loeb-Komlós-Sós conjecture.

6.1. Erdős-Sós conjecture

We have already discussed the Erdős-Gallai theorems. Since the extremal numbers for P_k and for the star $K_{1, k-1}$ are roughly the same, this led Erdős and T. Sós to the following famous conjecture.

Conjecture 6.1 (Erdős-Sós [63]). *Let T_k be an arbitrarily fixed k -vertex tree. If a graph G_n contains no T_k , then*

$$e(G_n) \leq \frac{1}{2}(k-2)n. \quad (6.1)$$

As we have seen – by (5.2) – the disjoint union of complete graphs K_{k-1} shows that $\mathbf{ex}(n, T_k) \geq \frac{1}{2}(k-2)n - \frac{1}{8}k^2$. Though several partial cases were

settled, the upper bound was unknown until Ajtai, Komlós, Simonovits and Szemerédi proved:

Theorem 6.2 (Main Theorem, Sharp [1, 2, 3]). *There exists an integer k_0 such that if $k > k_0$ and T_k is an arbitrarily fixed k -vertex tree, and the graph G_n contains no T_k , then*

$$e(G_n) \leq \frac{1}{2}(k-2)n. \quad (6.2)$$

Below we list a few subcases where this conjecture is verified, but we do not try to give a complete list.

Theorem 6.3 (Sidorenko [215]). *If T_k has a vertex x connected to at least $k/2$ vertices of degree 1 (i.e., leaves) then the Erdős-Sós conjecture holds for this T_k .*

Theorem 6.4 (McLennan [182]). *If the diameter of T_k is at most 4, then the Erdős-Sós conjecture holds for this T_k .*

Dobson (and coauthors) have several results in this area, under some strong condition of sparsity. We mention only the Brandt-Dobson theorem [34], or Sacle and Wozniak, [251], [211].

6.2. Sketch of the proof of Theorem 6.2

We are given a T_k , and a G_n violating (6.2). We wish to embed T_k into G_n ($T_k \hookrightarrow G_n$). The proof is very involved and will be given in three rather long papers. The following weakening plays a central role.

Theorem 6.5 (η -weakening [1]). *For any (small) constant $\eta > 0$ there exist a $k_0(\eta)$ such that for $n \geq k > k_0(\eta)$, if*

$$e(G_n) > \frac{1}{2}(k-2)n + \eta kn, \quad (6.3)$$

then each k -vertex tree T_k is contained in G_n .

(a) First, in [1] we prove this theorem. If, in addition, we assume that G_n is dense: for some $c > 0$, $k > cn$, then we can apply the Szemerédi Regularity Lemma [235]. The proof of this theorem follows basically the line which was later used to prove the Loeb Conjecture, by Ajtai, Komlós and Szemerédi [4]

and later by Yi Zhao [255]. Also it was used in the Komlós-Sós conjecture by Diana Piguet and Maya Stein [201], Cooley [54], Hladký and Piguet [144], in stronger and stronger form, and now the publication of that proof is almost finished by Hladký, Komlós, Simonovits, Stein, and Szemerédi [143].

(b) In the second part, [2] we prove several theorems asserting that under some very special conditions $T_k \subseteq G_n$. Some of these steps are “stability arguments”.

Analyzing the proof of Theorem 6.5, shows that either we can gain at some points, in some of the estimates ηkn edges, and therefore Theorem 6.5 (more precisely, its slightly modified proof) implies the sharp version, Theorem 6.2, or else G_n must have a very special structure: it contains a smaller copy of the conjectured extremal graphs: for some $m \approx k$,

(b₁) either it contains a G_m which is almost a K_m ;

(b₂) or a G_m which is almost a $K(m/2 - \varepsilon m, m + \delta m)$.

(c) In both cases, if many edges connect $G_n - G_m$ to G_m , then we can embed T_k into G_n , embedding a smaller part of T_k outside of G_m , a larger part in the dense G_m , concluding that $T_k \hookrightarrow G_n$.

(d) If, on the other hand, we have found such a “mini-almost-extremal” $G_m \subseteq G_n$, but $e(G_m, G_n - G_m)$ is “small”, then we prove that

$$e(G_n - G_m) > \frac{1}{2}(n - m)(k - 2).$$

Hence we may forget the larger G_n : replace it by the smaller $G_n - G_m$. (In other words, we can apply induction on n .)

(e) The real difficulty comes when we have sparse graphs: $e(G_n) = o(n^2)$. Then we partition $V(G_n)$ into three parts: \mathbb{C} contains the vertices of high degrees, \mathbb{B} contains a part of $V(G_n)$ not containing dense subgraphs, and therefore behaving in a pseudo-random way, and \mathbb{A} behaves very similarly to the graphs we have in the dense cases.

How do we handle the dense case?

(i) Applying the Regularity Lemma to G_n , we get a so called Cluster Graph H_ν . If this cluster graph has an (almost)-1-factor, then we can relatively easily embed T_k into G_n , using the extra ηkn edges of (6.3).

(ii) Next we extend this case to a more general situation, when G_n contains a so called Generalized 1-factor. We can prove the η -weakening in this case as well.

(iii) If the Cluster Graph H_ν does not contain an almost-1-factor, then we apply the Gallai-Edmonds structure-theorem (on graphs without 1-factors)

to H_ν . In this case we can either embed T_k into G_n directly, or reduce this case to Case (ii) above. Case (iii) is a very important subcase, with 3-4 subsubcases (depending on, how do we count them). Some of them go back to Case (ii) and in some others we directly (pseudo-greedily) embed T_k into G_n .

6.3. Komlós-Sós conjecture on median degree

The Komlós-Sós conjecture was already formulated in Section 1.4. This is a generalization of the Loebel conjecture:

Conjecture 6.6 (Loebel–Komlós–Sós Conjecture [79]). *If G_n has at least $n/2$ vertices of degree at least $k - 1$, then G_n contains all the k -vertex trees T_k .*

The authors of [143] plan to write up the sharp version as well, which asserts the following.

Theorem 6.7. *If k is sufficiently large, then the Loebel–Komlós–Sós conjecture is true.*

Remarks 6.8. (a) The Loebel conjecture originates from a paper of Erdős, Füredi, Loebel, and T. Sós, on the discrepancy of trees [79].

(b) Pósa’s theorem on the existence of Hamiltonian cycles also is – in some sense – a theorem asserting that if G has many vertices of sufficiently high degree, then it is Hamiltonian. There were earlier cases, when Woodall [250], proved an Erdős–Gallai type theorem on cycles, using the condition that there are many vertices of high degree. Also, Erdős, Faudree, Schelp and Simonovits – trying to prove some Ramsey type theorems, – found a similar statement [78], but not for all the trees, only for the paths, and they proved there an almost sharp theorem. Their sharp conjecture was later proved by Hao Li.

(c) There were many important steps to reach the theorem above. We should mention here Ajtai–Komlós–Szemerédi, [4], then Yi Zhao [255], next Piguet and Stein [201], [202], Cooley [54], Hladký and Piguet [144], and many others.

7. More complex excluded subgraphs

In this section we present three theorems, each leading to a reduction method to prove new results from old estimates. Still there is no general theory to

determine the bipartite Turán numbers.


The three results we pick are the Erdős-Simonovits cube theorem (Theorem 7.1), $\mathbf{ex}(n, Q_8) = O(n^{8/5})$ which led to the Erdős-Simonovits reduction, the Faudree-Simonovits theorem (Theorem 7.8) concerning Theta graphs, a generalization of the Erdős-Bondy-Simonovits theorem, $\mathbf{ex}(n, C_{2k}) = O(n^{1+(1/k)})$, and Füredi's theorem (Theorem 7.15) on two levels of the Boolean lattice which implies a general upper bound $\mathbf{ex}(n, L) = O(n^{2-(1/r)})$ for any graph L with vertices of degrees at most r on one side of L .

7.1. The Erdős-Simonovits Reduction and the Cube theorem

We have already mentioned Theorem 1.5, on the extremal number of the cube. Here we formulate a sharpening of it.

Theorem 7.1 ([90]). *Let Q_8 denote the graph determined by the 8 vertices and 12 edges of a cube, and Q_8^+ denote the graph obtained by joining two opposite vertices of this cube. Then*

$$\mathbf{ex}(n, Q_8) \leq \mathbf{ex}(n, Q_8^+) = O(n^{8/5}).$$



One reason why Erdős and Simonovits considered the extremal problem of the Cube graph was that this was one of Turán's originally posed problems. The reason that Q_8^+ was also considered was that Erdős and Simonovits got it for free: their proof of Theorem 1.5 gave the same upper bound for Q_8^+ .

Let L be a bipartite graph with partite sets X and Y , and let $K_{t,t} * L$ denote the graph obtained by completely joining one partite set of $K_{t,t}$ to X and the other to Y .


Theorem 7.2 (Erdős and Simonovits Reduction Theorem [90]). *If L is a bipartite graph with $\mathbf{ex}(n, L) = O(n^{2-a})$, $a \leq 1$, and b is defined by $\frac{1}{b} = \frac{1}{a} + t$, then $\mathbf{ex}(n, K_{t,t} * L) = O(n^{2-b})$*

The proof can go by induction on t and by counting the number of C_4 's.

Let H be the graph obtained by deleting just three independent edges from $K_{4,4}$. Since $H = K_{1,1} * C_6$, Theorem 7.2 and $\mathbf{ex}(n, C_6) = O(n^{4/3})$ (Corollary 4.9) imply Theorem 7.1.

Since $\mathbf{ex}(n, L) = O(n)$ if L is a tree, so we have the following result:

Corollary 7.3. For any tree L , $\mathbf{ex}(n, K_{t,t} * L) = O(n^{2-(1/(t+1))})$.



This can be considered as a generalization of the Kővári-T. Sós-Turán theorem, since for $L = K_2$ we have $K_{t+1,t+1} = K_{t,t} * K_{1,1}$. Since $Q_8 - e$ is a subgraph of $K_{1,1} * P_6$, Corollary 7.3 implies

Corollary 7.4 (Erdős and Simonovits). $\mathbf{ex}(n, Q_8 - e) = O(n^{3/2})$.

Further,

Theorem 7.5 (Erdős). Delete an edge from $K_{a,b}$.

For the resulting $L = K_{a,a} - e$ we have $\mathbf{ex}(n, L) = O(n^{2-\frac{1}{a-1}})$.

Indeed, for $a \geq 3$ the graph $K_{a,a} - e$ is a subgraph of $K_{a-2,a-2} * P_4$.

Since $K_{b,b} - K_{a,a}$ (for $b - 2 \geq a \geq 1$) can be written as $K_{b-a-1,b-a-1} * T$ where T is a double star, Corollary 7.3 also implies that $\mathbf{ex}(n, K_{b,b} - K_{a,a}) = O(n^{2-(1/(b-a))})$. For this important case Füredi and West gave a sharper upper bound.

Theorem 7.6 ([121]). For every $n \geq b > a$ we have

$$\mathbf{ex}(n, K_{b,b} - K_{a,a}) \leq \frac{1}{2}(b + a - 1)^{1/(b-a)} n^{2-(1/(b-a))} + \frac{1}{2}(b - a - 1)n.$$

In particular, it gives $\mathbf{ex}(n, K_{3,3} - e) \leq \frac{1}{2}\sqrt{3}n^{3/2} + O(n)$. This was further improved by J. Shen [214] to

$$\mathbf{ex}(n, K_{3,3} - e) \leq \frac{\sqrt{15}}{5}n^{3/2} + O(n).$$

He also showed that $\mathbf{ex}(n, n, K_{3,3} - e) \leq (4/\sqrt{7})n^{3/2} + (n/2)$.

Pinchasi and Sharir extended the cube theorem, using a somewhat different proof:

Theorem 7.7 (Pinchasi and Sharir [204]). A bipartite graph $G[A, B]$ with $|A| = m$ and $|B| = n$, not containing the cube Q has

$$O(n^{4/5}m^{4/5} + mn^{1/2} + nm^{1/2})$$

edges.

Another, more explicit proof for Theorem 7.7 was presented in [115].

Historical remarks:


(a) Erdős and Simonovits first proved the Cube theorem, using the Δ -almost-regularization.

(b) It seems that Theorem 7.2 covered all the cases known until that point.

(c) This (i.e. the Cube Recursion Theorem) was the first case, where one got an exponent, different from $2 - (1/a)$ and $1 + (1/a)$. Actually, Erdős thought earlier, that all the exponents must be of this form, (see [70]). This was disproved in their paper [90]: not by the cube, since there is no good lower bound for the cube: even $\mathbf{ex}(n, Q_8)/n^{3/2} \rightarrow \infty$ is not known. However, a more complicated example, for which the lower bound – using random graphs – was good enough, disproved Erdős’ conjecture. Actually, one thinks that each rational $\alpha \in (0, 1)$ is extremal exponent for some finite \mathcal{L}_α , see Conjecture 2.37.

To disprove the Erdős conjecture concerning the exponents are of the form $1 + (1/a)$ or $2 - (1/a)$ it is enough to notice that we have graphs H with

$$c_H n^{(8/5) - \varepsilon(H)} < \mathbf{ex}(n, H) = O(n^{8/5}), \quad (c_H > 0, \varepsilon(H) < \frac{1}{10}).$$



More generally, consider the graph $H(t, \ell)$ obtained by connecting a $\Theta(3, \ell)$ to $K(t, t)$, as described in Theorem 7.2. By the Theorem 2.26 (lower bound) and Theorem 7.2 and Theorem 7.8 (upper bound) we obtain

$$c_{\ell,t} n^{2 - \frac{2\ell+2t}{3\ell+t^2+2t(\ell+1)-1}} < \mathbf{ex}(n, H_{t,\ell}) \leq \tilde{c}_{\ell,t} n^{2 - \frac{2}{2t+3}}.$$


So all the numbers $2 - \frac{2}{2t+3}$ are points of accumulations of exponents, in this sense. Actually, applying this argument with $t = 1, \ell = 3$, we get a simple counterexample, with the upper bound $O(n^{8/5})$ and a lower bound $cn^{2-(8/17)}$ ($c > 0$).

(d) In [92], Erdős and Simonovits proved the Supersaturated graph theorem (see Section 11) corresponding to the cube, thus providing a second proof of the Cube Theorem, that needed “less regularization”.

7.2. Theta graphs and the Faudree-Simonovits reduction

There is an alternative proof for the Bondy–Simonovits Theorem in [99]. This proof enabled a generalization to Θ -graphs. Recall that $\Theta_{k,\ell}$ denotes the graph consisting of ℓ paths of length k with the same endpoints but no inner intersections. We have $v(\Theta_{k,\ell}) = 2 + (k - 1)\ell$ and $e(\Theta_{k,\ell}) = k\ell$.

Theorem 7.8 (Theta-graph, Faudree–Simonovits [99]). *For fixed k and $\ell \geq 2$ one has $\text{ex}(n, \Theta_{k,\ell}) = O(n^{1+(1/k)})$.*




This exponent is conjectured to be the best possible, see Conjecture 4.11.

Applying the Erdős–Rényi Random Lower bound (Theorem 2.26) in its simpler form to $\Theta_{k,\ell}$ we get

$$\text{ex}(n, \Theta_{k,\ell}) > c_{k,\ell} n^{1+\frac{1}{k}-\frac{2}{k\ell}},$$

asymptotically matching the upper bound’s exponent.



The proof of Theorem 7.8 came from a “Recursion” theorem, asserting that if one knows good upper bounds for an L , and L^* is built from L in a simple way, then one has a good upper bound on $\text{ex}(n, L^*)$ as well.

Definition 7.9. Let L be a bipartite graph, with a fixed 2-colouring ψ in RED-BLUE with h RED vertices. Let $x \notin V(L)$ be a vertex from which h independent paths of $k-1$ edges go the RED vertices of L , (these paths intersect only in x). Denote the obtained graph by $L_k(L, \psi)$.

Theorem 7.10 (Faudree-Simonovits Reduction, Trees [99]). *If L is a tree, then*

$$\text{ex}(n, L_k(L, \psi)) = O(n^{1+(1/k)}).$$

The Theta graph $\Theta_{k,\ell}$ is obtained from a star of ℓ edges. One has to be cautious with the next theorem, see Remark 7.12.

Theorem 7.11 (Faudree-Simonovits Reduction, General Case [99, 100]). *Let L be an arbitrary bipartite graph with a fixed coloring ψ and assume that*

$$\text{ex}^*(n, L) = O(n^{2-\alpha}). \tag{7.1}$$

Then for


$$\beta = \frac{\alpha + \alpha^2 + \dots + \alpha^{k-2}}{1 + \alpha + \alpha^2 + \dots + \alpha^{k-2}}$$

we have

$$\mathbf{ex}(n, L_k(L, \psi)) \leq \mathbf{ex}^*(n, L_k(L, \psi)) = O(n^{2-\beta}). \tag{7.2}$$

Remark 7.12. Most probably, this recursion is never sharp but for trees. In its proof one has to apply standard arguments to subgraphs of $K(m, n)$ where $n \gg m$. We very seldom have matching lower and upper bounds in such cases.

7.3. A universal graph and dependent random choice



Erdős asked the following question: what are the extremal numbers for the two graphs on the left: The left one will be called M_{10} , the right one M_{11} and they are described as special cases of the following

Definition 7.13. Let k, r and t be given positive integers. $U(k, r, t)$ is obtained from the k vertices x_1, \dots, x_k by joining to each of the r -element subsets of $\{x_1, \dots, x_k\}$ t distinct vertices y_{i_1, \dots, i_r}^j . $U^+(k, r, t)$ is obtained from $U(k, r, t)$ by joining a new vertex w to all $x_h, h = 1, \dots, k$.

Problem 7.14 (Erdős). Determine (or estimate) $\mathbf{ex}(n, L)$ for $L := M_{10} = U(4, 2, 1)$ and $L := M_{11} = U^+(4, 2, 1)$.

One could ask for the motivation: why these graphs? Perhaps having obtained the cube theorem, we had good upper and lower bounds only in very special cases, when L contained some sample graphs – say a C_4 for which we have already provided sharp lower bounds. $U(4, 2, 1)$ clearly needed a new approach, and e.g. $U^+(k, 2, 1)$ contains many C_4 's but the earlier methods did not yield appropriate upper bounds. Füredi [107] answered this question proving that $\mathbf{ex}(n, U^+(k, 2, 1)) < k^{3/2}n^{3/2}$. More generally,

Theorem 7.15 (Füredi [107]). Let $U^+(k, r, t)$ be the universal bipartite graph from Definition 7.13. Then there exists a $c = c_r^{k,t} > 0$ such that

$$\mathbf{ex}(n, U^+(k, r, t)) < cn^{2-(1/r)}. \tag{7.3}$$

Concerning the Erdős question we have $\mathbf{ex}(n, U^+(k, 2, 1)) < k^{3/2}n^{3/2}$ and more generally

$$\mathbf{ex}(n, U^+(k, 2, t)) < n^{3/2} \cdot \sqrt{\frac{tk(k-1)^2 + 2(k-2)(k-1)}{8}} + n \frac{k-1}{4}. \quad (7.4)$$

Multiplying each vertex $(k-1)$ times in a C_4 -free graph we get a $U^+(k, 2, 1)$ -free graph which yields $\mathbf{ex}(n, U^+(k, 2, 1)) \geq \Omega(k^{1/2}n^{3/2})$.

Erdős had the more general conjecture

Conjecture 7.16 (Erdős, [66], see also [92], [225]). *If every subgraph of the bipartite graph L has a vertex of degree at most r , then*

$$\mathbf{ex}(n, L) = O(n^{2-(1/r)}).$$

The upper bound (7.3) for the universal graph immediately gives

Corollary 7.17. *If L is bipartite and has a 2-coloring where in the first color class all but one vertex is of degree at most r , then*

$$\mathbf{ex}(n, L) = O(n^{2-(1/r)}).$$

Indeed, all such graphs can trivially be embedded into an appropriate $U^+(k, r, t)$.

Alon, Krivelevich, and Sudakov [9] gave a new probabilistic proof (for graphs where on one side all vertices are of degree at most r) with a better constant $c_r^{k,t}$. Their proof method became known as “dependent random choice”; for a survey see [101].

Lemma 7.18 (Dependent random choice, see, e.g., [101]). *Let k, t, r be positive integers. Let G_n be a graph with n vertices and average degree d , d be an integer. If there is a positive integer a such that*

$$\frac{d^a}{n^{a-1}} - \binom{n}{r} \left(\frac{t}{n}\right)^a \geq k,$$

then G_n contains a subset U of at least k vertices such that every r vertices in U have at least t common neighbors.

Note that in this lemma they do not claim that $U(k, r, t)$ is a subgraph. Nevertheless, using this lemma they improve the constant $c = c_r^{k,t}$ in (7.3) from $O((t+1)^{1/r}k^{2-(2/r)})$ to $c \leq 2^{-1+(2/t)}(t+1)^{1/r}k$.

As they mention at the end of [9], both proofs of Theorem 7.15 give a bit more (and thus Corollary 7.17 can be sharpened accordingly):

$$\mathbf{ex}(n, U^{+r}(k, r, t)) < cn^{2-(1/r)}, \quad (7.5)$$

where the graph U^{+r} is obtained from $U^+(k, r, t)$ by replacing the vertex w in Definition 7.13 by an independent set of r vertices with the same neighbors, x_1, \dots, x_k .

However, the method of Dependent random choice gives more. Call a graph L_h on h vertices *r-degenerate* if it satisfies the condition of Conjecture 7.16. In other words, there is an ordering of its vertices x_1, \dots, x_h such that for every $1 \leq i \leq h$ the vertex x_i has at most r neighbors x_j with $j < i$.

Theorem 7.19 (Alon, Krivelevich, and Sudakov [9]). *If L is bipartite r -degenerate graph on h vertices, then for every $n \geq h$*

$$\mathbf{ex}(n, L) \leq h^{1/(2r)} n^{2-(1/4r)}.$$

Applying the above results with $r = 2$, $t = 1$ and $k = c\sqrt{n}$ to find a $U(k, 2, 1)$ one immediately obtains the following. Any graph on n vertices with $c_1 n^2$ edges contains a 1-subdivision of K_k with $k = c_2 \sqrt{n}$ for some positive c_2 depending on c_1 . This answers a question of Erdős [72]. The theorems of Bollobás and Thomason [29] and Komlós and Szemerédi [161] also imply the existence of such a large topological clique but their subgraph is not necessarily a 1-subdivision.

Given any graph L , let \bar{d} denote the $\max_{X \subseteq V(L)} \{2e_L(X)/|X|\}$, the maximum *local average degree*. Then L is $\lfloor \bar{d} \rfloor$ -degenerate. Hence the upper bound of Theorem 7.19 and the random method lower bound in (2.9) yield that

Corollary 7.20. *For every bipartite graph L ,*

$$\Omega(n^{2-c}) \leq \mathbf{ex}(n, L) \leq O(n^{2-(c/8)}), \quad (7.6)$$

where $c = 2/\bar{d}$, is the same as in (2.9).

8. Eigenvalues and extremal problems

Let $A = A(G_n)$ be the adjacency matrix of G_n , and \mathbf{j} be the vector each entry of which is 1. Since

$$e(G_n) = \frac{1}{2} \mathbf{j} A \mathbf{j}^T \quad (8.1)$$

and, more generally,

$$w_k(G_n) = \frac{1}{2} \mathbf{j} A^k \mathbf{j}^T \quad (8.2)$$

counts the number of k -edge walks in G_n , therefore it is not so surprising that eigenvalues can be used in extremal graph problems. An easy to read source on spectra of graphs is Cvetkovič–Doob–Sachs [56].

Theorem 8.1 (Babai–Guiduli [12]). *Let $\Lambda(G) = \max |\lambda_i|$, where $\lambda_1, \dots, \lambda_n$ are the eigenvalues of $A(G_n)$. If $K_{a,b} \not\subseteq G_n$, (and $2 \leq a \leq b$) then*

$$\Lambda \leq \sqrt[a]{b-1} \cdot n^{1-(1/a)} + o(n^{1-(1/a)}). \quad (8.3)$$

Since trivially

$$2e(G_n) \leq \Lambda n \quad (8.4)$$

the inequality (8.3) implies Theorem 2.22 apart from the $o()$ term.

Remark 8.2. For regular or almost regular graphs $\Lambda(G_n) \approx \frac{2e(G_n)}{n}$, and then the two estimates are basically equivalent. The constant in the above theorem is not sharp since – as we know from Theorem 3.19, – the constant can be improved.

We have already mentioned Nikiforov’s result (Theorem 3.21) on the Zarankiewicz problem. In fact, he proved [198] that for all $n \geq b \geq a \geq 2$ and a $K_{a,b}$ -free graph G_n we have

$$\Lambda(G_n) \leq (b-a+1)^{1/a} n^{1-(1/a)} + (a-1)n^{1-(2/a)} + (a-2). \quad (8.5)$$

This improves the coefficient in Theorem 8.1. It also implies Füredi’s bound (3.11) for the $\mathbf{ex}(n, K_{a,b})$ according to (8.4). For C_4 -free graphs he has $\Lambda^2 - \Lambda + 1 \leq n$.

Recall that $T_{n,k}$ denotes the Turán graph, the k -partite complete graph of maximum size. Given a K_{k+1} -free graph G_n Nikiforov [197] showed that $\Lambda(G_n) < \lambda(T_{n,k})$ unless $G = T_{n,k}$. For a recent reference of a generalization see Z. L. Nagy [196].

9. Excluding topological subdivisions

9.1. Large topological subgraphs

We have already mentioned that our classification does not hold for infinite families of excluded subgraphs. One important phenomenon is that $\mathbf{ex}(n, \mathcal{L})$

can be linear for infinite \mathcal{L} even if \mathcal{L} contains only cycles.²⁷ Here we consider a very central graph theoretical problem strongly connected to the 4-colour conjecture.

Definition 9.1. Given a graph H , its *subdivision* (or a topological H) is obtained from it by replacing each edge e of H by some paths P_e so that these paths do not have their inner (new) vertices in common.

Wagner asked if for any integer ℓ there exists a $k = k_\ell$ such that any G with chromatic number $\chi(G) > k_\ell$ must contain a topological subdivision of K_ℓ . This was proved by Gabor Dirac and H. Jung (independently). Answering a question of Dirac, Mader proved the following important result.

Theorem 9.2 (Mader, [183]). *If G_n is an n -vertex graph, and*

$$e(G_n) \geq n(\ell - 1)2^{\binom{\ell-1}{2}-1},$$

then G_n contains a subdivision of the complete ℓ -graph.

This statement is stronger than the original Wagner conjecture, since a graph with large chromatic number contains a subgraph with large minimum degree. Mader, and independently, Erdős and Hajnal conjectured that

Conjecture 9.3 (Mader, Erdős-Hajnal). *There exists a constant $c > 0$ such that if $e(G_n) > c\ell^2 n$, then G_n contains a topological K_ℓ .*

A slightly weaker form of this conjecture was proved by Komlós and Szemerédi, [160], then – by a different method – Bollobás and Thomason [29] proved this conjecture and, almost immediately after that, Komlós and Szemerédi [161] proved Conjecture 9.3 as well.

Theorem 9.4 (Bollobás–Thomason). *Every graph G_n of size at least $256\ell^2 n$ edges contains a topological complete subgraph of order ℓ .*

As to the small values of ℓ , Dirac conjectured that every that for $n \geq 3$ every G_n , with $e(G_n) \geq 3n - 5$ contains a topological K_5 . This improvement of the famous Kuratowski theorem was proved by Mader in [184] and the corresponding extremal graphs were characterized in [186]. The reader is recommended the excellent “featured review” of Carsten Thomassen on the paper of W. Mader [184], on the MathSciNet.

An excellent survey of Mader on this topic is [185]

²⁷Here the simplest case is Theorems 5.1.

9.2. Turán numbers of subdivided graphs

Let ε be a positive real, $0 < \varepsilon < 1$. Kostochka and Pyber [163] proved that every n -vertex graph G_n with at least $4t^2 n^{1+\varepsilon}$ edges contains a subdivision of K_t on at most $(7t^2 \ln t)/\varepsilon$ vertices, where $0 < \varepsilon < 1$. This (for $t = 5$) answers a question of Erdős about finding a non-planar subgraph of size $c(\varepsilon)$ in a graph with $n^{1+\varepsilon}$ edges.

Recently, T. Jiang [150] improved the Kostochka-Pyber upper bound to $O(t^2/\varepsilon)$. On the other hand, for each $0 < \varepsilon < 1$ and $n > n_0(\varepsilon)$ there are n -vertex graphs of girth at least $1/\varepsilon$ (see Corollary 2.30). In such a graph any subdivision of K_t must contain $\Omega(t^2/\varepsilon)$ vertices, so Jiang’s result is sharp.

Theorem 9.5 (Jiang and Seiver [151]). *Let L be a subdivision of another graph H . For each edge $xy \in E(H)$ let $\ell(x, y)$ denote the length of the path in L replacing the edge xy . Suppose that $\ell(x, y)$ is even for each edge of H , and let $\min\{\ell(x, y) : xy \in E(H)\} = 2m$. Then $\mathbf{ex}(n, L) = O(n^{1+(8/m)})$.*


The main tools in the proof are the Dependent Random Choice, Lemma 7.18, and the Erdős-Simonovits Δ -almost-regularization, Theorem 2.19.

10. Hypergraph Extremal Problems

10.1. Positive Density problems

This is a short detour into Hypergraph Extremal Problems. Now our “Universe” is the class of r -uniform hypergraphs. Katona, Nemetz and Simonovits [154] showed (using a simple averaging) that

Theorem 10.1 (Katona, Nemetz and Simonovits [154]). $\mathbf{ex}_r(n, \mathcal{L})/\binom{n}{r}$ is monotone decreasing, and therefore convergent.



The hypergraph extremal problems are extremely hard. Even the simplest extension of Turán’s theorem is unsolved: Let $K_4^{(3)}$ be the three-uniform hypergraph with 4 vertices and 4 triples.

Construction 10.2 (Turán, the simplest case). *We partition n vertices into three classes C_1, C_2, C_3 and we take all the triples of the form (x, y, z) , where*

- (a) $x \in C_i, y \in C_i, z \in C_{i+1}$ (where the indices are taken mod 3);
- (b) the three vertices are in three different groups.

One can easily see that his construction contains no $K_4^{(3)}$.

Conjecture 10.3 (Turán). *Construction 10.2 is asymptotically extremal for $K_4^{(3)}$. (Perhaps it is extremal, not only asymptotically extremal, at least for $n > n_0$.)*

Here we cut it short and recommend the reader (among others) the survey of Füredi on hypergraph extremal problems [108], and also the papers of Füredi and Simonovits, [120], Keevash and Sudakov [156], Füredi-Pikhurko-Simonovits [119], and the survey of Keevash [155].

10.2. Degenerate hypergraph problems

For r -uniform hypergraphs the r -partite graphs generalize the bipartite graphs. An important illustration of this is the one below, extending Theorem 2.31.

Theorem 10.4 (Degenerated hypergraph problems). *For an r -uniform extremal hypergraph problem of $\mathcal{L}^{(r)}$, $\mathbf{ex}(n, \mathcal{L}^{(r)}) = o(n^r)$, if and only if there is an $L \in \mathcal{L}^{(r)}$ which can be r -vertex-colored so that each hyperedge of L gets r distinct colors.*

Theorem 10.4 is an easy corollary of the following theorem of Erdős, (which generalizes Theorem 2.22).

Theorem 10.5 (Erdős [65]). *Let $K^{(r)}(a_1, \dots, a_r)$ be the r -uniform hypergraph with r vertex-classes C_1, \dots, C_r , where $|C_i| = a_i$, and $a_1 = t$. Then*

$$\mathbf{ex}^{(r)}(n, K^{(r)}(a_1, \dots, a_r)) = O(n^{r-(1/t^{r-1})}).$$

Extending some problems and results for ordinary graphs, Brown, Erdős and Sós started investigating the following

Problem 10.6 (Brown, Erdős, Sós, [39], [40]). Consider r -uniform hypergraphs for some fixed r , and denote by $\mathbb{H}_{k,\ell}^r$ the family of r -uniform k -vertex hypergraphs with ℓ hyperedges. Determine or estimate $f_r(n, k, \ell) := \mathbf{ex}(n, \mathbb{H}_{k,\ell}^r)$.

Brown, Erdős, and Sós proved many upper and lower bounds for special cases of Problem 10.6. We have already mentioned one of them: the

$f_3(n, 6, 3)$ -problem.²⁸ It is easy to see that $f_3(n, 6, 3) < \frac{1}{6}n^2$. The real question was if $f_3(n, 6, 3) = o(n^2)$ or not. Ruzsa and Szemerédi [210] proved that the answer is YES. We formulated this in Theorem 1.9. This theorem became a very important one. We originate, among others, the “Removal Lemma” from here.

We shall return to this problem in the section on applications.

11. Supersaturated graphs

The theory of Supersaturated extremal problems is a very popular area today. Here we shall restrict ourselves to the supersaturated extremal graph problems related to bipartite excluded graphs, just mention a few further references, like Lovász and Simonovits [178], Razborov, [179], Lovász [177], Reiher [205].

Given a graph G , denote by $N(G, F)$ the number of subgraphs of G isomorphic to F . Here we have to be slightly cautious: if F has non-trivial automorphisms, then we can count isomorphisms or isomorphic subgraphs, and the ratio of these two numbers equal to the automorphism number.

A theorem which asserts that a graph G_n contains very many graphs L from a family \mathcal{L} is called a **theorem on supersaturated graphs**. Such theorems are not only interesting in themselves, but also are often useful in establishing other extremal results. At this point it is worthwhile mentioning such a result for complete bipartite graphs, obtained by Erdős and Simonovits [94]:

Theorem 11.1 (Number of complete bipartite graphs). *For any integers a and b there exists a constant $c_{a,b} > 0$ such that if G_n is a graph with m edges, then G_n contains at least $\lfloor c_{a,b} m^{ab} / n^{2ab-a-b} \rfloor$ copies of $K_{a,b}$.*

Corollary 11.2. *Let $c > 0$. If $e(G_n) = E > (1 + c)\text{ex}(n, C_4)$, then G_n contains at least $\gamma e^4 / n^4$ copies of C_4 , for some $\gamma(c) > 0$. The random graph with E edges shows that this is sharp.*

Proof of the Cube Theorem (Sketch). Apply Theorem 2.19 obtaining a Δ -almost-regular (bipartite) $\tilde{G}_n \subseteq G_n$. Apply the corollary to this \tilde{G}_n . It contains $\gamma \frac{E^4}{n^4}$ C_4 's. On the average, an edge of G_n is contained in $\gamma E^3 / n^4$ copies of C_4 . Take a typical edge xy : the bipartite graph $G[U, V]$ spanned

²⁸If $r = 3$, then we delete the subscript in f_3 .

by the neighbors $U := N(x)$ and $V := N(y)$ – by $\mathbf{ex}(m, C_6) = O(m^{4/3})$, – will contain a C_6 . Now, xy and this C_6 will provide a Q_8^+ : a cube with a diagonal. \square

Basically the same argument proves Theorem 7.2.

11.1. Erdős-Simonovits-Sidorenko conjecture

In this part $\chi(L) = 2$. Erdős and Simonovits [225] formulated three conjectures and also that the main idea behind these conjectures is that the number of copies of subgraphs $L \in G_n$ is minimized by the random graph if $E = e(G_n)$ is fixed and is not too small.

To formulate these conjectures, first we calculate the “expected number of copies” of $L \subseteq R_n$ if R_n is a random graph with edge probability $p = E/\binom{n}{2}$. Let $v = v(L)$, and $e = e(L)$. Clearly, if the edges are selected independently, with probability p , then R_n contains $\binom{n}{v}$ possible v -tuples, each containing the same number a_L of copies of L , and therefore

$$\mathbb{E}(\#(L \subseteq R_n)) = (a_L + o(1)) \frac{n^v}{v!} p^e = (a_L + o(1)) \frac{n^v}{v!} \left(\frac{2E}{n^2} \right)^e = a_L^* \frac{E^e}{n^{2e-v}} \quad (11.1)$$

Conjecture 11.3 (Erdős–Simonovits, [225]). *There are two constants, $\Omega = \Omega_L > 0$ and $c = c_L > 0$ such that if $E > \Omega \cdot \mathbf{ex}(n, L)$, then any graph G_n with E edges contains at least*

$$c_L \frac{E^e}{n^{2e-v}}$$

copies of L .

This was the weakest form. The strongest form of this conjecture was

Conjecture 11.4 (Erdős and Simonovits). *For every $\varepsilon > 0$, if $E > (1 + \varepsilon) \cdot \mathbf{ex}(n, L)$, then any graph G_n with E edges contains at least $(1 + \varepsilon) \mathbb{E}\mathbb{R}(n, L, E)$ copies of L , if $n > n_0(\varepsilon)$, where $\mathbb{E}\mathbb{R}(n, L, E)$ denotes the expected number of edges of a random Erdős-Rényi graph with n vertices and E edges.*

Obviously, one has to assume that $e(G_n) > \mathbf{ex}(n, L)$.

Remark 11.5 (Relation to Sidorenko’s Conjecture). At first sight Sidorenko’s conjecture [216] seems to be sharper than the above one. This is not the case. In fact, Sidorenko’s Conjecture applies only to dense host graphs. There, as Sidorenko points out in his papers, the two versions are equivalent.

Also, it is obvious that there is not much difference if we consider above the hypergeometric model of random graphs, where the number of vertices and edges are given, or if we fix only n but the edges are taken independently, and therefore $e(G_n)$ follows a binomial distribution.

Sidorenko, working on applications of extremal graph theorem to probability distribution translated the above conjecture to integrals and arrived at a conjecture [216], where the error terms disappeared. The meaning of his version was that if one considers dense graphs and defines $L \subseteq R$ for the case when G is a function, generalizing the notion of graphs, then the Random Continuous graph will have the least number of copies of L , more precisely, that will minimize the corresponding integral.

We skip the formulation of this problem, just refer to some papers of Lovász, and, Hatami [140], and to the book of Lovász [177].

Jagger, Štoviček, and Thomason [148] investigated the following problem originating from a conjecture of Erdős, disproved by Thomason.

Problem 11.6. Given a sample graph L , denote by $\rho_L(G_n)$ the sum of copies of L in G_n and in its complementary graph. What is the minimum $\Gamma_n(L)$ of this, taken over all n -vertex graphs?

Erdős conjectured that the random graph yields the minimum, for K_4 . This was disproved by Thomason [237]. Investigating the case of general L , Jagger, Štoviček, and Thomason proved some interesting results in connection with Sidorenko's conjecture.

Here we should emphasize that there is a slight difference between looking for copies of an L in G_n or for copies of homomorphic images: In the second case we allow vertices to map into G_n with some coincidences.

As to the Sidorenko Conjecture, the first unknown case (as Sidorenko mentions) is when we delete the edges of a Hamiltonian cycle from $K_{5,5}$.

Theorem 11.7 (Conlon, Fox, Sudakov [52] [53]). *The Sidorenko Conjecture holds if $L = L[A, B]$ is a bipartite graph with a vertex $x \in A$ completely joined to B .*

Remark 11.8. Unfortunately, we do not have sufficiently good lower bounds for the extremal problem of the cube. The Erdős–Simonovits Conjecture was proved for Q_8 in [92].

Hatami proved the Sidorenko conjecture for any cube (i.e. of any dimension), yet, that was not really enough to provide a reasonable upper bound

for the 4-dimensional cube. This reflects some difference between extremal problems and the corresponding Supersaturated Graph Problems (at least, for dense host graphs).

12. Ordered structures

12.1. Directed graphs, ordered graphs

There is an extensive literature on Digraph extremal problems, see e.g., the survey of Brown and Simonovits [42], or [41]. We skip here the general theory.

Denote $\overrightarrow{\text{ext}}(n, \vec{L})$ the maximum number of edges in a directed graph not containing the oriented subgraph \vec{L} . For every \vec{L} containing a directed path of length 2 one has $\overrightarrow{\text{ext}}(\vec{L}) \geq \lfloor n^2/4 \rfloor$. Indeed, it is easy to orient the edges of $K_{n/2, n/2}$ such that $\vec{L}_{1,a,b}$ does not appear. For bipartite \vec{L} it is more interesting to consider the minimum outdegree.

Consider the following directed graph $\vec{L}_{1,a,b}$ on $1 + a + b$ vertices $w, x_1, \dots, x_a, y_1, \dots, y_b$. The oriented edges are w to x_i and x_i to y_j ($1 \leq i \leq a, 1 \leq j \leq b$).

Theorem 12.1 (Erdős, Harcos and Pach [82]). *Given $a \geq b$ integers, there exists a $c = c_{a,b} > 0$ such that the following holds. Any oriented graph with minimum out-degree $\delta^+ \geq cn^{1-(1/a)}$ contains a copy of $\vec{L}_{1,a,b}$.*

This result opened up a new interesting field with many open problems.

Another ordered Turán function was defined by Timmons [239]. He showed that if a graph with vertex set $\{1, 2, \dots, n\}$ has at least

$$(1 + o(1))(2/3)n^{3/2}$$

edges, then it contains a C_4 with vertices $a_1b_1a_2b_2$ such that $a_1, a_2 < b_1, b_2$. He extended other ordinary Turán problems to these zig-zag type questions. Many problems remain unsolved.

12.2. Erdős-Moser conjecture on unit distances

Erdős and Leo Moser [83] conjectured that

Conjecture 12.2. *If n points of the plane are in convex position, then the number of unit distances among them is $O(n)$.*

Füredi proved a slightly weaker result:

Theorem 12.3 (Füredi [106]). *If n points are in convex position in the plane, then there are at most $O(n \log n)$ unit distances among them.*

To prove this, Füredi directly formulated the excluded Ordered Matrix Property and solved a matrix-containment problem. The crucial point of his proof was Theorem 12.6 below.

The best known lower bound in the Erdős-Moser problem, $2n - 7$, is due to Edelsbrunner and P. Hajnal [58].

12.3. Ordered submatrices

The ordered matrix problems partly came from geometric problems (see Binstock and Györi, [23], Füredi [106]), but they are interesting on their own, too. A geometric application, called Erdős-Moser conjecture, is discussed above in Subsection 12.2.

We have already indicated that most extremal graph problems have matrix forms, too: To determine $\mathbf{ex}^*(m, n, L)$ we considered all $m \times n$ 0-1 matrices not containing any permutation of the bipartite adjacency matrix of L .

In the ordered case here we exclude only those submatrices where the indexing of the rows and columns of \mathbf{M} is fixed. This way we exclude fewer subconfigurations.

Definition 12.4 (Matrix containment). Let \mathbf{M} and \mathbf{P} be two 0-1 matrices. We say that \mathbf{M} contains \mathbf{P} if we can delete some rows and columns of \mathbf{M} and then perhaps switch some 1's into 0 so that the resulting matrix be \mathbf{P} . Otherwise we say that \mathbf{M} avoids \mathbf{P} .

So, we can delete rows and columns of \mathbf{M} but can not permute them. Now we can define the Matrix Extremal Problems:

Problem 12.5 (Ordered Matrix Problem). Given an $a \times b$ 0-1 (sample) matrix \mathbf{P} , and a (huge) $m \times n$ 0-1 matrix \mathbf{M} , how many 1's can occur in \mathbf{M} under the condition that \mathbf{M} does not contain \mathbf{P} in the "ordered" way. Denote by $\mathbf{ext}_{\text{mat}}(m, n, \mathbf{P})$ the maximum.

One of the first nontrivial results was

Theorem 12.6 (Füredi [106]). *Let*

$$\mathbf{P} = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}.$$

If the $n \times n$ 0-1 matrix \mathbf{M} does not contain \mathbf{P} , then it has at most $O(n \log n)$ 1's. In fact, $\text{ext}_{\text{mat}}(n, n, \mathbf{P}) = \Theta(n \log n)$.

Tardos [236] proved that $\text{ext}_{\text{mat}}(n, \mathbf{P}) = n \log_2 n + O(n)$.

Completing earlier works of Füredi and Péter Hajnal [116] Tardos [236] classified the ordered matrix Turán numbers for all small submatrices. The extremely slow growing inverse Ackermann function is denoted by $\alpha(n)$.

Theorem 12.7 ([116], [236]). *If \mathbf{P} is a 0-1 matrix with at most four 1's, then*

$$\text{ext}_{\text{mat}}(n, n, P) = \begin{cases} 0 & \text{or} \\ \Theta(n), & \text{or} \\ \Theta(n\alpha(n)), & \text{or} \\ \Theta(n \log n), & \text{or} \\ \Theta(n^{3/2}). \end{cases}$$

12.4. Ordered matrices and the Stanley-Wilf conjecture on subpermutations

Trying to prove the Erdős-Moser Conjecture, Füredi and Péter Hajnal [116] arrived at the following conjecture, proved by Marcus and Tardos.

Theorem 12.8 (Füredi-Hajnal conj. [116]/Marcus-Tardos theorem [187]). *For all permutation matrices P we have $\text{ext}_{\text{mat}}(n, n, \mathbf{P}) = O(n)$.*

This time there was a famous Stanley-Wilf conjecture around, on the number of permutations “avoiding” a fixed permutation. To formulate it, we need to define the Permutation containment:

Definition 12.9 (Permutation containment). We say that a permutation $\sigma : [1, n] \rightarrow [1, n]$ contains a permutation $\pi : [1, k] \rightarrow [1, k]$, if there exist $1 \leq x_1 < x_2 < \dots < x_k \leq n$ for which

$$\sigma(x_i) < \sigma(x_j) \quad \text{if and only if} \quad \pi(i) < \pi(j).$$

The famous Stanley–Wilf conjecture²⁹ states that

Conjecture 12.10 (Stanley–Wilf). *For any permutation pattern q , if $S_n(q)$ is the number of permutations of length n avoiding the pattern q , then there is a constant c_q so that $S_n(q) \leq c_q^n$.*

Theorem 12.11 (Klazar [158]). *The Füredi–Hajnal conjecture implies the Stanley–Wilf conjecture.*

So Marcus and G. Tardos settled this conjecture as well.

Remark 12.12. The permutation containment is just a subcase of the more general question. In some other cases there are definite differences between ordinary Turán type extremal problems and the ordered matrix problems. For a special matrix, where the corresponding graph is a tree, hence it has linear Turán function, our threshold function turns out to be $\Theta(n \log n)$.

13. Applications in Geometry

13.1. Applicability of the Kővári–T. Sós–Turán bound

We have mentioned that Theorem 2.22 is applicable in several cases. Here we mention only two.

(A) The Unit Distance Graph of the Plane contains no $K(2, 3)$. Erdős used this to estimate the number of unit distances by $O(n^{3/2})$.

(B) G. Megyesi and Endre Szabó³⁰ answered a question of F. E. P. Hirzebruch using this theorem.

Assume that we are given k smooth curves in the the Complex Projective Plane and assume that their union has only nodes and tacnodes³¹ as singularities. Let $t(k)$ denote the maximum number of tacnodes in such cases. Hirzebruch proved that $t(k) \leq \frac{4}{9}k^2 + \frac{4}{4}k$. Hirzebruch asked if $\limsup t(k)/k > 0$.

Theorem 13.1 (G. Megyesi, and E. Szabó [191]). *There exist three positive constants, A, B and C for which*

$$Ak^{1+(B/\log \log k)} \leq t(k) \leq Ck^{2-(1/7633)}.$$

²⁹Marcus and Tardos [187] write that it is difficult to locate the corresponding reference.

³⁰We use the longer versions of the names whenever we see chances to mix up authors of similar names.

³¹Tacnode means roughly that the curve is touching itself.

13.2. Unit Distances

Erdős was interested in the following problem:

Problem 13.2 (Unit distances). Given an n -element set in the d -dimensional Euclidean space \mathbb{E}^d , how many of the distances can be the same, say equal to 1?

Conjecture 13.3 (Unit distances). *For any $\varepsilon > 0$, there is an n_0 such that if $n > n_0$ and given an n -element set in the plane \mathbb{E}^2 , then the number of unit distances is at most $n^{1+\varepsilon}$.*

The motivation of this conjecture is – as Erdős observed – that if we arrange the $n = k \times k$ points into a $k \times k$ grid, and rescale this grid so that the “most popular” distance be 1, then this distance will occur at most $n^{1+\varepsilon}$ times, (actually, approximately $n^{1+(c/(\log \log n))}$ times). So Erdős conjectured that the number of unit distances in the plane has an upper bound of roughly this form.

The first upper bound was a trivial application of Theorem 2.22:

Theorem 13.4 (Unit distances, Erdős 1946). *Given n points in the plane, the number of unit distances among them is at most*

$$\text{ex}(n, K_{2,3}) < \left(\frac{1}{\sqrt{2}} + o(1) \right) n^{3/2}.$$

In \mathbb{E}^3 the number of unit distances is at most

$$\text{ex}(n, K_{3,3}) < c_{3,3} n^{5/3}. \tag{13.1}$$

Proof. Since two circles intersect in at most 2 points, the Unit Distance Graph of \mathbb{E}^2 contains no $K_{2,3}$. This implies the first inequality. Since 3 unit balls intersect in at most 2 points, the Unit Distance Graph of \mathbb{E}^3 does not contain any $K_{3,3}$. This implies (13.1). \square

Remarks 13.5. (a) Everything is different for the higher dimensions: \mathbb{E}^4 contains two orthogonal circles of radii $\frac{1}{\sqrt{2}}$, and these form a $K(\infty, \infty)$ in the corresponding unit graphs of \mathbb{E}^d , for $d \geq 4$. (This is the so called Lenz Construction.) (See also Section 3.6.)

(b) How sharp is this application? As the reader can see, it is very far from the conjectured upper bound. However, just to improve it to $o(n^{3/2})$ is

non-trivial (Józsa-Szemerédi [152]). Actually, for the plane an $O(n^{4/3})$ upper bound was proved by Beck and Spencer [20] and Spencer, Szemerédi and Trotter [233], which is sharp if we do not insist on Euclidean metric, only on “normed spaces”. For this see the results of Peter Brass [35] and Pavel Valtr [245].

13.3. Cells in line arrangements

Let $\mathcal{I}(m, n)$ denote the maximum number of edges in m distinct cells determined by an arrangement of n lines in the plane. Canham [45] showed that for an absolute constant $c > 0$

$$\mathcal{I}(m, n) < c(m\sqrt{n} + n). \quad (13.2)$$

Indeed, if we construct a bipartite graph where one side of the vertex set consists of the m cells (or any other family of m convex sets with disjoint interiors), the other side of the vertex set consists of the n (tangent) lines and two vertices are joined if the corresponding geometric objects are incident, then it is easy to see that this graph does not contain a $K_{5,2}$. \square

This was a first nontrivial step toward the determination of the exact order of the magnitude of $\mathcal{I}(m, n)$ by Clarkson, Edelsbrunner, Guibas, Sharir, and Welzl [49]; it is $\Theta(n^{2/3}m^{2/3} + n)$. More about this and other geometric applications see the monograph of Pach and Agarwal [199].

14. Further connections and problems

14.1. Connections of hypergraphs and critical graphs

We discussed Degenerate Hypergraph Extremal Problems in Section 10. Here we continue that line.

Excluding the 3-uniform hypergraph cones

Many of the other results, problems of [39] were also degenerate ones. One of them was where \mathcal{T} is the family of triangulations of the 3-dimensional sphere. This problem gave the name to this paper [39]. The crucial point was excluding the double cones:

Definition 14.1 (r -cones). The vertices of the 3-uniform hypergraph $Q_{r,t}$ are x_1, x_2, \dots, x_r , and y_1, y_2, \dots, y_t for some t , and the hyperedges are $x_i y_j y_{j+1}$,

for all the possible i, j , where $y_{t+1} = y_1$. Further, $\mathbb{Q}_r := \{Q_{r,t} : t = 3, 4, 5, \dots\}$.

Theorem 14.2 (Brown, Erdős, Sós, $r = 2$, [39], Simonovits [219] $r \geq 2$).

$$\text{ex}(n, \mathbb{Q}_r) := O(n^{3-(1/r)}).$$

For $r = 2, 3$ there are matching lower bounds here. Actually, for $r = 2$ Brown, Erdős and Sós gave a construction, where not only the double-cone was excluded, but all the triangulations of the sphere. In Simonovits' lower bound only the double cone was considered.

In [220] Simonovits returned to this question and – using the main idea of Brown's construction [36] – he proved

Theorem 14.3 (Simonovits [220]). *There are (finite geometric) 3-uniform hypergraphs without triple-cones (i.e. without hypergraphs from \mathbb{Q}_3) and still having at least $cn^{3-(1/3)}$ triples.*

We saw that for the family of triangulations of the sphere, and for the family of Double Cones the extremal number is $O(n^{3-(1/2)})$ [39], (see [219]).

Brown, Erdős and T. Sós arrived at their question (most probably) since they wanted to generalize certain results from ordinary graphs to hypergraphs. Simonovits came from a completely different direction: he used this to disprove a conjecture of Gallai on independent vertices in 4-colour-critical graphs.

G is colour-edge-critical, if deleting any edge of G , we get a $(\chi(G) - 1)$ -chromatic graph. The 3-colour-critical graphs are the odd cycles, so the problem of critical graphs becomes interesting for the 4-chromatic case. Here we shall restrict ourselves to this case and suggest the reader to read Bjarne Toft's results on this topic in general.

Erdős asked if a 4-colour-critical graph can have cn^2 edges and Bjarne Toft constructed such a 4-chromatic graph [240] of $\approx \frac{n^2}{16}$ edges. This and some related questions can also be found in Lovász' book: Combinatorial Exercises [176].

Gallai had many beautiful conjectures on 4-colour-critical graphs. One of them, however, was “completely demolished”. He conjectured that if G_n is 4-colour-critical, then $\alpha(G_n) \leq n/2$. $G_{4m+2} = C_{2m+1} \otimes C_{2m+1}$ is 6-critical, with $d_{\min}(G_{4m+2}) = 2m + 3$. Simonovits – “blowing up” the vertices in one of the two odd cycles, – proved that there are 6-critical graphs G_n with $\alpha(G_n) = n - o(n)$.

It turned out that slightly earlier Brown and Moon [38] already disproved Gallai’s conjecture for the 4-chromatic case, with a “clever but simple” construction.

Theorem 14.4 (Brown and Moon [38]). *There exist 4-chromatic edge-critical graphs G_n with $\alpha(G_n) > n - c\sqrt{n}$, for some constant $c > 0$.*

Next, Bjarne Toft came up with his construction, mentioned above. Using this and a hypergraph extremal theorem, Simonovits proved

Theorem 14.5. *There exists a constant $c_2 > 0$ such that if G_n is 4-colour-critical, then $\alpha(G_n) \leq n - c_2n^{2/5}$.*

This was obtained as follows: Simonovits reduced the original problem to estimating the number of independent vertices of degree 3 in a 4-colour-critical graph. The neighborhoods of these vertices generated a 3-uniform hypergraph $\mathcal{H}_m^{(3)}$ on the remaining vertices. Simonovits – using the Sperner Lemma from Topology proved that if I is a set of independent vertices of degree 3, in $V(G_n)$, then for $m := n - |I|$, $|I| < \mathbf{ex}_3(m, \mathbb{Q}_2) = O(m^{5/2})$, see [219]. He observed that $\mathcal{H}_m^{(3)}$ cannot contain double cones. This proved that $|I| < n - cn^{2/5}$. \square

(b) Lovász observed that instead of excluding the graphs from \mathbb{Q}_2 one can exclude a larger family, $\tilde{\mathbb{Q}}$: those 3-uniform hypergraphs which obey the conclusion of Sperner’s lemma [175]: each pair (x, y) is contained in an even number of hyperedges. This enabled him to completely settle *this* Gallai problem on colour-critical graphs. He proved that $\mathbf{ex}(n, \tilde{\mathbb{Q}}_2) \leq \binom{n}{2}$. So he obtained $|I| < n - c\sqrt{n}$, in Gallai’s problem. Besides proving and using a more applicable extremal graph theorem he also generalized the Brown–Moon construction.

(c) It was an interesting feature of Lovász’ solution that to get an upper bound on $\mathbf{ex}(n, \tilde{\mathbb{Q}})$ he used linear algebra.

We finish this part by sketching the proof of Lovász on the upper bound.

Theorem 14.6 (Lovász [175]). *Let $\mathbb{E}^{(3)}$ denote the family of 3-uniform hypergraphs H in which each pair of vertices is contained in an even number of triplets (i.e. hyperedges). Then $\mathbf{ex}(n, \mathbb{E}^{(3)}) \leq \binom{n}{2}$.*

Proof (Sketch). Assume that $H_n^{(3)}$ contains no subgraphs from $\mathbb{E}^{(3)}$. Consider that vectorspace over $GF(2)$ of dimension $\binom{n}{2}$ where the coordinates

are indexed by pairs from $1, \dots, n$. Represent each triple of $H_n^{(3)}$ by such a vector, where we have 1 in those coordinates which are pairs from our triple. The condition that $H_n^{(3)}$ contains no subgraphs from $\mathbb{E}^{(3)}$ translates into the fact, that these vectors are linearly independent. Hence their number is at most the dimension of the vector-space. \square

Now, repeating the original argument of Simonovits, Lovász obtained

Theorem 14.7. *There exists a constant $c_3 > 0$ such that if G_n is 4-colour-critical, then $\alpha(G_n) \leq n - c_3 n^{1/2}$,*

This with the Brown-Moon construction completely settles Gallai's original problem, providing a matching lower bound. Lovász proved a more general theorem, and extended the Brown-Moon construction as well. We close this part with a beautiful conjecture of Erdős:

Problem 14.8. Is it true that if (G_n) is a sequence of 4-colour-critical graphs, then $d_{\min}(G_n) = o(n)$?

(Simonovits [219] and Toft [240] succeeded in constructing 4-color-critical graphs with minimum degrees around $c\sqrt[3]{n}$.)

Further sources to read: Several related results can be found in Lovász [176].

14.2. A multiplicative Sidon problem and C_{2k} -free graphs

As it was explained in Subsection 1.5, the Erdős problem about $\text{ex}(n, C_4)$ in [60] was obtained from a multiplicative Sidon type question. He investigated subsets of integers of $A \subset \{1, 2, \dots, n\}$ with the property that for any four members of A the pairwise products are distinct, $a_i a_j \neq a_k a_\ell$.

A. Sárközy, P. Erdős, and V. T. Sós [88] started investigating the more general problem.

Problem 14.9. Fix an integer k . How many integers can we take from $[1, n]$ if the product of no k of them is a square.

Interestingly, this Problem also lead to Turán type questions, namely to $\text{ex}(m, n, C_{2k})$ with $m \gg n$. Their conjecture (Conjecture 4.32 above) was proved by Győri [134], see Theorem 4.33. We shall not go into the number theoretic details; just refer the reader again to [134].

14.3. Cycle-free subgraphs of the d -dimensional hypercube

The d -dimensional hypercube, Q^d , is the graph whose vertex set is $\{0, 1\}^d$ and whose edge set is the set of pairs that differ in exactly one coordinate, $e(Q^d) = d2^{d-1}$. Let $\gamma(C_\ell) = \lim_{d \rightarrow \infty} \mathbf{ex}(Q^d, C_\ell)/e(Q^d)$. Note that $\gamma(C_\ell)$ exists, because $\mathbf{ex}(Q^d, C_\ell)/e(Q^d)$ is a non-increasing and bounded function of d . Considering the edges between the levels $2i$ to $2i + 1$ one can see that $\mathbf{ex}(Q^d, C_4) \geq (1/2)e(Q^d)$. The following conjecture is still open.

Conjecture 14.10 (Erdős [74]). $\mathbf{ex}(Q^d, C_4) = (\frac{1}{2} + o(1))e(Q^d)$.

The best upper bound $\gamma(C_4) \leq 0.6226$ was obtained by Thomason and Wagner [238], slightly improving the result of Chung [46].

Erdős [74] also asked whether $\mathbf{ex}(Q^d, C_{2k})$ is $o(d)2^d$ for $k > 2$. This was answered negatively for C_6 by Chung [46], showing that $\gamma(C_6) \geq 1/4$. The best known results for C_6 are $1/3 \leq \gamma(C_6) < 0.3941$ due to Conder [50] and Lu [180], respectively.

On the other hand, for every $t \geq 2$ the inequalities

$$\mathbf{ex}(Q^d, C_{4t}) \leq O(d^{\frac{1}{2} - \frac{1}{2t}} 2^d) \quad \text{and} \quad \mathbf{ex}(Q^d, C_{4t+6}) = O(d^{\frac{15}{16} - \frac{1}{16t}} 2^d) \quad (14.1)$$

were proved by Chung [46] and Füredi and Özkahya [118], respectively. Hence $\gamma(C_{2k}) = 0$, except $\gamma(C_4) \geq 1/2$, $\gamma(C_6) \geq 1/3$ and the problem of deciding whether $\gamma(C_{10}) = 0$ is still open.

Conlon [51] generalized (14.1) by showing $\mathbf{ex}(Q^d, H) = o(e(Q^d))$ for all H that admit a k -partite representation, also satisfied by each $H = C_{2k}$ except for $k \in \{2, 3, 5\}$.

14.4. Two problems of Erdős

Of course, we should close with two open problems of Erdős. The first one is the general version of that problem which was solved in [124] and [141], see Section 1.1.

Conjecture 14.11 (Erdős [75]). *Suppose that G is a graph on $(2k + 1)n$ vertices and of odd girth $2k + 1$. Then G contains at most n^{2k+1} induced cycles of length $2k + 1$.*

The next conjecture is also very famous and is motivated by the blown up pentagon (if we restrict it to $k = 2$.)

Conjecture 14.12 (Erdős [75]). *Suppose that G is a graph on $(2k + 1)n$ vertices and of odd girth at least $2k + 1$. Then G can be made bipartite by omitting at most n^2 edges.*

For the best known results here, for $k = 1$, see Erdős, Faudree, Pach, and Spencer [77] and Erdős, Gyóri, and Simonovits [81].

15. Acknowledgements

The authors are greatly indebted for fruitful discussions and helps to a great number of colleagues, among others to R. Faudree, E. Gyóri, and Z. Nagy.

References

- [1] M. Ajtai, J. Komlós, M. Simonovits and E. Szemerédi: On the approximative solution of the Erdős-Sós conjecture on trees, (manuscript).
- [2] M. Ajtai, J. Komlós, M. Simonovits and E. Szemerédi: Some elementary lemmas on the Erdős-T. Sós conjecture for trees, (manuscript).
- [3] M. Ajtai, J. Komlós, M. Simonovits and E. Szemerédi: The solution of the Erdős-Sós conjecture for large trees, (manuscript, in preparation).
- [4] M. Ajtai, J. Komlós, and E. Szemerédi: On a conjecture of Loeb, in Graph theory, Combinatorics, and Algorithms, Vol. 1, 2 (Kalamazoo, MI, 1992), Wiley-Intersci. Publ., pp. 1135–1146. Wiley, New York, 1995.
- [5] P. Allen, P. Keevash, B. Sudakov and J. Verstraëte: Turán numbers of bipartite graphs plus an odd cycle, submitted.
- [6] N. Alon: Eigenvalues and expanders, *Combinatorica* 6 (1983), 83–96.
- [7] N. Alon: Tools from higher algebra, in : "Handbook of Combinatorics", R. L. Graham, M. Grötschel and L. Lovász, eds, North Holland (1995), Chapter 32, pp. 1749–1783.
- [8] N. Alon, S. Hoory, and N. Linial: The Moore bound for irregular graphs, *Graphs Combin.* 18 (2002), no. 1, 53–57.
- [9] N. Alon, M. Krivelevich, B. Sudakov: Turán numbers of bipartite graphs and related Ramsey-type questions, *Combin. Probab. Comput.* 12 (2003), no. 5-6, 477–494.
- [10] N. Alon and V. D. Milman: λ_1 -isoperimetric inequalities for graphs and superconcentrators, *J. Combin. Theory Ser. B* 38 (1985), 73–88.
- [11] N. Alon, L. Rónyai, and T. Szabó: Norm-graphs: variations and applications, *J. Combin. Theory Ser. B* 76 (1999), 280–290.

- [12] L. Babai and B. Guiduli: Spectral extrema for graphs: the Zarankiewicz problem, *Electronic J. Combin.* 15 (2009), R123.
- [13] R. Baer: Polarities in finite projective planes, *Bull. Amer. Math. Soc.* 52 (1946), 77–93.
- [14] C. Balbuena, P. García-Vázquez, X. Marcote, and J. C. Valenzuela: New results on the Zarankiewicz problem, *Discrete Math.* 307 (2007), no. 17-18, 2322–2327.
- [15] C. Balbuena, P. García-Vázquez, X. Marcote, and J. C. Valenzuela: Counterexample to a conjecture of Győri on C_{2l} -free bipartite graphs, *Discrete Math.* 307 (2007), no. 6, 748–749.
- [16] C. Balbuena, P. García-Vázquez, X. Marcote, and J. C. Valenzuela: Extremal $K(s,t)$ -free bipartite graphs, *Discrete Math. Theor. Comput. Sci.* 10 (2008), no. 3, 35–48.
- [17] P. N. Balister, B. Bollobás, O. M. Riordan, and R. H. Schelp: Graphs with large maximum degree containing no odd cycles of a given length, *J. Combin. Theory B* 87 (2003), 366–373.
- [18] P. N. Balister, E. Győri, J. Lehel, and R. H. Schelp: Connected graphs without long paths, *Discrete Math* 308 (2008), no. 19, 4487–4494.
- [19] S. Ball and V. Pepe: Asymptotic improvements to the lower bound of certain bipartite Turán numbers, *Combin. Probab. Comput.* 21 (2012), no. 3, 323–329.
- [20] J. Beck and J. Spencer: Unit distances, *J. Combin. Theory Ser. A*, 37 (1984), 231–238.
- [21] F. Behrend: On sets of integers which contain no three terms in arithmetic progression, *Proc. Nat. Acad. Sci. US.* 32 (1956), 331–332.
- [22] C. T. Benson: Minimal regular graphs of girths eight and twelve, *Canad. J. Math.* 18 (1966), 1091–1094.
- [23] D. Bienstock and E. Győri: An extremal problem on sparse 0-1 matrices, *SIAM J. Discrete Math.* 4 (1991), no. 1, 17–27.
- [24] P. Blagojević, B. Bukh, and R. Karasev: Turán numbers for $K_{s,t}$ -free graphs: topological obstructions and algebraic constructions, arXiv:1108.5254v3, 3 Jun 2012.
- [25] B. Bollobás: Cycles modulo k , *Bull. London Math. Soc.* 9 (1977), no. 1, 97–98.
- [26] B. Bollobás: *Extremal Graph Theory*, Academic Press, London, 1978.
- [27] B. Bollobás: *Random Graphs*, Academic Press, London, 1985.
- [28] B. Bollobás: Extremal graph theory, in: R. L. Graham, M. Grötschel, and L. Lovász (Eds.), *Handbook of Combinatorics*, Elsevier Science, Amsterdam, 1995, pp. 1231–1292.
- [29] B. Bollobás and A. Thomason: Proof of a conjecture of Mader, Erdős and

- Hajnal on topological subgraphs, *European J. Combin* 19 (1998), 883–887.
- [30] J. A. Bondy: Basic graph theory: paths and circuits, *Handbook of combinatorics*, Vol. I., pp. 3–110, Elsevier, Amsterdam, 1995.
- [31] J. A. Bondy: Extremal problems of Paul Erdős on circuits in graphs, *Paul Erdős and his mathematics, II* (Budapest, 1999), 135–156, *Bolyai Soc. Math. Stud.*, 11, János Bolyai Math. Soc., Budapest, 2002.
- [32] J. A. Bondy and M. Simonovits: Cycles of even length in graphs, *J. Combin. Theory Ser. B*, 16 (1974), 97–105.
- [33] J. A. Bondy and A. Vince: Cycles in a graph whose lengths differ by one or two, *J. Graph Theory* 27 (1998), 11–15.
- [34] S. Brandt and E. Dobson: The Erdős-Sós conjecture for graphs of girth 5, *Selected papers in honour of Paul Erdős on the occasion of his 80th birthday* (Keszthely, 1993), *Discrete Math.* 150 (1996), no. 1-3. 411–414.
- [35] P. Brass: Erdős distance problems in normed spaces, *Comput. Geom.* 6 (1996), no. 4, 195–214.
- [36] W. G. Brown: On graphs that do not contain a Thomsen graph, *Canad. Math. Bull.* 9 (1966), 281–285.
- [37] W. G. Brown: On the non-existence of a type of regular graphs of girth 5, *Canad. J. Math.* 19 (1967), 644–648.
- [38] W. G. Brown and J. W. Moon: Sur les ensembles de sommets indépendants dans les graphes chromatiques minimaux, (French), *Canad. J. Math.* 21 (1969), 274–278.
- [39] W. G. Brown, P. Erdős and V. T. Sós: On the existence of triangulated spheres in 3-graphs, and related problems, *Period Math. Hungar.* 3 (1973), 221–228.
- [40] W. G. Brown, P. Erdős and V. T. Sós: Some extremal problems on r -graphs, *New Directions in the Theory of Graphs* (ed. F. Harary), Academic Press, New York, 1973, pp. 53–63.
- [41] W. G. Brown and M. Simonovits: Digraph extremal problems, hypergraph extremal problems, and the densities of graph structures, *Discrete Math.* 48 (1984), no. 2-3, 147–162.
- [42] W. G. Brown, and M. Simonovits: Extremal multigraph and digraph problems, *Paul Erdős and his mathematics, II* (Budapest, 1999), pp. 157–203, *Bolyai Soc. Math. Stud.*, 11, János Bolyai Math. Soc., Budapest, 2002.
- [43] L. Caccetta and K. Vijayan: Long cycles in subgraphs with prescribed minimum degree, *Discrete Math.* 97 (1991), no. 1–3, 69–81.
- [44] D. de Caen, L. A. Székely: The maximum size of 4- and 6-cycle free bipartite graphs on m, n Vertices, *Sets, Graphs and Numbers* (Budapest, 1991), *Colloquium Mathematical Society János Bolyai*, vol. 60, North-Holland, Amsterdam, 1992, pp. 135–142.

- [45] R. Canham: A theorem on arrangements of lines in the plane, *Israel J. Math.* 7 (1969), 393–397.
- [46] F. Chung: Subgraphs of a hypercube containing no small even cycles, *J. Graph Theory* 16 (1992), 273–286.
- [47] F. R. K. Chung, and R. L. Graham: *Erdős on Graphs: His Legacy of Unsolved Problems*, A. K. Peters Ltd., Wellesley, MA, 1998.
- [48] C. R. J. Clapham, A. Flockart, and J. Sheehan: Graphs without four-cycles, *J. Graph Theory* 13 (1989), 29–47.
- [49] K. Clarkson, H. Edelsbrunner, L. J. Guibas, M. Sharir, and E. Welzl: Combinatorial complexity bounds for arrangements of curves and spheres, *Discrete Comput. Geom.* 5 (1990), no. 2, 99–160.
- [50] M. Conder: Hexagon-free subgraphs of hypercubes, *J. Graph Theory* 17 (1993), 477–479.
- [51] D. Conlon: An extremal theorem in the hypercube, *Electron. J. Combin.* 17 (2010), Research Paper 111.
- [52] D. Conlon, J. Fox, and B. Sudakov: An approximate version of Sidorenko’s conjecture, *Geom. Funct. Anal.* 20 (2010), no. 6, 1354–1366.
- [53] D. Conlon, J. Fox, and B. Sudakov: Sidorenko’s conjecture for a class of graphs: an exposition, <http://arxiv.org/abs/1209.0184>
- [54] O. Cooley: Proof of the Loebel-Komlós-Sós conjecture for large, dense graphs, *Discrete Math.*, 309 (2009), no. 21, 6190–6228.
- [55] K. Čulík: Teilweise Lösung eines verallgemeinerten Problems von K. Zarankiewicz, *Ann. Polon. Math.* 3 (1956), 165–168.
- [56] D. M. Cvetkovič, M. Doob, and H. Sachs: *Spectra of Graphs*, Academic Press Inc., New York, 1980.
- [57] G. Damásdi, T. Héger, and T. Szőnyi: The Zarankiewicz problem, cages, and geometries, manuscript 2013.
- [58] H. Edelsbrunner and P. Hajnal: A lower bound on the number of unit distances between the vertices of a convex polygon, *J. Combin. Theory Ser. A* 56 (1991), no. 2, 312–316.
- [59] Paul Erdős: Erdős homepage (his scanned in papers up to 1989): www.renyi.hu/~p_erdos.
- [60] P. Erdős: On sequences of integers no one of which divides the product of two others, and some related problems, *Mitt. Forschungsinst. Math. u. Mech. Tomsk* 2 (1938), 74–82.
- [61] P. Erdős: Graph theory and probability I, *Canad. J. Math.* 11 (1959), 34–38.
- [62] P. Erdős: Graph theory and probability II, *Canad. J. Math.* 13 (1961), 346–352.
- [63] P. Erdős: Extremal problems in graph theory, *Proc. Sympos. Smolenice, 1963*, pp. 29–36, *Publ. House Czechoslovak Acad. Sci., Prague*, 1964.

- [64] P. Erdős: Some applications of probability to graph theory and combinatorial problems, *Theory of Graphs and its Applications* (Proc. Sympos. Smolenice, 1963), pp. 133–136, Publ. House Czech. Acad. Sci., Prague, 1964.
- [65] P. Erdős: On some extremal problems in graph theory, *Israel J. Math.* 3 (1965), 113–116.
- [66] P. Erdős: Some recent results on extremal problems in graph theory, *Theory of Graphs* (ed P. Rosenstiehl), (Internat. Sympos., Rome, 1966), Gordon and Breach, New York, and Dunod, Paris, 1967, pp. 117–123.
- [67] P. Erdős: On some new inequalities concerning extremal properties of graphs, *Theory of Graphs* (ed P. Erdős and G. Katona), Academic Press, Nev. York, 1968, pp. 77–81.
- [68] P. Erdős: *The Art of Counting* (ed. J. Spencer), The MIT Press, Cambridge, Mass., 1973.
- [69] P. Erdős: Problems and results on finite and infinite combinatorial analysis, *in Infinite and Finite Sets* (Proc. Conf., Keszthely, Hungary, 1973), pp. 403–424, Proc. Colloq. Math. Soc. J. Bolyai 10, Bolyai–North-Holland, 1975.
- [70] P. Erdős: Some recent progress on extremal problems in graph theory, *Congr. Numerantium* 14 (1975), 3–14.
- [71] P. Erdős: Problems and results in combinatorial analysis, *Colloquio Internazionale sulle Teorie Combinatorie* (Rome, 1973), Tomo II, Atti dei Convegni Lincei, No. 17, pp. 3–17, Accad. Naz. Lincei, Rome, 1976.
- [72] P. Erdős: Problems and results in graph theory and combinatorial analysis, *Graph theory and related topics* (Proc. Conf., Univ. Waterloo, Waterloo, Ont., 1977), pp. 153–163, Academic Press, New York-London.
- [73] P. Erdős: On the combinatorial problems which I would most like to see solved, *Combinatorica* 1 (1981), no. 1, 25–42.
- [74] P. Erdős: On some problems in graph theory, combinatorial analysis and combinatorial number theory, *Graph Theory and Combinatorics* (Cambridge, 1983), pp. 1–17, Academic Press, London, 1984.
- [75] P. Erdős: Two problems in extremal graph theory. *Graphs Combin.* 2 (1986), no. 1, 189–190.
- [76] P. Erdős: On some of my favourite theorems, *Combinatorics, Paul Erdős is eighty*, Vol. 2 (Keszthely, 1993), 97–132, Bolyai Soc. Math. Stud., 2, János Bolyai Math. Soc., Budapest, 1996.
- [77] P. Erdős, R. J. Faudree, J. Pach, and J. Spencer: How to make a graph bipartite, *J. Combin. Theory Ser. B* 45 (1988), no. 1, 86–98.
- [78] P. Erdős, R. J. Faudree, R. H. Schelp, and M. Simonovits: An extremal result for paths, *Graph theory and its applications: East and West* (Jinan, 1986), 155–162, Ann. New York Acad. Sci., 576, New York Acad. Sci., New York, 1989.

- [79] P. Erdős, Z. Füredi, M. Loebel, and V. T. Sós: Discrepancy of trees, *Studia Sci. Math. Hungar.* 30 (1995), no. 1-2, 47–57.
- [80] P. Erdős and T. Gallai: On maximal paths and circuits of graphs, *Acta Math. Acad. Sci. Hungar.* 10 (1959), 337–356.
- [81] P. Erdős, E. Györi, and M. Simonovits: How many edges should be deleted to make a triangle-free graph bipartite? *Sets, graphs and numbers* (Budapest, 1991), pp. 239–263, *Colloq. Math. Soc. János Bolyai*, 60, North-Holland, Amsterdam, 1992.
- [82] P. Erdős, G. Harcos, and J. Pach: Popular distances in 3-space, *Discrete Math.* 200 (1999), no. 1-3, 95–99.
- [83] P. Erdős and L. Moser: Problem 11, *Canad. Math. Bull.* 2 (1959), 43.
- [84] P. Erdős and A. Rényi: On the evolution of random graphs, *Magyar Tud. Akad. Mat. Kutató Int. Közl.* 5 (1960), 17–61.
- [85] P. Erdős and A. Rényi: On a problem in the theory of graphs, *Magyar Tud. Akad. Mat. Kutató Int. Közl.* 7 (1962), 623–641.
- [86] P. Erdős, A. Rényi, and Vera T. Sós: On a problem of graph theory, *Stud Sci. Math. Hung.* 1 (1966), 215–235.
- [87] P. Erdős and H. Sachs: Reguläre Graphen gegebener Tailenweite mit minimaler Knotenzahl (in German), *Wiss. Z. Martin-Luther-Univ. Halle-Wittenberg Math.-Natur. Reihe* 12 (1963), 251–257.
- [88] P. Erdős, A. Sárközy, and V. T. Sós: On product representation of powers, I, *European J. Combinatorics*, 16 (1995), 567–588.
- [89] P. Erdős and M. Simonovits: A limit theorem in graph theory, *Studia Sci. Math. Hungar.* 1 (1966), 51–57.
- [90] P. Erdős and M. Simonovits: Some extremal problems in graph theory, *Combinatorial Theory and Its Applications, I*. (Proc. Colloq. Balatonfüred, 1969), North Holland, Amsterdam, 1970, pp. 377–390.
- [91] P. Erdős and M. Simonovits: An extremal graph problem, *Acta Math. Acad. Sci. Hungar.* 22 (1971/72), 275–282.
- [92] P. Erdős and M. Simonovits: Cube-supersaturated graphs and related problems, *Progress in graph theory* (Waterloo, Ont., 1982), pp. 203–218, Academic Press, Toronto, Ont., 1984.
- [93] P. Erdős and M. Simonovits: Compactness results in extremal graph theory, *Combinatorica* 2 (1982), no. 3, 275–288.
- [94] P. Erdős and M. Simonovits: Supersaturated graphs and hypergraphs, *Combinatorica* 3 (1983), 181–192.
- [95] P. Erdős and A. M. Stone: On the structure of linear graphs, *Bull. Amer. Math. Soc* 52 (1946), 1087–1091.
- [96] G. Fan: Distribution of cycle lengths in graphs, *J. Combin. Theory Ser. B* 84 (2002), 187–202.

- [97] G. Fan, Xuezheng Lv, and Pei Wang: Cycles in 2-connected graphs, *J. Combin. Theory Ser. B* 92 (2004), no. 2, 379–394.
- [98] R. J. Faudree and R. H. Schelp: Path Ramsey numbers in multicolorings, *J. Combin. Theory Ser. B* 19 (1975), no. 2, 150–160.
- [99] R. J. Faudree and M. Simonovits: On a class of degenerate extremal graph problems, *Combinatorica* 3 (1983), 83–93.
- [100] R. J. Faudree and M. Simonovits: On a class of degenerate extremal problems II, preprint.
- [101] J. Fox and B. Sudakov: Dependent random choice, *Random Structures Algorithms* 38 (2011), no. 1-2, 68–99.
- [102] F. A. Firke, P. M. Kosek, E. D. Nash, and J. Williford: Extremal Graphs Without 4-Cycles, <http://arxiv.org/pdf/1201.4912v1.pdf>
- [103] Z. Füredi: Graphs without quadrilaterals, *J. Combin. Theory Ser. B* 34 (1983), 187–190.
- [104] Z. Füredi: Quadrilateral-free graphs with maximum number of edges, preprint 1988, http://www.math.uiuc.edu/~z-furedi/PUBS/furedi_C4from1988.pdf
- [105] Z. Füredi: Graphs of diameter 3 with the minimum number of edges, *Graphs Combin.* 6 (1990), no. 4, 333–337.
- [106] Z. Füredi: The maximum number of unit distances in a convex n -gon, *J. Combin. Theory Ser. A* 55 (1990), no. 2, 316–320.
- [107] Z. Füredi: On a Turán type problem of Erdős, *Combinatorica*, 11 (1991), 75–79.
- [108] Z. Füredi: Turán type problems, in *Surveys in Combinatorics*, London Math. Soc. Lecture Note Ser. 166, Cambridge University Press, Cambridge, UK, 1991, pp. 253–300.
- [109] Z. Füredi: The maximum number of edges in a minimal graph of diameter 2, *J. Graph Theory* 16 (1992), no. 1, 81–98.
- [110] Z. Füredi: Extremal hypergraphs and combinatorial geometry, *Proceedings of the International Congress of Mathematicians*, Vol. 1, 2 (Zürich, 1994), pp. 1343–1352, Birkhäuser, Basel, 1995.
- [111] Z. Füredi: On the number of edges of quadrilateral-free graphs, *J. Combin. Theory Ser. B* 68 (1996), 1–6.
- [112] Z. Füredi: An upper bound on Zarankiewicz problem, *Combin. Probab. Comput.* 5 (1996), no. 1, 29–33.
- [113] Z. Füredi: New asymptotics for bipartite Turán numbers, *J. Combin. Theory Ser. A* 75 (1996), no. 1, 141–144.
- [114] Z. Füredi: On the number of fivecycles, manuscript, unpublished, superseded by [124]
- [115] Z. Füredi: On a theorem of Erdős and Simonovits on graphs not containing

the cube, to appear

- [116] Z. Füredi and Peter Hajnal: Davenport–Schinzel theory of matrices, *Discrete Math.* 103 (1992), 231–251.
- [117] Z. Füredi, A. Naor, and J. Verstraëte: On the Turán number for the hexagon, *Adv. Math.* 203 (2006), no. 2, 476–496.
- [118] Z. Füredi and L. Özkahya: On even-cycle-free subgraphs of the hypercube, *J. Combin. Theory Ser. A* 118 (2011), 1816–1819.
- [119] Z. Füredi, O. Pikhurko, and M. Simonovits: The Turán density of the hypergraph $\{abc, ade, bde, cde\}$, *Electronic J. Combin.* 10 (2003), R18.
- [120] Z. Füredi and M. Simonovits: Triple systems not containing a Fano configuration, *Combin. Probab. Comput.* 14 (2005), no. 4, 467–484.
- [121] Z. Füredi and D. West: Ramsey theory and bandwidth of graphs, *Graphs and Combin.* 17 (2001), 463–471.
- [122] D. K. Garnick, Y. H. H. Kwong, and F. Lazebnik: Extremal graphs without three-cycles or four-cycles, *J. Graph Theory* 17 (1993), no. 5, 633–645.
- [123] D. K. Garnick, and N. A. Nieuwejaar, Non-isomorphic extremal graphs without three-cycles and four-cycles, *J. Combin. Math. Combin. Comput.* 12 (1992), 33–56.
- [124] A. Grzesik: On the maximum number of five-cycles in a triangle-free graph, *J. Combin. Theory Ser. B* 102 (2012), no. 5, 1061–1066.
- [125] J. R. Griggs and Chih-Chang Ho: On the half-half case of the Zarankiewicz problem, *Discrete Math.* 249 (2002), no. 1-3, 95–104.
- [126] J. Griggs, J. Ouyang: $(0, 1)$ -matrices with no half-half submatrix of ones, *European J. Combin.* 18 (1997), 751–761.
- [127] J. R. Griggs, M. Simonovits, and George Rubin Thomas: Extremal graphs with bounded densities of small subgraphs, *J. Graph Theory* 29 (1998), no. 3, 185–207.
- [128] R. K. Guy: A problem of Zarankiewicz, in: *Theory of Graphs (Proc. Colloq., Tihany, 1966)*, pp. 119–150. Academic Press, New York 1968.
- [129] R. K. Guy and S. Znám: A problem of Zarankiewicz, *Recent Progress in Combinatorics (Proc. Third Waterloo Conf. on Combinatorics, 1968)*, pp. 237–243. Academic Press, New York 1969.
- [130] A. Gyárfás: Graphs with k odd cycle lengths, *Discrete Math.* 103 (1992), 41–48.
- [131] A. Gyárfás, J. Komlós and E. Szemerédi: On the distribution of cycle lengths in graphs, *J. Graph Theory* 8 (1984), 441–462.
- [132] A. Gyárfás, C. C. Rousseau, and R. H. Schelp: An extremal problem for paths in bipartite graphs, *J. Graph Theory* 8 (1984), 83–95.
- [133] E. Györi: On the number of C_5 's in a triangle-free graph, *Combinatorica* 9 (1989), 101–102.

- [134] E. Györi: C_6 -free bipartite graphs and product representation of squares, *Graphs Combin.* (Marseille, 1995), *Discrete Math.* 165/166 (1997), 371–375.
- [135] E. Györi: Triangle-free hypergraphs, *Combin. Prob. Comput.* 15 (2006), 185–191.
- [136] E. Györi, B. Rothschild, and A. Ruciński: Every graph is contained in a sparsest possible balanced graph, *Math. Proc. Cambridge Philos. Soc.* 98 (1985), no. 3, 397–401.
- [137] R. Häggkvist and A. D. Scott: Arithmetic progressions of cycles, *Tech. Rep. Mat. Inst. Umeå Univ.* 16, (1998).
- [138] R. Häggkvist and A. Scott: Cycles of nearly equal length in cubic graphs, *Preprint*.
- [139] S. Hartman, J. Mycielski, C. Ryll-Nardzewski: Systèmes spéciaux de points à coordonnées entières, *Colloq. Math.* 3 (1954), 84–85, (Bericht Über di Tagung der Poln Math Gesellschaft, Wroclaw, am 20. September 1951.)
- [140] H. Hatami: Graph norms and Sidorenko’s conjecture, *Israel J. Math.*, 175 (2010), 125–150.
- [141] H. Hatami, J. Hladký, D. Král, S. Norine, and A. Razborov: On the number of pentagons in triangle-free graphs, *J. Combin. Theory Ser. A* 120 (2013), no. 3, 722–732.
- [142] H. Hatami and S. Norine: Undecidability of linear inequalities in graph homomorphism densities, *J. Amer. Math. Soc.* 24 (2011), no. 2, 547–565.
- [143] J. Hladký, J. Komlós, M. Simonovits, M. Stein, and E. Szemerédi: An approximate version of the Loebl-Komlós-Sós Conjecture for sparse graphs, submitted, on arXiv:1211.3050.v1, 2012, Nov 13.
- [144] J. Hladký and D. Piguet: Loebl-Komlós-Sós Conjecture: dense case, *Manuscript* (arXiv:0805:4834).
- [145] M. N. Huxley and H. Iwaniec: Bombieri’s theorem in short intervals, *Mathematika* 22 (1975), 188–194.
- [146] C. Hyltén-Cavallius: On a combinatorial problem, *Colloq. Math.*, 6 (1958), 59–65.
- [147] W. Imrich: Explicit construction of graphs without small cycles, *Combinatorica* 4 (1984), 53–59.
- [148] C. Jagger, P. Štoviček, and A. Thomason: Multiplicities of subgraphs, *Combinatorica* 16 (1996), no. 1, 123–141.
- [149] S. Janson, T. Łuczak, and A. Ruciński: *Random Graphs*, Wiley-Interscience Series in Discrete Mathematics and Optimization. Wiley-Interscience, New York, 2000. xii+333 pp.
- [150] T. Jiang: Compact topological minors in graphs, *J. Graph Theory*, 67 (2011), 139–152.
- [151] T. Jiang and R. Seiver: Turán numbers of subdivided graphs, *SIAM J.*

- Discrete Math. 26 (2012), no. 3, 1238–1255.
- [152] S. Józsa and E. Szemerédi: The number of unit distance on the plane, Infinite and finite sets (Colloq., Keszthely, 1973; dedicated to P. Erdős on his 60th birthday), Vol. II, pp. 939–950. Colloq. Math. Soc. János Bolyai, Vol. 10, North-Holland, Amsterdam, 1975.
 - [153] G. O. H. Katona: Turán’s graph theorem and probability theory, Turán Memorial: Number theory, Analysis and Combinatorics, de Gruyter, Berlin, to appear.
 - [154] Gy. Katona, T. Nemetz, and M. Simonovits: On a problem of Turán in the theory of graphs, *Mat. Lapok* 15 (1964), 228–238.
 - [155] P. Keevash: Hypergraph Turan problems, *Surveys in Combinatorics*, Cambridge University Press, 2011, 83–140.
 - [156] P. Keevash and B. Sudakov: The Turan number of the Fano plane, *Combinatorica* 25 (2005), 561–574.
 - [157] P. Keevash, B. Sudakov and J. Verstraëte: On a conjecture of Erdős and Simonovits: even cycles, *Combinatorica*, to appear.
 - [158] M. Klazar: The Füredi–Hajnal conjecture implies the Stanley–Wilf conjecture, in: D. Krob, A. A. Mikhalev, A. V. Mikhalev (Eds.), *Formal Power Series and Algebraic Combinatorics*, Springer, Berlin, 2000, pp. 250–255.
 - [159] J. Kollár, L. Rónyai, and T. Szabó: Norm graphs and bipartite Turán numbers, *Combinatorica*, 16 (1996), 399–406.
 - [160] J. Komlós and E. Szemerédi: Topological cliques in graphs, *Combin. Probab. Comput.* 3 (1994), no. 2, 247–256.
 - [161] J. Komlós and E. Szemerédi: Topological cliques in graphs II, *Combin. Probab. Comput.* 5 (1996), 79–90.
 - [162] G. N. Kopylov: Maximal paths and cycles in a graph, *Dokl. Akad. Nauk SSSR* 234 (1977), no. 1, 19–21. (English translation: *Soviet Math. Dokl.* 18 (1977), no. 3, 593–596.)
 - [163] A. Kostochka and L. Pyber: Small topological complete subgraphs of “dense” graphs, *Combinatorica*, 8 (1988), 83–86.
 - [164] T. Kóvári, V. T. Sós, and P. Turán: On a problem of K. Zarankiewicz, *Colloq. Math.* 3 (1954), 50–57.
 - [165] D. Kühn and D. Osthus: Four-cycles in graphs without a given even cycle, *J. Graph Theory* 48 (2005), 147–156.
 - [166] T. Lam and J. Verstraëte: A note on graphs without short even cycles, *Electron. J. Combin.* 12 (2005), Note 5, 6 pp.
 - [167] F. Lazebnik and D. Mubayi: New lower bounds for Ramsey numbers of graphs and hypergraphs, *Adv. in Appl. Math.* 28 (2002), no. 3-4, 544–559.
 - [168] F. Lazebnik and V. A. Ustimenko, New examples of graphs without small cycles and of large size, *European J. Combin.* 14 (1993), no.5, 445–460.

- [169] F. Lazebnik, V. A. Ustimenko, and A. J. Woldar: Properties of certain families of $2k$ -cycle-free graphs, *J. Combin. Theory Ser. B* 60 (1994), no. 2, 293–298.
- [170] F. Lazebnik, V. A. Ustimenko, and A. J. Woldar: A new series of dense graphs of high girth, *Bull. Amer. Math. Soc.* 32 (1995), no. 1, 73–79.
- [171] F. Lazebnik, V. A. Ustimenko, and A. J. Woldar, Polarities and $2k$ -cycle-free graphs, *Discrete Math.* 197/198 (1999), 503–513.
- [172] F. Lazebnik and A. J. Woldar: General properties of some families of graphs defined by systems of equations, *J. Graph Theory* 38 (2001), no. 2, 65–86.
- [173] J. Lenz and D. Mubayi: Multicolor Ramsey numbers for complete bipartite versus complete graphs, arXiv 1201.2123, 26 pp.
- [174] B. Lidický, Hong Liu and C. Palmer: On the Turán number of forests, arXiv 1204.3102.
- [175] L. Lovász: Independent sets in critical chromatic graphs, *Studia Sci. Math. Hungar.* 8 (1973), 165–168.
- [176] L. Lovász: *Combinatorial Problems and Exercises*, 2nd Ed., North-Holland, Amsterdam, 1993.
- [177] L. Lovász: *Large Networks and Graph Limits*, Colloquium Publications 2012, 475 pp.
- [178] L. Lovász and M. Simonovits: On the number of complete subgraphs of a graph II, *Studies in Pure Mathematics*, pp. 459–495, (dedicated to the memory of P. Turán), Akadémiai Kiadó and Birkhäuser Verlag 1982.
- [179] A. A. Razborov: Flag algebras, *J. Symbolic Logic* 72 (2007), no. 4, 1239–1282.
- [180] Linyuan Lu, Hexagon-free subgraphs in hypercube Q_n , private communication.
- [181] A. Lubotzky, R. Phillips, and P. Sarnak: Ramanujan graphs, *Combinatorica* 8 (1988), no. 3, 261–277.
- [182] A. McLennan: The Erdős-Sós conjecture for trees of diameter four, *J. Graph Theory* 49 (2005), no. 4, 291–301.
- [183] W. Mader: Homomorphieeigenschaften und mittlere Kantendichte von Graphen, *Math. Ann.*, 174 (1967), 265–268.
- [184] W. Mader: Topological subgraphs in graphs of large girth, *Combinatorica* 18 (1998), no. 3, 405–412.
- [185] W. Mader: Topological minors in graphs of minimum degree n , *Contemporary trends in discrete mathematics (Štířín Castle, 1997)*, 199–211, DIMACS Ser. Discrete Math. Theoret. Comput. Sci., 49, Amer. Math. Soc., Providence, RI, 1999.
- [186] W. Mader: Graphs with $3n - 6$ edges not containing a subdivision of K_5 , *Combinatorica* 25 (2005), no. 4, 425–438.

- [187] A. Marcus and G. Tardos: Excluded permutation matrices and the Stanley-Wilf conjecture, *J. Combin. Theory Ser. A* 107 (2004), no. 1, 153–160.
- [188] G. A. Margulis: Explicit construction of graphs without short cycles and low density codes, *Combinatorica* 2 (1982), 71–78.
- [189] G. A. Margulis: Arithmetic groups and graphs without short cycles, in: 6th Int. Symp. on Information Theory, Tashkent, Abstracts 1, 1984, pp. 123–125 (in Russian).
- [190] G. A. Margulis: Explicit group-theoretical construction of combinatorial schemes and their application to the design of expanders and concentrators, *J. Problems of Inform. Trans.* 24 (1988), 39–46; translation from *Problemy Peredachi Informatsii* 24 (January–March 1988), 51–60.
- [191] G. Megyesi and E. Szabó: On the tacnodes of configurations of conics in the projective plane, *Math. Ann.* 305 (1996), no. 4, 693–703.
- [192] M. Molloy and B. Reed: *Graph Colouring and the Probabilistic Method, Algorithms and Combinatorics*, 23. Springer-Verlag, Berlin, 2002. xiv+326 pp.
- [193] B. Montágh: Unavoidable substructures, PHD Thesis, University of Memphis, May 2005.
- [194] M. Mörs: A new result on the problem of Zarankiewicz, *J. Combin. Theory Ser. A* 31 (1981), no. 2, 126–130.
- [195] D. Mubayi and Gy. Turán: Finding bipartite subgraphs efficiently, *Inform. Process. Lett.* 110 (2010), no. 5, 174–177.
- [196] Z. L. Nagy: A multipartite version of the Turán problem-density conditions and eigenvalues, *Electron. J. Combin.* 18 (2011), no. 1, Paper 46, 15 pp.
- [197] V. Nikiforov: Bounds on graph eigenvalues II, *Linear Algebra Appl.* 427 (2007), 183–189.
- [198] V. Nikiforov: A contribution to the Zarankiewicz problem, *Linear Algebra Appl.* 432 (2010), no. 6, 1405–1411.
- [199] J. Pach and P. K. Agarwal: *Combinatorial Geometry*, Wiley-Interscience, New York, 1995. xiv+354 pp.
- [200] P. P. Pálffy and M. Szalay, in *Turán Memorial: Number theory, Analysis and Combinatorics*, de Gruyter, Berlin, to appear.
- [201] D. Piguet and M. J. Stein: Loebel-Komlós-Sós conjecture for trees of diameter 5, *Electron. J. Combin.*, 15 (2008), Research Paper 106, 11 pp.
- [202] D. Piguet and M. J. Stein: An approximate version of the Loebel-Komlós-Sós conjecture, *J. Combin. Theory Ser. B* 102 (2012), no. 1, 102–125.
- [203] O. Pikhurko: A note on the Turán Function of even cycles, *Proc. Amer. Math Soc.* 140 (2012), 3687–3992.
- [204] R. Pinchasi and M. Sharir: On graphs that do not contain the cube and related problems, *Combinatorica* 25 (2005), no. 5, 615–623.

- [205] C. Reiher: The clique density theorem, arxiv1212.2454.
- [206] I. Reiman: Über ein Problem von K. Zarankiewicz, *Acta Math. Acad. Sci. Hungar.* 9 (1958), no. 3-4, 269–273.
- [207] I. Reiman: An extremal problem in graph theory, (Hungarian). *Mat. Lapok* 12 (1961), 44–53.
- [208] A. Rényi: Selected Papers of Alfréd Rényi, Akadémiai Kiadó, 1976 (ed. Paul Turán).
- [209] V. Rödl, and M. Schacht: Extremal results for random graphs, in this volume.
- [210] I. Z. Ruzsa and E. Szemerédi: Triple systems with no six points carrying three triangles, *Combinatorics (Proc. Fifth Hungarian Colloq., Keszthely, 1976)*, Vol. II, pp. 939–945, *Colloq. Math. Soc. János Bolyai*, 18, North-Holland, Amsterdam-New York, 1978.
- [211] J.-F. Saclé and M. Woźniak: A note on the Erdős-Sós conjecture for graphs without C_4 , *J. Combin. Theory Ser. B* 70 (1997), no.2, 367–372.
- [212] G. N. Sárközy: Cycles in bipartite graphs and an application in number theory, *J. Graph Theory*, 19 (1995), 323–331.
- [213] A. Scott: Szemerédi’s regularity lemma for matrices and sparse graphs, *Combin. Probab. Comput.* 20 (2011), no. 3, 455–466.
- [214] Jian Shen: On two Turán numbers, *J. Graph Theory* 51 (2006), 244–250.
- [215] A. F. Sidorenko: Asymptotic solution for a new class of forbidden r -graphs, *Combinatorica* 9 (1989), no. 2, 207–215.
- [216] A. Sidorenko: A correlation inequality for bipartite graphs, *Graphs Combin.* 9 (1993), no. 2, 201–204.
- [217] A. F. Sidorenko: What do we know and what we do not know about Turán Numbers, *Graphs Combin.* 11 (1995), no. 2, 179–199.
- [218] M. Simonovits: A method for solving extremal problems in graph theory, *Theory of Graphs, Proc. Colloq. Tihany, (1966)*, (Ed. P. Erdős and G. Katona), pp. 279–319, Acad. Press, New York, 1968.
- [219] M. Simonovits: On colour-critical graphs, *Studia Sci. Math. Hungar.* 7 (1972), 67–81.
- [220] M. Simonovits: Note on a hypergraph extremal problem, *Hypergraph Seminar, Columbus Ohio USA, 1972*, (ed. C. Berge and D. K. Ray-Chaudury), *Lecture Notes in Mathematics* 411, pp. 147–151, Springer Verlag, 1974.
- [221] M. Simonovits: Extremal graph problems with symmetrical extremal graphs, additional chromatic conditions, *Discrete Math.* 7 (1974), 349–376.
- [222] M. Simonovits: On Paul Turán’s influence on graph theory, *J. Graph Theory* 1 (1977), no. 2, 102–116.
- [223] M. Simonovits: Extremal graph problems and graph products, *Studies in Pure Mathematics*, pp. 669–680, (dedicated to the memory of P. Turán),

Akadémiai Kiadó and Birkhäuser Verlag 1982.

- [224] M. Simonovits: Extremal graph theory, in: L. W. Beineke, R. J. Wilson (Eds.), *Selected Topics in Graph Theory II.*, pp. 161–200, Academic Press, London, 1983.
- [225] M. Simonovits: Extremal graph problems, degenerate extremal problems and supersaturated graphs, *Progress in graph Theory*, (Bondy and Murty, Eds.), pp. 419–438, Academic Press, 1984.
- [226] M. Simonovits: How to solve a Turán type extremal graph problem? (linear decomposition), *Contemporary trends in discrete mathematics* (Stirin Castle, 1997), pp. 283–305, DIMACS Ser. Discrete Math. Theoret. Comput. Sci., 49, Amer. Math. Soc., Providence, RI, 1999.
- [227] M. Simonovits: Paul Erdős’ influence on extremal graph theory, *The mathematics of Paul Erdős, II.*, pp. 148–192, *Algorithms Combin.*, 14, Springer, Berlin, 1997.
- [228] M. Simonovits: Paul Erdős’ Influence on Extremal Graph Theory, The new version of the old paper [227].
- [229] M. Simonovits: Paul Turán’s influence in Combinatorics, in *Turán Memorial: Number Theory, Analysis, and Combinations*, De Gruyter, to appear.
- [230] M. Simonovits and V. T. Sós: Ramsey-Turán theory, *Combinatorics, graph theory, algorithms and applications*, *Discrete Math.* 229 (2001), no. 1-3, 293–340.
- [231] R. R. Singleton: On minimal graphs of maximum even girth, *J. Combinatorial Theory* 1 (1966), 306–332.
- [232] V. T. Sós: Remarks on the connection of graph theory, finite geometry and block designs, *Colloquio Internazionale sulle Teorie Combinatorie* (Roma, 1973), Tomo II, pp. 223–233, *Atti dei Convegni Lincei*, No. 17, Accad. Naz. Lincei, Rome, 1976.
- [233] J. Spencer, E. Szemerédi, and W. T. Trotter: Unit distances in the Euclidean plane, *Graph theory and combinatorics* (Cambridge, 1983), pp. 293–303, Academic Press, London, 1984.
- [234] B. Sudakov and J. Verstraëte: Cycle lengths in sparse graphs, *Combinatorica* 28 (2008), no. 3, 357–372.
- [235] E. Szemerédi: Regular partitions of graphs, *Problemes Combinatoires et Theorie des Graphes* (ed. I.-C. Bermond et al.), pp. 399–401, CNRS, Paris, 1978.
- [236] G. Tardos: On 0-1 matrices and small excluded submatrices, *J. Combin. Th. Ser. A* 111 (2005), 266–288.
- [237] A. G. Thomason: A disproof of a conjecture of Erdős in Ramsey Theory, *J. London Math. Soc.* 39 (1989), 246–255.
- [238] A. Thomason and P. Wagner: Bounding the size of square-free subgraphs

- of the hypercube, *Discrete Math.* 309 (2009), 1730–1735.
- [239] C. M. Timmons: Ordered Turán Problems, Lecture no. 1086-05-1067 on the Joint Mathematics Meetings, San Diego, CA, January 9, 2013.
- [240] B. Toft: Two theorems on critical 4-chromatic graphs, *Studia Sci. Math. Hungar.* 7 (1972), 83–89.
- [241] P. Turán: On a theorem of Hardy-Ramanujan, *Journal of London Math Soc.* 9 (1934), 274–276.
- [242] P. Turán: On an extremal problem in graph theory, (Hungarian), *Mat. Fiz. Lapok* 48 (1941), 436–452.
- [243] P. Turán: On the theory of graphs, *Colloq. Math.* 3 (1954), 19–30.
- [244] P. Turán: A note of welcome, *J. Graph Theory* 1 (1977), 7–9.
- [245] P. Valtr: Strictly convex norms allowing many unit distances and related touching questions, manuscript.
- [246] J. Verstraëte: On arithmetic progressions of cycle lengths in graphs, *Combin. Probab. Comput.* 9 (2000), no.4, 369–373.
- [247] R. Wenger: Extremal graphs with no C_4 's, C_6 's, or C_{10} 's, *J. Combin. Theory Ser. B* 52 (1991), no. 1, 113–116.
- [248] R. M. Wilson: An existence theory for pairwise balanced designs, III. Proof of the existence conjectures, *J. Combin. Theory Ser. A* 18 (1975), 71–79.
- [249] D. R. Woodall: Maximal circuits of graphs I, *Acta Math. Acad. Sci. Hungar.* 28 (1976), no. 1–2, 77–80.
- [250] D. R. Woodall: Maximal circuits of graphs II, *Studia Sci. Math. Hungar.* 10 (1975), no. 1–2, 103–109.
- [251] M. Woźniak: On the Erdős–Sós conjecture, *J. Graph Theory*, 21 (1996), no. 2, 229–234.
- [252] Y. Yuansheng and P. Rowlinson: On graphs without 6-cycles and related Ramsey numbers, *Utilitas Math.* 44 (1993), 192–196.
- [253] Y. Yuansheng and P. Rowlinson: On extremal graphs without four-cycles, *Utilitas Math.* 41 (1992), 204–210.
- [254] K. Zarankiewicz: Problem 101, *Colloquium Mathematicum* 2 (1951), p. 301.
- [255] Yi Zhao: Proof of the $(n/2 - n/2 - n/2)$ conjecture for large n , *Electron. J. Combin.* 18 (2011), Paper 27.
- [256] Š. Znám: On a combinatorial problem of K. Zarankiewicz, *Colloq. Math.* 11 (1963), 81–84.
- [257] Š. Znám: Two improvements of a result concerning a problem of K. Zarankiewicz, *Colloq. Math.* 13 (1964/1965), 255–258.