

On the largest prime factors of n and $n + 1$

PAUL ERDÖS AND CARL POMERANCE

§1. Introduction

If $n \geq 2$ is an integer, let $P(n)$ denote the largest prime factor of n . For every $x > 0$ and every t , $0 \leq t \leq 1$, let $A(x, t)$ denote the number of $n \leq x$ with $P(n) \geq x^t$. A well-known result due to Dickman [4] and others is

THEOREM A. *The function*

$$a(t) = \lim_{x \rightarrow \infty} x^{-1} A(x, t)$$

is defined and continuous on $[0, 1]$.

In fact it is even shown that $a(t)$ is strictly decreasing and differentiable. Note that $a(0) = 1$ and $a(1) = 0$.

If $0 \leq t, s \leq 1$, denote by $B(x, t, s)$ the number of $n \leq x$ with $P(n) \geq x^t$ and $P(n+1) \geq x^s$. One might guess that

$$b(t, s) = \lim_{x \rightarrow \infty} x^{-1} B(x, t, s)$$

exists and is continuous on $[0, 1]^2$. In fact, one could guess that

$$b(t, s) = a(t)a(s);$$

that is, the largest prime factors of n and $n + 1$ are “independent events.” We do not know how to prove the above guesses. In fact, we cannot even prove the almost certain truth that the density of integers n with $P(n) > P(n + 1)$ is $\frac{1}{2}$.

However we can prove:

THEOREM 1. *For each $\epsilon > 0$, there is a $\delta > 0$ such that for sufficiently large x ,*

AMS (1970) subject classification: Primary 10H15, 10J15.

Received May 19, 1976.

the number of $n \leq x$ with

$$x^{-\delta} < P(n)/P(n+1) < x^{\delta} \quad (1)$$

is less than ϵx .

That is, $P(n)$ and $P(n+1)$ are usually not close. We use Brun's method in the proof. One corollary is that the lower density of integers n for which $P(n) > P(n+1)$ is positive (see §6).

If the canonical prime factorization of $n > 1$ is $\prod p_i^{a_i}$, let $f(n) = \sum a_i p_i$; and let $f(1) = 0$. Several authors have considered this function or the closely related $g(n) = \sum p_i$ or $h(n) = \sum p_i^{a_i}$, among them Alladi and Erdős [1], Chawla [2], Dane [3], Hall [7], Lal [10], LeVan [12], and Nicolas [14]. In Nelson, Penney, and Pomerance [13] the following problem is raised: does the set of n for which $f(n) = f(n+1)$ have density 0? If $f(n) = f(n+1)$, we call n an *Aaron number* (see [13]). We prove here the Aaron numbers do indeed have density 0. The result follows as a corollary to Theorem 1 and

THEOREM 2. *For every $\epsilon > 0$, there is a $\delta > 0$ such that for sufficiently large x there are at least $(1 - \epsilon)x$ choices for $n \leq x$ such that*

$$P(n) < f(n) < (1 + x^{-\delta})P(n). \quad (2)$$

Theorem 2 implies that usually $f(n) \approx P(n)$ and $f(n+1) \approx P(n+1)$. But Theorem 1 implies $P(n)$ and $P(n+1)$ are usually not close. Hence $f(n)$ and $f(n+1)$ are usually not close, and in particular, we usually have $f(n) \neq f(n+1)$. This then establishes that the Aaron numbers have density 0. However we can prove a sharper result:

THEOREM 3. *For every $\epsilon > 0$, the number of $n \leq x$ for which $f(n) = f(n+1)$ is $O(x/(\log x)^{1-\epsilon})$.*

Actually we can prove the sharper estimate $O(x/\log x)$, but the proof is more difficult than the proof of Theorem 3 and we do not present it here. We suspect that the estimate $O(x/(\log x)^k)$ is true for every k , but we cannot prove this for any $k > 1$. In fact, we cannot even get $o(x/\log x)$. On the other hand, we cannot prove that there are infinitely many Aaron numbers (this would follow if Schinzel's Conjecture *H* is true – see [13]). But by a consideration of those n for which $P(n)$ and $P(n+1)$ are both relatively small, we believe the number of Aaron numbers up to x is $\Omega(x^{1-\epsilon})$ for every $\epsilon > 0$.

There are integers n for which $f(n) = f(n+1) = f(n+2)$. The least example, kindly found for us by David E. Penney in a computer search, is $n = 417162$. We cannot prove that the number of such $n \leq x$ is $o(x/\log x)$. We conjecture that for every k there are integers n with $f(n) = f(n+1) = \dots = f(n+k)$.

§2. Preliminaries

In this section we record several lemmas which will be useful in our discussion. The letter p denotes a prime.

LEMMA 1. *There is an absolute constant C , such that if $3 < u < v$, then*

$$\sum_{u \leq p \leq v} \frac{1}{p} < \frac{C + \log(v/u)}{\log u}.$$

This lemma is used when u is large compared with v/u . The proof follows easily from the classical result (see Hardy and Wright [8], Theorem 427 and its proof): there are absolute constants B, D such that if $x \geq 3$, then

$$\left| \sum_{p \leq x} \frac{1}{p} - \log \log x - B \right| < \frac{D}{\log x}.$$

Lemma 1 easily follows with $C = 2D$.

$$\text{LEMMA 2. } \sum_{p \geq t} \frac{1}{p \log p} \sim \frac{1}{\log t}.$$

Proof. If p_k denotes the k -th prime, then $p_k \sim k \log k$ and

$$\sum_{p \geq t} \frac{1}{p \log p} \sim \sum_{k \geq \pi(t)} \frac{1}{k \log^2 k} \sim \frac{1}{\log \pi(t)} \sim \frac{1}{\log t}.$$

LEMMA 3. *If $P(n) \geq 5$, then $f(n) \leq P(n) \log n / \log P(n)$.*

Proof. We use the fact that $t/\log t$ is increasing for $t \geq e$ and $2/\log 2 < 5/\log 5$. Write $n = \prod p_i^{a_i}$ where $p_1 = P(n)$. Then

$$f(n) = \sum a_i p_i \leq \sum a_i p_1 \log p_i / \log p_1 = P(n) \log n / \log P(n).$$

§3. Proof of Theorem 1

Let $\epsilon > 0$. From Theorem A it follows there is a $\delta_0 = \delta_0(\epsilon)$ such that $\frac{1}{4} > \delta_0 > 0$ and for large x the number of $n \leq x$ with

$$P(n) < x^{\delta_0} \quad \text{or} \quad x^{1/2 - \delta_0} \leq P(n) < x^{1/2 + \delta_0}$$

is less than $\epsilon x/3$. We now consider the remaining $n \leq x$. There are 2 cases:

- (i) $x^{\delta_0} \leq P(n) < x^{1/2 - \delta_0}$,
- (ii) $x^{1/2 + \delta_0} \leq P(n)$.

For each pair of primes p, q , the number of $n \leq x$ for which $P(n) = p$, $P(n+1) = q$ is at most $1 + [x/pq]$. Then for large x , the number of $n \leq x$ in case (i) for which (1) holds is at most (assume $0 < \delta < \delta_0/4$)

$$\begin{aligned} \sum_{\substack{x^{\delta_0} \leq p < x^{1/2 - \delta_0} \\ px^{-\delta} < q < px^{\delta}}} 1 + [x/pq] &< x^{1 - 2\delta_0 + \delta} + x \sum \frac{1}{p} \sum \frac{1}{q} \\ &< x^{1 - 2\delta_0 + \delta} + x \sum \frac{1}{p} \cdot \frac{C + \log(x^{2\delta})}{\log(px^{-\delta})} \quad (\text{Lemma 1}) \\ &< x^{1 - 2\delta_0 + \delta} + 3\delta x \log x \sum \frac{1}{p \log p} \\ &< x^{1 - 2\delta_0 + \delta} + 4\delta x / \delta_0 \quad (\text{Lemma 2}) \end{aligned} \tag{3}$$

Hence if we choose δ so that

$$0 < \delta < \delta_0 \epsilon / 13, \tag{4}$$

then (3) implies there are fewer than $\epsilon x/3$ choices of such n .

Suppose now $n \leq x$ is in case (ii) and (1) holds. Let $a = n/P(n)$, $b = (n+1)/P(n+1)$. Then $a \leq x^{1/2 - \delta_0}$, $b < x^{1/2 - \delta_0 + \delta}$, and $x^{-\delta}/2 < a/b < 2x^{\delta}$. On the other hand, given integers a, b , the number of $n \leq x$ for which $n = aP(n)$ and $n+1 = bP(n+1)$ is at most the number of primes $p \leq x/a$ such that $(ap+1)/b$ is prime. (Note that there is at most one such prime p unless $(a, b) = 1$ and $2 \mid ab$.) All such primes p are in a fixed residue class mod b , say $p = kb + c$ for some $k \geq 0$. Let $d = (ac+1)/b$. Then we are counting integers k with $0 \leq k < x/ab$ such that $kb + c$ and $ka + d$ are simultaneously prime. By Brun's method (see Halberstam

and Richert [6], Theorem 2.3, p. 70), we have the number of such k is at most

$$\frac{Ax}{ab \log^2(x/ab)} \prod_{p|ab} \left(1 - \frac{1}{p}\right)^{-1} = \frac{Ax}{\varphi(a)\varphi(b) \log^2(x/ab)}$$

where A is an absolute constant (independent of the choice of a, b) and φ is Euler's function. Hence for sufficiently large x , the number of $n \leq x$ in case (ii) for which (1) holds is at most

$$Ax \sum_{\substack{1/2 - \delta_0 \\ a \leq x \\ ax^{2\delta_0} < b < 2ax^{\delta_0}}} 1/\varphi(a)\varphi(b) \log^2(x/ab) \quad (5)$$

$$< \frac{2Ax}{(2\delta_0 - \delta)^2 \log^2 x} \sum \frac{1}{\varphi(a)} \sum \frac{1}{\varphi(b)}.$$

We now use the result of Landau [11], that if $E = \zeta(2)\zeta(3)/\zeta(6)$, then

$$\sum_{n \leq x} 1/\varphi(n) = E \log x + o(1).$$

Hence for large x the quantity in (5) is less than

$$\begin{aligned} & \frac{3EAx}{(2\delta_0 - \delta)^2 \log^2 x} \sum \frac{\log(x^{2\delta_0})}{\varphi(a)} \\ &= \frac{6\delta EAx}{(2\delta_0 - \delta)^2 \log x} \sum \frac{1}{\varphi(a)} \\ &< \frac{7\delta E^2 Ax}{(2\delta_0 - \delta)^2 \log x} \log(x^{1/2 - \delta_0}) \\ &< \frac{4\delta E^2 Ax}{(2\delta_0 - \delta)^2}. \end{aligned} \quad (6)$$

If we now choose δ so that

$$0 < \delta < \delta_0^2 \epsilon / 4E^2 A \quad \text{and} \quad \delta < \delta_0/4, \quad (7)$$

then (6) implies there are fewer than $\epsilon x/3$ choices for such n . Hence if we choose δ so that (4) and (7) hold, it follows that the number of $n \leq x$ for which (1) holds is

less than ϵx for every sufficiently large value of x (depending, of course, on ϵ). This completes our proof.

Note that using a known explicit estimate for the upper bound sieve result we may take $A = 8 + o_x(1)$.

§4. The proof of Theorem 2

Since any integer $n \leq x$ is divisible by at most $\log x / \log 2$ primes, we have for large x and composite $n \leq x$

$$\begin{aligned} f(n) &= P(n) + f(n/P(n)) \leq P(n) + P(n/p(n)) \log x / \log 2 \\ &< P(n) + P(n/P(n))x^\delta. \end{aligned} \quad (8)$$

If (2) fails, then, but for $o(x)$ choices of $n \leq x$, we have

$$f(n) \geq (1 + x^{-\delta})P(n), \quad (9)$$

so that from (8) and (9) we have

$$P(n/P(n)) > x^{-2\delta}P(n). \quad (10)$$

Let $\epsilon > 0$. From Theorem A there is a $\delta_0 = \delta_0(\epsilon) > 0$ such that for large x , the number of $n \leq x$ with $P(n) < x^{\delta_0}$ is at most $\epsilon x / 3$. For each pair of primes p, q the number of $n \leq x$ with $P(n) = p$ and $P(n/P(n)) = q$ is at most $[x/pq]$. Hence from (10), for large x the number of $n \leq x$ for which (2) fails is at most (assume $0 < \delta < \delta_0/7$)

$$o(x) + \epsilon x / 3 + \sum_{\substack{x^{\delta_0} \leq p \\ x^{-2\delta} p < q \leq p}} [x/pq] < \epsilon x / 2 + x \sum \frac{1}{p} \sum \frac{1}{q}$$

$$< \epsilon x / 2 + x \sum \frac{1}{p} \cdot \frac{C + \log(x^{2\delta})}{\log(x^{-2\delta}p)} \quad (\text{Lemma 1})$$

$$< \epsilon x / 2 + 3\delta x \log x \sum \frac{1}{p \log p}$$

$$< \epsilon x / 2 + 4\delta x / \delta_0 \quad (\text{Lemma 2})$$

$$\leq \epsilon x,$$

if we take $\delta = \delta_0 \epsilon / 8$. This completes the proof.

§5. Aaron numbers

In this section we prove Theorem 3. Let x be large, $n \leq x$, and $f(n) = f(n+1)$. We distinguish two cases:

- (i) $P(n) > x^{1/2}$,
- (ii) $P(n) \leq x^{1/2}$.

Let n be in case (i). We first show that

$$P(n+1) > P(n)/3. \quad (11)$$

Indeed we have

$$x^{1/2} < P(n) \leq f(n) = f(n+1) \leq P(n+1) \log(x+1)/\log 2$$

so that $P(n+1) > x^{1/2} \log 2 / \log(x+1)$. Hence Lemma 3 implies

$$P(n) < P(n+1) \log(x+1) / \log(x^{1/2} \log 2 / \log(x+1)) < 3P(n+1)$$

for large x , which proves (11). We next show that

$$|P(n) - P(n+1)| < 4x/P(n). \quad (12)$$

Indeed, $f(n) = f(n+1)$ implies

$$\begin{aligned} P(n+1) - P(n) &= f(n/P(n)) - f((n+1)/P(n+1)) \leq n/P(n), \\ P(n) - P(n+1) &\leq (n+1)/P(n+1), \end{aligned}$$

so that using (11) we have (12). We next show that

$$P(n) < 3x^{2/3}. \quad (13)$$

We use the congruence

$$(P(n+1) - P(n)) \frac{n+1}{P(n+1)} \equiv 1 \pmod{P(n)}. \quad (14)$$

From (11) we have $P(n)$ and $P(n+1)$ both odd primes so the left side of (14) is

not 1. Then (11), (12), and (14) imply

$$\begin{aligned}
 P(n) &\leq |P(n) - P(n+1)| \frac{n+1}{P(n+1)} + 1 < \frac{4x}{P(n)} \cdot \frac{x+1}{P(n+1)} + 1 \\
 &< \frac{12x(x+1)}{P(n)^2} + 1 < \frac{14x^2}{P(n)^2}
 \end{aligned}$$

for large x . so that (13) follows.

If p, q are primes with $x^{1/2} < p, q > p/3$, then there are at most 3 integers $n \leq x$ with $P(n) = p$ and $P(n+1) = q$. Hence from (11), (12), (13) we have for large x that the number of $n \leq x$ in case (i) for which $f(n) = f(n+1)$ is at most

$$\begin{aligned}
 3 \sum_{\substack{x^{1/2} < p < 3x^{2/3} \\ |p-q| < \epsilon x/p}} 1 &\ll \sum_{x^{1/2} < p < 3x^{2/3}} \frac{x/p}{\log(x/p)} \\
 &\ll \sum \frac{x}{p \log x} \ll \frac{x}{\log x},
 \end{aligned}$$

where we use the well-known result of Hardy and Littlewood (see [9], p. 66) for the number of primes in an interval and Lemma 1.

We now turn our attention to case (ii). We have (see Erdős [5], proof of Lemma 1 or Rankin [15], Lemma II) the number of $n \leq x$ for which we do not have

$$P(n) > x^{1/3 \log \log x} \tag{16}$$

is $O(x/\log x)$. So we may assume (16) holds. Then using Lemma 3 and the argument which establishes (11), we have from the equation $f(n) = f(n+1)$ that

$$P(n)/4 \log \log x < P(n+1) < 3P(n) \log \log x. \tag{17}$$

For each pair of primes p, q , there are at most $1 + [x/pq]$ integers $n \leq x$ with $P(n) = p$ and $P(n+1) = q$. Hence from (16) and (17), for large x the number of $n \leq x$ in case (ii) for which $f(n) = f(n+1)$ is at most

$$\begin{aligned}
 \sum_{\substack{x^{1/3 \log \log x} < p \leq x^{1/2} \\ p/4 \log \log x < q < 3p \log \log x}} 1 + [x/pq] &\leq \pi(x^{1/2}) \pi(3x^{1/2} \log \log x) + x \sum \frac{1}{p} \sum \frac{1}{q} \\
 &\ll \frac{x}{\log x} + x \sum \frac{1}{p} \cdot \frac{\log \log \log x}{\log p} \tag{Lemma 1}
 \end{aligned}$$

$$\ll \frac{x \log \log x \log \log \log x}{\log x}. \quad (\text{Lemma 2})$$

This completes the proof of Theorem 3.

§6. The probability that $P(n) > P(n + 1)$.

Using some computer estimates of the function $a(t)$ made with the generous assistance of Don R. Wilhelmsen, it can be shown that the number of integers $n \leq x$ such that

$$x^{0.31} \leq P(n) < x^{0.46} \quad (18)$$

is more than $0.2002x$ for sufficiently large x . By an elementary argument similar to the proof of case (i) in Theorem 1 (see §3) one can show the number of $n \leq x$ for which (18) holds and for which

$$P(n) < P(n + 1) < P(n)x^{0.08} \quad (19)$$

is less than $0.0763x$ for sufficiently large x . Hence the number of $n \leq x$ for which (19) fails is more than

$$0.2002x - 0.0763x = 0.1239x$$

for sufficiently large x . Now for every k choices of $n \leq x$ for which $P(n + 1) \geq P(n)x^{0.08}$, there must be at least $[0.08k]$ integers n in the same interval for which $P(n) > P(n + 1)$. Hence the lower density of integers n for which $P(n) > P(n + 1)$ is at least

$$(0.08) \cdot (0.1239) > 0.0099.$$

Note that the same is true for integers n for which $P(n) < P(n + 1)$. Undoubtedly improvements in this type of result are possible.

§7. Comments on three or more consecutive numbers.

It is easy to show that the patterns

$$P(n) < P(n + 1), P(n + 1) > P(n + 2);$$

$$P(n) > P(n + 1), P(n + 1) < P(n + 2),$$

both occur infinitely often. However we cannot prove either of these two patterns occurs for a positive density of n , although this certainly must be the case. Suppose now p is an odd prime and

$$k_0 = \inf \{k : P(p^{2^k} + 1) > p\}$$

(note that $P(p^{2^{k_0}} + 1) \equiv 1 \pmod{2^{k_0+1}}$, so $k_0 < \infty$). Then

$$P(p^{2^{k_0}} - 1) < P(p^{2^{k_0}}) < P(p^{2^{k_0}} + 1).$$

On the other hand, we cannot find infinitely many n for which

$$P(n) > P(n+1) > P(n+2), \tag{20}$$

but perhaps we overlook a simple proof.

Suppose now

$$\epsilon_n = \begin{cases} 1, & \text{if } P(n) > P(n+1), \\ 0, & \text{if } P(n) < P(n+1). \end{cases}$$

Then $\sum_{n=2}^{\infty} \epsilon_n/2^n$ is irrational. Indeed, suppose not, so that $\{\epsilon_n\}$ is eventually periodic with period length K . Let $p > K$ be a fixed prime. An old and well-known result of Pólya implies that there are only finitely many pairs of consecutive integers in the set $M = \{n : P(n) \leq p\}$. (In fact, from the work of Baker, the largest consecutive pair in M is effectively computable.) Note that $p^i, 2p^i, \dots, Kp^i$ are all in M for every i . Hence for large i , none of $p^i + 1, 2p^i + 1, \dots, Kp^i + 1$ is in M , so that $\epsilon_m = 0$ for $m = p^i, 2p^i, \dots, Kp^i$. But these numbers form a complete residue system mod K . Hence $\epsilon_n = 0$ for every large n , an absurdity.

For each k , let $h(k)$ denote the number of different patterns of k consecutive terms of $\{\epsilon_n\}$ which occur infinitely often. Surely we must have $h(k) = 2^k$. This is easy for $k = 1$, but already for $k = 2$, all we can prove is $h(2) \geq 3$. (If there are infinitely many n for which (20) holds, then $h(2) = 4$.) It follows from the non-periodicity of $\{\epsilon_n\}$ that for every k ,

$$h(k) \geq k + 1.$$

To see this, it is sufficient to show $h(k)$ is strictly increasing (since $h(1) = 2$). But if $h(k) = h(k+1)$ (clearly $h(k) > h(k+1)$ is impossible), then sufficiently far out in the sequence $\{\epsilon_n\}$ we have each term determined by the previous k terms. Then as soon as a k -tuple repeats, the sequence repeats and hence is periodic.

We remark that $h(k) = 2^k$ can be seen to follow from the prime k -tuples conjecture.

REFERENCES

- [1] ALLADI, K. and ERDŐS, P., *On an additive arithmetic function*. To appear.
- [2] CHAWLA, L. M., *On a pair of arithmetic functions*. J. Natur. Sci. and Math. 8 (1968), 263–269.
- [3] DANE, P., *The "prime derivative"*. J. Recreational Math. 7 (1974), 111–115.
- [4] DICKMAN, K., *On the frequency of numbers containing prime factors of a certain relative magnitude*. Ark. Mat. Astr. Fys. 22 (1930), no. A10.
- [5] ERDŐS, P., *On the density of the abundant numbers*. J. London Math. Soc. 9 (1934), 278–282.
- [6] HALBERSTAM, H. and RICHERT, H.-E., *Sieve Methods*. Academic Press, London, 1974.
- [7] HALL, R. R., *On the probability that n and $f(n)$ are relatively prime*. Acta Arith. 17 (1970), 169–183.
- [8] HARDY, G. H. and WRIGHT, E. M., *An Introduction to the Theory of Numbers*. Fourth Edition, Oxford, 1968.
- [9] HARDY, G. H., and LITTLEWOOD, J. E., *Some problems of "partitio numerorum", III: On the expression of a number as a sum of primes*. Acta Math. 44 (1923), 1–70.
- [10] LAL, M., *Iterates of a number-theoretic function*. Math. Comp. 23 (1969), 181–183.
- [11] LANDAU, E., *Ueber die zahlentheoretische Function $\varphi(n)$ und ihre Beziehung zum Goldbachschen Satz*. Nachr. Königl. Ges. Wiss. Göttingen, math.-phys. Kl. (1900), 177–186.
- [12] LEVAN, M. O., *On the generalization of Chawla's two arithmetic functions*. J. Natur. Sci. and Math. 9 (1969), 57–66.
- [13] NELSON, C., PENNEY, D. E., and POMERANCE, C., *714 and 715*. J. Recreational Math. 7 (1974), 87–89.
- [14] NICOLAS, J.-L., *Sur l'ordre maximum d'un élément dans le groupe S_n des permutations*. Acta Arith. 14 (1968), 315–332.
- [15] RANKIN, R. A., *The difference between consecutive prime numbers*. J. London Math. Soc. 13 (1938), 242–247.

Math. Inst. Hung. Acad. Sci.,
Budapest, Hungary

Department of Mathematics,
University of Georgia,
Athens, Georgia 30602
U.S.A.