

Erdős-Szekeres-type theorems for monotone paths and convex bodies

Jacob Fox^{*} János Pach[†] Benny Sudakov[‡] Andrew Suk[§]

April 29, 2011

Dedicated to the 75th anniversary of the publication of the Happy Ending Theorem

Abstract

For any sequence of positive integers $j_1 < j_2 < \dots < j_n$, the k -tuples $(j_i, j_{i+1}, \dots, j_{i+k-1})$, $i = 1, 2, \dots, n-k+1$, are said to form a *monotone path* of length n . Given any integers $n \geq k \geq 2$ and $q \geq 2$, what is the smallest integer N with the property that no matter how we color all k -element subsets of $[N] = \{1, 2, \dots, N\}$ with q colors, we can always find a monochromatic monotone path of length n ? Denoting this minimum by $N_k(q, n)$, it follows from the seminal 1935 paper of Erdős and Szekeres that $N_2(q, n) = (n-1)^q + 1$ and $N_3(2, n) = \binom{2n-4}{n-2} + 1$. Determining the other values of these functions appears to be a difficult task. Here we show that

$$2^{(n/q)^{q-1}} \leq N_3(q, n) \leq 2^{n^{q-1} \log n},$$

for $q \geq 2$ and $n \geq q+2$. Using a “stepping-up” approach that goes back to Erdős and Hajnal, we prove analogous bounds on $N_k(q, n)$ for larger values of k , which are towers of height $k-1$ in n^{q-1} . As a geometric application, we prove the following extension of the Happy Ending Theorem. Every family of at least $M(n) = 2^{n^2 \log n}$ plane convex bodies in general position, any pair of which share at most two boundary points, has n members in convex position, that is, it has n members such that each of them contributes a point to the boundary of the convex hull of their union.

1 Introduction

The classic 1935 paper of Erdős and Szekeres [16] published in *Compositio Mathematica* was a starting point of a very rich discipline within combinatorics: Ramsey theory (see, e.g., [20]). Erdős liked to call the main result of this paper the “Happy Ending Theorem,” as its discovery was

^{*}Massachusetts Institute of Technology, Cambridge, MIT. Supported by a Simons Fellowship. Email: fox@math.mit.edu

[†]City College and Courant Institute, New York and EPFL, Lausanne. Supported by Grants from NSF, NSA, PSC-CUNY, BSF, OTKA, and SNF. Email: pach@cims.nyu.edu

[‡]Department of Mathematics, UCLA, Los Angeles, CA 90095. Email: bsudakov@math.ucla.edu. Research supported in part by NSF CAREER award DMS-0812005 and by a USA-Israeli BSF grant.

[§]Courant Institute, New York and EPFL, Lausanne. Email: suk@cims.nyu.edu

triggered by a geometric observation of Esther Klein, and the authors' collaboration with her eventually led to the marriage of Klein and Szekeres.

At the early stages of its development, Ramsey theory focused on the emergence of large monochromatic *complete* subgraphs in colored graphs and hypergraphs. In the 1960s and 70s, many initial results of Ramsey theory were revisited in a more general setting. For instance, Gerencsér and Gyárfás [19] proved that for any 2-coloring of the edges of a complete graph with roughly $3n/2$ vertices, there is a monochromatic path of length n . Recently, Figaj and Luczak [17] and, in a more precise form, Gyárfás et al. [21] have settled an old conjecture of Faudree and Schelp, according to which any 3-colored complete graph with roughly $2n$ vertices has a monochromatic path of length n . It follows from a theorem of Erdős and Gallai [15] that for every positive integer q there exists a constant $c(q) \leq q$ such that no matter how we color the edges of the complete graph $K_{c(q)n}$ with q colors, we can always find a monochromatic path of length n . The smallest value of the constant $c(q)$ is not known.

The problem becomes simpler if we consider *ordered* complete graphs and we want to find a long *increasing* path such that all of its edges are of the same color. It follows from the results of Erdős and Szekeres [16], and also from Dilworth's theorem on partially ordered sets [13], that for any 2-coloring of the edges of a complete graph with more than $(n-1)^2$ vertices, one can find a monochromatic monotone path with n vertices. An easy construction shows that this statement does not remain true for graphs with $(n-1)^2$ vertices. The result and its proof readily generalize to colorings with $q \geq 2$ colors: in this case, we have to replace $(n-1)^2$ with $(n-1)^q$.

The corresponding problems for hypergraphs are quite hard and their solution usually utilizes some variant or extension of the celebrated hypergraph regularity lemma. The only asymptotically sharp result of this kind was proved by Haxell, Łuczak et al. [22]. Let $H = (V(H), E(H))$ be a k -uniform hypergraph. A *tight path* or, shortly, a *path of length n* in H is comprised of a set of n distinct vertices $v_1, v_2, \dots, v_n \in V(H)$ and the set of $n-k+1$ hyperedges of the form

$$\{v_i, v_{i+1}, \dots, v_{i+k-1}\} \in E(H),$$

where $i = 1, 2, \dots, n-k+1$. It was shown in [22] that no matter how we 2-color all triples of a complete 3-uniform hypergraph on $4n/3$ vertices, one of the color classes will contain a path of length roughly n . This result is best possible. It was generalized by Nagle, Olsen et al. [26], Cooley, Fountoulakis et al. [12], and Conlon et al. [9]. These results imply that every q -coloring of the hyperedges of a complete k -uniform hypergraph on Cn vertices contains a monochromatic copy of any n -vertex hypergraph with maximum degree Δ , where $C = C(k, q, \Delta)$ is a suitable constant depending only on k , q , and Δ .

In the present paper, we consider the analogous problem for *ordered* hypergraphs. Let K_N^k denote the *complete k -uniform hypergraph*, consisting of all k -element subsets (k -tuples) of the vertex set $[N] = \{1, 2, \dots, N\}$. For any n positive integers, $j_1 < j_2 < \dots < j_n$, we say that the hyperedges

$$\{j_i, j_{i+1}, \dots, j_{i+k-1}\},$$

$i = 1, 2, \dots, n-k+1$, form a *monotone (tight) path of length n* . We will present several results on finding monochromatic monotone paths in edge-colored ordered hypergraphs. As a geometric application, in Section 7 we will show that every family of at least $2^{n^2 \log n}$ plane convex bodies in general position, any pair of which share at most two boundary points, has n members in convex

position. This substantially improves the previous double-exponential bound of Hubard et al. [23] for this problem. Here, and throughout the paper, all logarithms unless otherwise stated are in base 2.

1.1 Ramsey numbers for monotone paths

Let $N_k(q, n)$ denote the smallest integer N such that for every q -coloring of the hyperedges (k -tuples) of K_N^k , there exists a monotone path of length n such that all of its hyperedges are of the same color. Using this notation for graphs ($k = 2$), the corollary of the Erdős-Szekeres theorem or the Dilworth theorem mentioned above can be rephrased as

$$N_2(q, n) = (n - 1)^q + 1. \quad (1)$$

For 3-uniform hypergraphs, a straightforward generalization of the cup-cap argument of Erdős and Szekeres [16] gives

Theorem 1.1. *Let $N = N_3(2, n)$ be the smallest integer such that for every 2-coloring of all triples in $[N]$, there exists a monochromatic monotone path of length n . Then we have*

$$N_3(2, n) = \binom{2n - 4}{n - 2} + 1.$$

For a larger (but fixed) number of colors, we prove the following theorem which provides lower and upper bounds that are tight apart from a logarithmic factor in the exponent.

Theorem 1.2. *For any integers $q \geq 2$ and $n \geq q + 2$, we have*

$$2^{(n/q)^{q-1}} \leq N_3(q, n) \leq 2^{n^{q-1} \log n}.$$

Define the tower function $t_i(x, n)$ recursively as follows. Let $t_1(x, n) = x$ and $t_{i+1}(x, n) = n^{t_i(x, n)}$, so $t_i(x, n)$ is a tower of $i - 1$ n 's with an x on top. We let $t_i(x) = t_i(x, 2)$. Using a “stepping-up” approach, developed by Erdős and Hajnal (see [20]) and strengthened in [9], we show the following extension of Theorem 1.2 to k -uniform hypergraphs, for any $k \geq 3$.

Theorem 1.3. *For every $k \geq 3$ and q , there are positive constants c, c' depending only on k and q such that, for any $n \geq 4k$, we have*

$$t_{k-1}(cn^{q-1}) \leq N_k(q, n) \leq t_{k-1}(c'n^{q-1} \log n).$$

In Section 2, we prove the recursive upper bound $N_k(q, n) \leq N_{k-1}((n - k + 1)^{q-1}, n)$. This upper bound, together with equation (1), implies that $N_k(q, n) \leq t_k(q - 1, n)$ for $k \geq 3$. The upper bounds in Theorems 1.2 and 1.3 follow from this inequality.

In Sections 3 and 4, we establish the lower bound in Theorem 1.2. We also prove a general statement there (Theorem 4.3), providing a lower bound on $N_k(q, n)$, which is exponential in $N_{k-1}(q, n)$, for every $k \geq 4$. Putting these results together, the lower bound in Theorem 1.3 readily follows.

The case $k = 3$ is crucial to understanding the growth of $N_k(q, n)$. Indeed, the stepping-up lower bound mentioned above (see Theorem 4.3) together with the recursive upper bound in Theorem 1.4 and inequality 2, described in the next subsection, show that closing the gap between the upper and lower bounds on $N_k(q, n)$ in the special case $k = 3$ would also close the gap for all larger values of k .

1.2 Online and size Ramsey numbers

Consider the following game played by two players, *builder* and *painter*. For $t \geq 1$, at the beginning of stage t , a new vertex v_t is added (the vertices v_i for $1 \leq i < t$ already are present), and for each $(k-1)$ -tuple of vertices $(v_{i_1}, \dots, v_{i_{k-1}})$ with $1 \leq i_1 < \dots < i_{k-1} < t$, builder decides (in any order) whether to draw the edge $(v_{i_1}, \dots, v_{i_{k-1}}, v_t)$. If builder draws the edge, then painter has to immediately color it in one of q colors $1, \dots, q$. The *(vertex) online Ramsey number* $V_k(q, n)$ is the minimum number of edges builder has to draw to guarantee a monochromatic monotone path of length n .

An ordered k -uniform hypergraph G is said to be (q, n) -*path Ramsey* if for every q -edge-coloring of G one can find a monochromatic monotone path of length n . The *size Ramsey number* $S_k(q, n)$ is the minimum number of edges of an ordered k -uniform hypergraph G which is (q, n) -path Ramsey. It follows from the definitions that

$$N_k(q, n - k + 1) \leq V_k(q, n) \leq S_k(q, n) \leq \binom{N_k(q, n)}{k}. \quad (2)$$

Indeed, the first inequality comes from the following painter strategy. Let w_i be the i th vertex that builder adds in which there is an edge whose largest vertex is w_i (there may be vertices not given labels). That is, $w_i = v_{j(i)}$ where $j(i)$ is the i th stage for which at least one edge is added. Painter colors each edge $(w_{i_1}, w_{i_2}, \dots, w_{i_k})$ with $i_1 < \dots < i_k < N_k(q, n - k + 1)$ the same color as the color of (i_1, \dots, i_k) in a q -edge-coloring of the ordered complete k -uniform hypergraph on $N_k(q, n - k + 1) - 1$ vertices with no monochromatic path of length $n - k + 1$. All other edges, i.e., those containing a vertex which is not the largest vertex in some edge, can be colored arbitrarily. Note that every monotone path of length n contains a path of length $n - k + 1$ such that each vertex is the largest vertex in an edge of the ordered hypergraph. Therefore, this coloring has no monochromatic monotone path of length n , and painter guarantees that there are at least $N_k(q, n - k + 1)$ edges when the first monochromatic monotone path of length n appears. On the other hand, if builder selects all edges of a (q, n) -path Ramsey ordered k -uniform hypergraph, she will definitely win; which proves the second inequality. The third inequality is trivial, taking into account that the ordered complete k -uniform hypergraph K_N^k with $N = N_k(q, n)$ is (q, n) -path Ramsey.

Conlon, Fox, and Sudakov [10] used online Ramsey numbers to give an upper bound for the Ramsey number of a *complete* k -uniform hypergraph with n vertices. Here we apply this technique to establish an upper bound for the Ramsey number of a *monotone path* of length n . More precisely, we show

Theorem 1.4. *For every $k \geq 3$ and for every $q \geq 1$, $n \geq k$, we have*

$$N_k(q, n) \leq q^{V_{k-1}(q, n+k-2)} + k - 2.$$

In the special case $k = 3$, this bound is worse than the upper bound in Theorem 1.2. However, Theorem 1.4 is already useful for $k \geq 4$. Indeed, any improvement on the upper bound in Theorem 1.3 for $k = 3$, together with the trivial upper bound on $V_{k-1}(q, n + k - 2)$ in (2), will lead to analogous improvements for all larger k . For example, if, for fixed q , we would know that $N_3(q, n) = 2^{O(n^{q-1})}$, then using Theorem 1.4 we would also know that $N_4(q, n) = 2^{2^{O(n^{q-1})}}$. Note that the recursive bound in Theorem 2.1 does not give such an improvement even if we knew that $N_3(q, n)$

is exponential in n^{q-1} . Similarly, the stepping-up lower bound in Theorem 4.3 shows that any improvement on the lower bound in Theorem 1.3 for $k = 3$ will lead to analogous improvements for all larger k .

In addition to being useful for bounding classical Ramsey numbers as discussed above, online and size Ramsey numbers are now well-studied topics in their own right. The *size Ramsey number* $r_e(H)$ of a graph H is the minimum number of edges of a graph G in which every 2-edge-coloring of G contains a monochromatic copy of H . The study of these numbers was initiated by Erdős et al. [14]. A fundamental problem of Erdős in this area is to determine the growth of the size Ramsey number of paths. Beck [2] solved this problem, proving that these numbers grow linearly. That is, $r_e(P_n) \leq cn$, where c is an absolute constant. In contrast, Rödl and Szemerédi [31] disproved a conjecture of Beck by showing that there are graphs of maximum degree 3 whose size Ramsey number grows superlinear in the number of vertices. In the other direction, Kohayakawa et al. [24] recently showed that graphs on n vertices with fixed maximum degree Δ have size Ramsey number at most $O(n^{2-1/\Delta} \log^{1/\Delta} n)$.

Another online Ramsey game which is quite close to ours was introduced independently by Beck [3] and Kurek and Ruciński [25]. In this game, there are two players, Builder and Painter, who move on the originally empty graph with an unbounded number of vertices. At each step, Builder draws a new edge and Painter has to color it either red or blue immediately. The *edge online Ramsey number* $\bar{r}(H)$ is the minimum number of edges that builder has to draw in order to force painter to create a monochromatic H . As Builder can simply draw the edges of a given graph, we have the inequality $\bar{r}(H) \leq r_e(H)$. A basic conjecture in this area due to Rödl is to show that $\lim_{n \rightarrow \infty} \frac{\bar{r}(K_n)}{r_e(K_n)} = 0$. Conlon [8] made substantial progress on this conjecture, showing that $\bar{r}(K_n) \leq c^n r_e(K_n)$ holds infinitely often, where $c < 1$ is an absolute constant. Randomized variants of the edge online Ramsey number were studied in [1], [4], [18].

Note that the problem of estimating $V_k(q, n)$ and $S_k(q, n)$ is most interesting in the case of graphs ($k = 2$). Indeed, for larger k , $N_k(q, n)$ grows roughly $(k-2)$ -fold exponential in n^{q-1} , so the lower and upper bounds on these numbers are roughly determined by $N_k(q, n)$. We will therefore focus our attention on the interesting case $k = 2$.

In view of (2), one may think that the functions $V_k(q, n)$ and $S_k(q, n)$ cannot differ too much. Rather surprisingly, already for graphs ($k = 2$) this is not the case. In Section 5, we show that the size Ramsey number satisfies $S_2(q, n) \geq c_q n^{2q-1}$. On the other hand, also in that section we prove

Theorem 1.5. *We have*

$$V_2(2, n) = (1 + o(1))n^2 \log_2 n,$$

and for every fixed $q \geq 2$, there are constants c_q and c'_q such that

$$c_q n^q \log n \leq V_2(q, n) \leq c'_q n^q \log n.$$

Moreover, the proof of Theorem 1.5 shows that builder has a winning strategy which uses $N_2(q, n)$ vertices and each vertex v_t belongs to at most $c'_q \log n$ edges (v_j, v_t) with $j < t$. This theorem shows that, by using the information on the initial portion of the coloring, builder can substantially reduce the number of edges that guarantee a monochromatic monotone path.

1.3 Noncrossing convex bodies in convex position

In the last section, we apply Theorem 1.2 and Lemma 6.1 to a problem for families of convex bodies. To formulate our question, we need some definitions. A family \mathcal{C} of convex bodies (compact convex

sets with nonempty interior) in the plane is said to be in *convex position* if none of its members is contained in the convex hull of the union of the others. We say that \mathcal{C} is in *general position* if

1. every three members of \mathcal{C} are in convex position;
2. no two members $C, C' \in \mathcal{C}$ have a common tangent that meets $C \cap C'$; and
3. no three members of \mathcal{C} share a common tangent.

Bisztriczky and Fejes-Tóth [5] generalized the Happy Ending theorem of Erdős and Szekeres [16], [6] as follows. They proved that for every n , there exists a function $D(n)$ such that any family of $D(n)$ pairwise disjoint convex bodies in general position in the plane contains n members in convex position. It was proved in [27] that $D(n) \leq \binom{2n-4}{n-2} + 1$.

In [28], the Bisztriczky-Fejes Tóth theorem has been extended to families of *noncrossing* convex bodies, that is, to convex bodies, no pair of which share more than *two* boundary points. It was proved that there exists a function $N = M(n)$ such that from every family of N noncrossing convex bodies in general position in the plane one can select n members in convex position. More recently, Hubard et al. [23] showed that the function $M(n)$ grows at most double exponentially in n . We use Theorem 1.2 to obtain a much better bound.

Theorem 1.6. *Let $N = M(n)$ denote the smallest integer such that every family of N noncrossing convex bodies in general position in the plane has n members in convex position. Then we have*

$$M(n) \leq N_3(3, n) \leq 2^{n^2 \log n}.$$

Note that no analogue of the last theorem is true if we drop the assumption that the bodies are noncrossing. One can construct a family of arbitrarily many pairwise crossing rectangles that which are in general position, but no four of them are in convex position (see [28]).

Theorem 1.6 can be established, as follows. First, we choose a coordinate system, in which the *left endpoint* (i.e., the leftmost point) of every member of the family is unique, and the x -coordinates of the left endpoints are different. Let C_1, C_2, \dots, C_N denote the members of our family, listed in the increasing order of the x -coordinates of their left endpoints. Suppose that $N = N_3(3, n)$.

In the last section, we define a 3-coloring of all ordered triples (C_i, C_j, C_k) with $i < j < k$. We will show that this coloring has the property that if a sequence of sets C_{i_1}, C_{i_2}, \dots ($i_1 < i_2 < \dots$) induces a monochromatic monotone path, then they are in convex position.

According to Theorem 1.2, $N \leq n^{n^2}$. Therefore, our colored triple system contains a monochromatic monotone path of length n . In other words, there exists a sequence $i_1 < i_2 < \dots < i_n$ such that all ordered triples induced by the sets C_{i_j} are of the same color. It follows from the special property of our coloring mentioned above that the sets C_{i_1}, \dots, C_{i_n} are in convex position. This will complete the proof of Theorem 1.6.

Organization: The rest of the paper is organized as follows. In the next section we prove a recursive upper bound on $N_k(q, n)$. In Section 3, we prove results on monochromatic walks in digraphs and the minimum chromatic number of a (q, n) -path Ramsey graph which will be used in the subsequent two sections. In Section 4, we prove lower bounds on $N_k(q, n)$. In Section 5, we prove bounds on size and online Ramsey numbers of paths, as well as an alternative upper bound on $N_k(q, n)$ in terms of online Ramsey numbers. In Section 6, we prove a lemma relating monochromatic cliques in “transitive” colorings to monochromatic monotone paths in general colorings. We

use this lemma in Section 7 in the proof of Theorem 1.6. We finish with some concluding remarks and open problems in Section 8.

2 An upper bound on $N_k(q, n)$

Given any two positive integers $i \leq j$, let $[i, j]$ denote the set of positive integer h with $i \leq h \leq j$, and $[i] = [1, i]$. We begin by proving a recursive upper bound on $N_k(q, n)$.

Theorem 2.1. *For any integers $n \geq k \geq 2$ and $q \geq 2$, we have*

$$N_k(q, n) \leq N_{k-1}((n - k + 1)^{q-1}, n).$$

Proof. Suppose for contradiction that there is a q -coloring

$$\chi : \binom{[N]}{k} \rightarrow [q]$$

of the edges of the complete k -uniform ordered hypergraph K_N^k on $N = N_{k-1}((n - k + 1)^{q-1}, n)$ vertices with no monochromatic path of length n . Define the auxiliary coloring

$$\phi : \binom{[N]}{k-1} \rightarrow [k-1, n-1]^{q-1},$$

as follows. For any $v_1 < \dots < v_{k-1}$, we let $\phi(v_1, \dots, v_{k-1}) = (n_1, \dots, n_{q-1})$, where n_i is the length of the longest monochromatic path in color i ending with v_1, \dots, v_{k-1} . By convention, any $k-1$ vertices v_1, \dots, v_{k-1} form a monochromatic monotone path of length $k-1$ with respect to coloring χ . Thus, we have $k-1 \leq n_i \leq n-1$, for every i ($1 \leq i \leq q-1$).

By our assumption, $N = N_{k-1}((n-k+1)^{q-1}, n)$, so that in coloring ϕ , there is a monochromatic monotone path of length n . Let $u_1 < \dots < u_n$ denote the vertices of this path. To complete the proof of the theorem, it is sufficient to show that these vertices form a monochromatic monotone path in color q , with respect to the coloring χ . Suppose that this is not the case. Then, for some $1 \leq j \leq n - k + 1$ and $1 \leq i \leq q - 1$, we have $\chi(u_j, u_{j+1}, \dots, u_{j+k-1}) = i$. Since the vertices u_1, \dots, u_n form a monochromatic monotone path with respect to the coloring ϕ , it follows that in coloring χ , the length of the longest monochromatic monotone path of color i ending with $u_j, u_{j+1}, \dots, u_{j+k-2}$ is the same as the length of the longest monochromatic monotone path of color i ending with $u_{j+1}, u_{j+2}, \dots, u_{j+k-1}$. However, any longest monochromatic monotone path of color i ending with $u_j, u_{j+1}, \dots, u_{j+k-2}$ can be extended to a longer monochromatic monotone path of color i , by adding the vertex u_{j+k-1} . This contradiction completes the proof of the theorem. \square

Now we are in a position to prove the upper bound in Theorem 1.2.

Proof of the upper bounds in Theorems 1.2 and 1.3. We prove the stronger inequality

$$N_k(q, n) \leq t_k(q-1, n),$$

for every $k \geq 3$. The proof is by induction on k . By (1), we have $N_2(q, n) = (n-1)^q + 1$. According to Theorem 2.1,

$$N_3(q, n) \leq N_2((n-2)^{q-1}, n) = (n-1)^{(n-2)^{q-1}} + 1 < n^{n^{q-1}} = t_3(q-1, n).$$

Now suppose that the desired inequality holds for k . Then it also holds for $k + 1$, because, again by Theorem 2.1, we have

$$N_{k+1}(q, n) \leq N_k((n - k)^{q-1}, n) \leq t_k((n - k)^{q-1}, n) \leq t_k(n^{q-1}, n) = t_{k+1}(q - 1, n).$$

□

3 Monochromatic walks in digraphs

Let D_N denote the *complete digraph* on N vertices, that is, the directed graph in which each pair of distinct vertices is connected by two edges with opposite orientations. A *walk of length n* in a digraph is a sequence of n vertices v_1, \dots, v_n with possible repetitions such that for every i ($1 \leq i < n$), the directed edge $\overrightarrow{v_i v_{i+1}}$ belongs to the digraph. If $v_1 = v_n$, then the walk is called *closed*. Note that closed walks can be used to construct walks which are arbitrarily long. A digraph with no closed walk is *acyclic*.

Let $f(q, n)$ be the smallest number N such that for every q -coloring of the edges of D_N , there is a monochromatic walk of length n . First, we show that for a fixed q , the order of magnitude of $f(q, n)$ is n^{q-1} . In the next section, this fact is used to establish the lower bound on $N_3(q, n)$, stated in Theorem 1.2.

Theorem 3.1. *For any integers $n, q \geq 2$, we have*

$$(n/q)^{q-1} \leq f(q, n) \leq N_2(q - 1, n) = (n - 1)^{q-1} + 1.$$

Proof. We first establish the upper bound. Consider a q -coloring of the edges of D_N with $N = N_2(q - 1, n)$. We have to show that there is a monochromatic walk with n vertices. Suppose that the set of edges of color q does not determine a walk of length n . Then these edges form an acyclic digraph. Hence, there is an ordering of the vertices of D_N so that the edges of color q go backwards. Thus, removing the backwards edges, we have an ordered complete graph on $N = N_2(q - 1, n)$ vertices, whose edges are colored with $q - 1$ colors. In one of the color classes, we can find a monotone path with n vertices, which is a monochromatic directed path in D_N .

The lower bound follows from Lemma 3.2 below. Indeed, letting $m = 1 + \lfloor \frac{n-2}{q-1} \rfloor$, it is easy to check that $2 + (q - 1)(m - 1) \leq n$ and $m \geq n/q$, so

$$f(q, n) \geq f(q, 2 + (q - 1)(m - 1)) \geq m^{q-1} \geq (n/q)^{q-1}.$$

□

Lemma 3.2. *For any integers $n, q \geq 2$, we have $f(q, 2 + (q - 1)(n - 1)) > n^{q-1}$.*

Proof. Define the coloring ϕ of the edges of the complete digraph on $[n]^{q-1}$, as follows. For any pair of distinct vertices, $a = (a_1, \dots, a_{q-1}), b = (b_1, \dots, b_{q-1})$, define the color of the directed edge (a, b) as the smallest coordinate i for which $a_i < b_i$. If there is no such coordinate, that is, if $a_i \geq b_i$ for all i ($1 \leq i \leq q - 1$), then color the directed edge (a, b) with color q . For each color $i \leq q - 1$, the length of a longest monochromatic walk of color i is n , because traversing any edge of color i , the i th coordinate must increase. On the other hand, along any walk in color class q , the coordinates

are never allowed to increase, and in each step at least one of them must strictly decrease, so that the sum of the coordinates strictly decreases. The sum of the coordinates of a point in $[n]^{q-1}$ is at least $q-1$ and is at most $(q-1)n$. Therefore, the length of such a monochromatic walk does not exceed $(q-1)n - (q-1) + 1 = 1 + (q-1)(n-1)$. \square

Recall from Subsection 1.2 that an ordered graph G is called (q, n) -path Ramsey if for every q -edge-coloring of G there exists a monochromatic monotone path of length n . Let $\chi(q, n)$ denote the minimum chromatic number of an ordered graph G which is (q, n) -path Ramsey.

We close this section by showing that the functions $\chi(q, n)$ and $f(q, n)$ are actually identical. Therefore, Theorem 3.1 implies that, for a fixed q , $\chi(q, n)$ also grows on the order of n^{q-1} .

The (classical) Ramsey number $R(n; q)$ is the minimum N such that every q -edge-coloring of the complete graph on N vertices contains a monochromatic clique on n vertices.

Theorem 3.3. *For any integers $q, n \geq 2$, we have*

$$\chi(q, n) = f(q, n).$$

Proof. We first show that $\chi(q, n) \geq f(q, n)$. Consider a q -edge-coloring ϕ of the edges of the complete digraph with vertex set $[f(q, n) - 1]$ without a monochromatic walk of length n . Assume for contradiction that there exists an ordered graph G with $\chi(G) < f(q, n)$, which is (q, n) -path Ramsey, and consider a partition $V(G) = V_1 \cup \dots \cup V_t$ into independent sets with $t = \chi(G)$. Define a q -coloring ρ of the edges of G , as follows. If v, w are adjacent with $v < w$ and $v \in V_i$ and $w \in V_j$, let $\rho(v, w) = \phi(i, j)$. If the vertices $v_1 < \dots < v_n$ form a monochromatic monotone path with respect to the edge-coloring ρ of G , then denoting by i_k the integer for which $v_k \in V_{i_k}$, we have that i_1, \dots, i_n form a monochromatic walk with respect to the edge-coloring ϕ . This is a contradiction, showing that our assumption $\chi(G) < f(q, n)$ was wrong.

To show that $\chi(q, n) \leq f(q, n)$, we define an ordered graph H , which is $f(q, n)$ -colorable and (q, n) -path Ramsey. Let $V(H) = [Rt]$, where $t = f(q, n)$ and $R = R(n; Q)$ is the Q -color Ramsey number for the complete graph on n vertices with $Q = q^{t^2-t}$. Connect two vertices i, j ($1 \leq i < j \leq Rt$) by an edge in H if $j - i$ is not a multiple of t . The partition of $V(H)$ modulo t defines a proper t -coloring of H . Since the first t vertices of H form a clique, we have $\chi(H) = t$. It remains to prove that H is (q, n) -path Ramsey. Consider an edge-coloring ψ of H with q colors. Define an auxiliary Q -coloring τ of the complete graph on $[R]$ vertices, where the color of the edge between u, v ($1 \leq u < v \leq R$) is given as a $t \times t$ matrix $A = (a_{ij})$ with 0's in the diagonal and $a_{ij} = \psi((u-1)t + i, (v-1)t + j)$ for every $i \neq j$. Obviously, the number of colors used in this coloring is at most the number of possible matrices A , which is equal to $q^{t^2-t} = Q$. By the definition of the Ramsey number $R = R(n; Q)$, $[R]$ contains a monochromatic clique on n vertices, with respect to the coloring τ . Denote the vertices of such a clique by $u_1 < u_2 < \dots < u_n$. Define a q -edge-coloring ξ of the complete digraph on $[t]$, where $\xi(i, j) = \psi((u_1-1)t + i, (u_2-1)t + j)$. Since $t = f(q, n)$, there is a monochromatic walk i_1, \dots, i_n of length n with respect to the coloring ξ . Then the vertices $(u_1-1)t + i_1, (u_2-1)t + i_2, \dots, (u_n-1)t + i_n$ form a monochromatic monotone path in the q -edge-coloring ψ of H , which completes the proof. \square

4 Lower bounds on $N_k(q, n)$

In this section we adapt the stepping-up approach of Erdős and Hajnal on hypergraph Ramsey numbers to provide lower bounds for $N_k(q, n)$, stated in Theorems 1.2 and 1.3. First we address

the case $k = 3$.

Theorem 4.1. *For any integers $q, n \geq 2$, we have*

$$N_3(q, n) > 2^{f(q, n-1)-1}.$$

Proof. Let ϕ be a q -coloring of the edges of the complete digraph D on vertex set $[0, f(q, n-1) - 2]$ without a monochromatic walk on $n-1$ vertices. We use ϕ to define a q -coloring χ of the hyperedges (triples) of the complete ordered 3-uniform hypergraph K_N^3 on the vertex set $V = [N]$ with $N = 2^{f(q, n-1)-1}$, as follows.

For any $a \in V$, write $a - 1 = \sum_{i=0}^{f(n-1, q)-2} a(i)2^i$ with $a(i) \in \{0, 1\}$ for each i . For $a \neq b$, let $\delta(a, b)$ denote the largest i for which $a(i) \neq b(i)$. Obviously, we have $\delta(a, b) \neq \delta(b, c)$ for every triple $a < b < c$.

Given any triple $a < b < c$, define $\chi(a, b, c) = \phi(\delta(a, b), \delta(b, c))$. To complete the proof of the theorem, it is enough to show that, with respect to this coloring, there is no monochromatic monotone path of length n . Suppose for contradiction that there is such a path, and denote its vertices by $a_1 < a_2 < \dots < a_n$. Letting $\delta_j = \delta(a_j, a_{j+1})$ for every j ($1 \leq j < n$), it follows from the definition of the coloring χ that the integers $\delta_1, \dots, \delta_{n-1} \in [0, f(q, n-1) - 2]$ induce a monochromatic walk in the digraph D , with respect to the coloring ϕ . This contradiction completes the proof. \square

From Lemma 3.2, we have the estimate $f(q, n-1) - 1 \geq m^{q-1}$ with $m = 1 + \lfloor \frac{n-3}{q-1} \rfloor$. It is easy to check that $m \geq n/q$ if $n \geq q + 2$. Together with Theorem 4.1, we immediately obtain the following corollary, which is the same as the lower bound in Theorem 1.2.

Corollary 4.2. *For any integers $q \geq 2$ and $n \geq q + 2$, we have*

$$N_3(q, n) \geq 2^{(n/q)^{q-1}}.$$

We next give a recursive lower bound on $N_k(q, n)$ for $k \geq 4$. The proof is an adaptation of an improved version of the stepping-up technique, due to Conlon, Fox, and Sudakov [11]. It is easy to see that, together with Corollary 4.2, this gives the lower bound in Theorem 1.3.

Theorem 4.3. *For any integers $n \geq k \geq 4$ and $q \geq 2$, we have*

$$N_k(q, n+3) > 2^{N_{k-1}(q, n)-1}.$$

Proof. We start the proof in a way similar to Theorem 4.1.

Let ϕ be a q -coloring with colors $1, \dots, q$ of the edges of the complete ordered $(k-1)$ -uniform hypergraph on $N_{k-1}(q, n) - 1$ vertices without a monochromatic monotone path on n vertices. We use ϕ to define a q -coloring χ of the edges of the complete ordered k -uniform hypergraph K_N^k on the vertex set $V = [N]$ with $N = 2^{N_{k-1}(q, n)-1}$, as follows.

For any $a \in V$, write $a - 1 = \sum_{i=0}^{N_{k-1}(q, n)-2} a(i)2^i$ with $a(i) \in \{0, 1\}$ for each i . For $a \neq b$, let $\delta(a, b)$ denote the largest i for which $a(i) \neq b(i)$. As in the proof of Theorem 4.1, we have $\delta(a, b) \neq \delta(b, c)$ for every triple $a < b < c$.

Given any k -tuple $a_1 < a_2 < \dots < a_k$ of V , consider the integers $\delta_i = \delta(a_i, a_{i+1})$, $1 \leq i \leq k-1$. If $\delta_1, \dots, \delta_{k-1}$ form a monotone sequence, then let $\chi(a_1, a_2, \dots, a_k) = \phi(\delta_1, \delta_2, \dots, \delta_{k-1})$.

Now we have to color the k -tuple (a_1, \dots, a_k) in the case when $\delta_1, \dots, \delta_{k-1}$ is not monotone. We say that i is a *local minimum* if $\delta_{i-1} > \delta_i < \delta_{i+1}$, a *local maximum* if $\delta_{i-1} < \delta_i > \delta_{i+1}$, and a *local extremum* if it is either a local minimum or a local maximum. Since $\delta_{i-1} \neq \delta_i$ for every i , every nonmonotone sequence $\delta_1, \dots, \delta_{k-1}$ has a local extremum. Let i_1 denote the first local extremum. If $\delta_1, \dots, \delta_{k-1}$ is not monotone, we define $\chi(a_1, \dots, a_k) = 1$ if i_1 is even and a local maximum, or if i_1 is odd and a local minimum. Otherwise, let $\chi(a_1, \dots, a_k) = 2$.

Suppose for contradiction that the vertices $a_1 < \dots < a_{n+3}$ induce a monochromatic monotone path with respect to the q -coloring χ . For $1 \leq i \leq n+2$, let $\delta_i = \delta(a_i, a_{i+1})$. Since there is no monochromatic monotone path on n vertices with respect to the coloring ϕ , the sequence $\delta_1, \dots, \delta_{n+2} \in [0, N_{k-1}(q, n) - 2]$ must have a local extremum. Moreover, if i_1 denotes the location of the first local extremum, we have $2 \leq i_1 \leq n-1$.

CASE 1: The first local extremum i_1 satisfies $i_1 > 2$. We claim that, if $i_1 < k-1$, then the first two edges of the path, $e_1 = (a_1, \dots, a_k)$ and $e_2 = (a_2, \dots, a_{k+1})$, receive different colors, contradicting our assumption that the path is monochromatic. If $i_1 \geq k-1$, then the edges $e_1 = (a_{i_1-k+3}, \dots, a_{i_1+2})$ and $e_2 = (a_{i_1-k+4}, \dots, a_{i_1+3})$ receive different colors, which is again a contradiction. Indeed, in either case the *type* (maximum or minimum) of the first local extremum is the same for these two edges, but their locations differ by one and hence have different parity, which implies that e_1 and e_2 receive different colors.

CASE 2: The first local extremum is $i_1 = 2$.

CASE 2(A): 3 is a local extremum. Note that consecutive extrema have different types, so that the types of 2 and 3 are different. Therefore, the first two edges of the path, $e_1 = (a_1, \dots, a_k)$ and $e_2 = (a_2, \dots, a_{k+1})$, must have different colors, contradicting our assumption that the path is monochromatic. Indeed, for each of these edges, the first local extremum is the second δ , but these extrema are of different types, and hence e_1 and e_2 receive different colors.

CASE 2(B): 3 is not a local extremum. As the sequence $\delta_2, \dots, \delta_{n+1}$ cannot be monotone, the sequence of δ 's has a second local extremum i_2 satisfying $3 < i_2 \leq n$. If $i_2 < k$, then the edges $e_1 = (a_2, \dots, a_{k+1})$ and $e_2 = (a_3, \dots, a_{k+2})$ have different colors. If $i_2 \geq k$, then the edges $e_1 = (a_{i_2-k+3}, \dots, a_{i_2+2})$ and $e_2 = (a_{i_2-k+4}, \dots, a_{i_2+3})$ have different colors. Indeed, in either case, the first local extremum for the pair of edges are of the same type, but their locations differ by one and hence have different parity, which implies they have different colors. This completes the proof of the theorem. \square

5 Size and Vertex Online Ramsey numbers for monotone paths

We begin this section by estimating vertex online Ramsey numbers, introduced in Subsection 1.2 of the Introduction. We then prove an alternative upper bound on $N_k(q, n)$, in terms of the online Ramsey number $V_{k-1}(q, n)$. We end this section by estimating the size Ramsey number $S_2(q, n)$.

5.1 Games

In this subsection we study vertex online Ramsey numbers. We first relate the online Ramsey game to another game, which we call the (q, n) -*lattice game* (or *lattice game* for short). In this game there are two players, builder and coordinator. During the game, a sequence of (not necessarily

distinct) points in the q -dimensional grid $\mathbb{Z}_{>0}^q$ with positive coordinates is built. After stage i , the sequence has length i . In stage $i + 1$, a new point p_{i+1} is added to the sequence as described below.

In each step of stage $i + 1$, builder picks a point $p_j = (x_{1,j}, \dots, x_{q,j})$ with $j \leq i$ already in the sequence. Coordinator then decides a coordinate $k \in [q]$. The point p_{i+1} must satisfy that its k th coordinate is greater than the k th coordinate of p_j , i.e., $x_{k,j} < x_{k,i+1}$. After some steps, builder decides it is time to end this stage, and coordinator picks a point p_{i+1} satisfying the conditions provided by the steps. Note that in stage 1, as there are no points yet in the sequence, no steps are taken and the only thing that happens is coordinator picks a point p_1 in $\mathbb{Z}_{>0}^q$ to begin the sequence. Let $L(q, n)$ be the minimum (total) number of steps needed for builder to guarantee that there is a point in the sequence with a coordinate at least n . The following lemma relates these numbers with online Ramsey numbers.

Lemma 5.1. *For all q and n , we have*

$$L(q, n) = V_2(q, n).$$

Proof. We first prove the bound $L(q, n) \geq V_2(q, n)$. As builder and painter are playing the Ramsey game, builder and coordinator play a corresponding lattice game. We will show that as long as painter guarantees in the Ramsey game there is no monochromatic monotone path of length n , coordinator can guarantee in the lattice game that no point in the sequence will have a coordinate at least n . Each vertex v_i in the Ramsey game corresponds to a point p_i in the lattice game. To prove the desired result, coordinator will guarantee that the k th coordinate of the point p_i in the lattice game is the length of the longest monotone monochromatic path in color k ending at v_i in the Ramsey game. At each step of stage i in the lattice game, builder picks the point p_j . Then, in the Ramsey game builder picks the edge (v_j, v_i) to add. The color k , $1 \leq k \leq q$, that painter picks to color the edge (v_j, v_i) is the coordinate which coordinator picks so that the k th coordinate of p_i is greater than the k th coordinate of p_j . At the end of this stage, coordinator picks the point p_i such that the k th coordinate of p_i is the length of the longest monotone monochromatic path in color k in the Ramsey game ending at v_i . This shows that, no matter what builder's strategy is, coordinator can mimic painter's strategy and continue the lattice game for as long as the Ramsey game.

Next we show that $L(q, n) \leq V_2(q, n)$, which would complete the proof. As builder and coordinator are playing the lattice game, builder and painter play a corresponding Ramsey game. We will show that as long as coordinator guarantees in the lattice game that no point in the sequence will have a coordinate at least n , painter can guarantee in the Ramsey game there is no monochromatic monotone path of length n . Each point p_i in the lattice game will have a corresponding vertex v_i in the Ramsey game. To prove the desired result, painter will guarantee that for each vertex v_i and each color k , the length of the longest monochromatic path in color k ending at v_i is at most the k th coordinate of p_i .

At each step of stage i in the Ramsey game, the builder picks an edge (v_j, v_i) to add. Then, in the corresponding lattice game, builder picks the point p_j to compare p_i with. The coordinate k that coordinator picks so that the k th coordinate of p_i is greater than the k th coordinate of p_j will be the color k that painter uses to color the edge (v_j, v_i) . The length of the longest monochromatic monotone path in color k with last edge (v_j, v_i) is one more than the length of the longest monochromatic monotone path in color k ending at v_j , and this length is at most the k th coordinate of p_j by construction. Since the k th coordinate of p_i is greater than the k th coordinate of

p_j , this guarantees that the length of the longest monochromatic monotone path in color k ending at v_i is at most the k th coordinate of p_i . This shows that, no matter what builder's strategy is, painter can mimic coordinator's strategy and continue the Ramsey game for as long as the lattice game. \square

The next lemma tells us the minimum number of steps required for builder to win the (q, n) -lattice game. The lattice \mathbb{Z}^q naturally comes with a partial order \prec , where $p \prec q$ if $p \neq q$ and the k th coordinate of q is at least the k th coordinate of p for $1 \leq k \leq q$. We will use this order further on.

Lemma 5.2. *In the (q, n) -lattice game, the minimum number of stages (i.e., the number of points in the constructed sequence) builder needs to guarantee a point in the sequence with a coordinate at least n is $(n - 1)^q + 1$.*

Proof. Indeed, in each stage, builder, by picking all previous points at steps, can ensure that the points in the constructed sequence are distinct. Hence, as there are only $(n - 1)^q$ points in $\mathbb{Z}_{>0}^q$ with coordinates at most $n - 1$, by the pigeonhole principle, the minimum number of stages required for builder is at most $(n - 1)^q + 1$. Picking a linear extension of the partial order \prec on $[n - 1]^q$ described above, coordinator can guarantee that p_i is the i th smallest point in the linear extension. In this way, the minimum number of stages required for builder is at least $(n - 1)^q + 1$, which completes the proof. \square

For a finite subset $S \subset \mathbb{Z}^q$ and a point $p \in \mathbb{Z}^q$, define the *position of p with respect to S* to be the maximum t such that for each $k \in [q]$, there are at least t points $s \in S$ such that the k th coordinate of p is at least the k th coordinate of s .

Lemma 5.3. *Let S be a finite nonempty subset of \mathbb{Z}^q .*

1. *There is a point $p \in S$ whose position with respect to S is at least $|S|/q$.*
2. *If $p \in \mathbb{Z}^q$ is such that there is no $s \in S$ satisfying $s \prec p$, then the position of p with respect to S is at most $(1 - \frac{1}{q})|S|$.*

Proof. We first show the first part. For each t , the number of elements of position less than t is less than tq . Indeed, for each of the q coordinates, less than t elements of S have position less than t in that coordinate. All other elements have position at least t . Setting $t = |S|/q$, there is a point of S of position at least $|S|/q$.

We now show the second part. Suppose the position of p with respect to S is t . Delete from S all elements s for which there is $k \in [q]$ such that the k th coordinate of s is larger than the k th coordinate of p . Since p has position t , the number of deleted elements is at most $q(|S| - t)$. If $|S| > q(|S| - t)$, then p is at least as large as an element of S , a contradiction. Hence, $|S| \leq q(|S| - t)$, and we have $t \leq (1 - \frac{1}{q})|S|$, which completes the proof. \square

We now prove an upper bound on $L(q, n)$.

Lemma 5.4. *Let $a(q, n) = 1 + (q - 1) \frac{\log n}{\log(\frac{n}{q-1})}$ and $b(q, n) = (n - 1)^q + 1$. In the (q, n) -lattice game, builder has a winning strategy which uses the minimum possible number of stages, which is $b(q, n)$, such that each stage uses at most $a(q, n)$ steps. In particular, we have*

$$L(q, n) \leq b(q, n)a(q, n).$$

Proof. By Lemma 5.2, the minimum possible number of stages builder needs to win the (q, n) -lattice game is indeed $b(q, n)$.

As there are at most $(n - 1)^q$ points in \mathbb{Z}^q with positive coordinates at most $n - 1$, it suffices for builder to guarantee at each stage i , using at most $a(q, n)$ steps, that coordinator picks a point p_i not already in the sequence. To accomplish this, it suffices for builder to guarantee that for each point p_j in the sequence with $j < i$, there is $k \in [q]$, such that the k th coordinate of p_i must be greater than the k th coordinate of p_j .

Suppose we have finished $i - 1$ stages, and we are now on stage i . Let M_0 denote the set of maximal elements in the already constructed sequence of $i - 1$ points. We have $|M_0| \leq (n - 1)^{q-1}$ as M_0 cannot contain two points with the same $q - 1$ first coordinates.

In the first step of stage i , builder picks a point $p_{j_1} \in M_0$ of position in M_0 at least $|M_0|/q$. Such a point exists by the first part of Lemma 5.3. Coordinator then picks a coordinate k_1 . By the definition of position, there are at least $|M_0|/q$ points in M_0 whose k_1 th coordinate is at most the k_1 th coordinate of p_{j_1} . Therefore, point p_i must have k_1 th coordinate larger than the k_1 th coordinate of these at least $|M_0|/q$ points in M_0 . Let M_1 denote those points in M_0 whose k_1 th coordinate is larger than the k_1 th coordinate of p_{j_1} , so $|M_1| \leq (1 - 1/q)|M_0|$.

After step h , we have a set M_h with $|M_h| \leq (1 - 1/q)^h |M_0|$ such that for every point $p_j \in M_0 \setminus M_h$, there is $k \in [q]$ such that the k th coordinate of p_i is guaranteed to be larger than the k th coordinate of p_j . If M_h is nonempty, then builder moves on to step $h + 1$ and picks a point $p_{j_{h+1}} \in M_h$ of position in M_h at least $|M_h|/q$. Such a point exists by the first part of Lemma 5.3. Coordinator then picks a coordinate k_{h+1} . By the definition of position, there are at least $|M_h|/q$ points in M_h whose k_{h+1} th coordinate is at most the k_{h+1} th coordinate of $p_{j_{h+1}}$. Therefore, point p_i must have k_{h+1} th coordinate larger than the k_{h+1} th coordinate of these at least $|M_h|/q$ points in M_h . Let M_{h+1} denote those points in M_h whose k_{h+1} th coordinate is larger than the k_{h+1} th coordinate of $p_{j_{h+1}}$, so $|M_{h+1}| \leq (1 - 1/q)|M_h|$.

If M_h is empty, then builder decides to end the stage and coordinator must pick a point p_i satisfying the required conditions. Note that we eventually will have an empty M_h as M_0 is a finite set, and M_h is a proper subset of M_{h-1} . Since in the previous step M_{h-1} is nonempty, we have

$$1 \leq |M_{h-1}| \leq (1 - 1/q)^{h-1} |M_0| \leq (1 - 1/q)^{h-1} (n - 1)^{q-1},$$

and so the number h of steps satisfies

$$h \leq 1 + (q - 1) \frac{\log n}{\log(\frac{q}{q-1})} = a(q, n).$$

Since M_0 are the maximal elements of the already constructed sequence, and for every point $p \in M_0$, there is $k \in [q]$ such that the k th coordinate of p_i is greater than the k th coordinate of p , then p_i cannot be an element of the already constructed sequence. This completes the proof. \square

We remark that one can do a little better in the above proof by improving the bound on $|M_0|$ by using a result of de Bruijn, van Ebbenhorst Tengbergen, and Kruyswijk [7] which extends Sperner's theorem. This result says that the set of elements in $[n - 1]^q$ whose coordinates sum to $\lfloor nq/2 \rfloor$ forms a maximum antichain in the poset on $[n - 1]^q$. Since M_0 is an antichain, the size of this set is an upper bound on $|M_0|$.

We next present a strategy for coordinator which gives a lower bound for $L(q, n)$.

Lemma 5.5. *For each fixed $q \geq 2$, we have $L(q, n) \geq (q - 1 - o(1)) n^q \log_q n$.*

Proof. We present a strategy for coordinator which satisfies that the number of steps in the stage the i th point not already in the sequence is added is at least a certain number depending on i which will be defined later in the proof. Level r of the grid $[n - 1]^q$ consists of all points in this grid whose coordinates sum to r . The new points will be added in levels, so no points of larger level are added until all points of lower level are already in the sequence.

At stage i , for some r , all points in levels j with $j < r$ are already in the sequence, and not all of level r is in the sequence. Let S_0 denote the set of points in level r not already in the sequence. As S_0 is disjoint from the already constructed sequence, and no element in the already constructed sequence is larger than any point in S_0 , then no matter what point p_{j_1} in the already constructed sequence builder chooses for the first step to compare p_i with, by the second part of Lemma 5.3, the position of p_{j_1} with respect to S_0 is at most $(1 - \frac{1}{q})|S_0|$. Hence, there is $k \in [q]$ such that at least $\frac{1}{q}|S_0|$ points in S_0 have k th coordinate more than the k th coordinate of p_{j_1} . Coordinator picks this coordinate k to compare with, and lets S_1 denote the points in S_0 whose k th coordinate is larger than the k th coordinate of p_{j_1} , so $|S_1| \geq |S_0|/q$.

After h steps of stage i , we have a subset S_h of level r with $|S_h| \geq |S_0|/q^h$ and every point in S_h satisfies the conditions that must be satisfied by p_i . Notice this is satisfied for $h = 1$. Builder has two options, either continue on to step $h + 1$, or declare the turn over.

In the first case, builder picks a point $p_{j_{h+1}}$ in the already constructed sequence to compare p_i with. By the second part of Lemma 5.3, the position of $p_{j_{h+1}}$ with respect to S_h is at most $(1 - \frac{1}{q})|S_h|$. Hence, there is $k \in [q]$ such that at least $\frac{1}{q}|S_h|$ points in S_h have k th coordinate more than the k th coordinate of $p_{j_{h+1}}$. Coordinator picks this coordinate k to compare with, and lets S_{h+1} denote the points in S_h whose k th coordinate is larger than the k th coordinate of $p_{j_{h+1}}$, so $|S_{h+1}| \geq |S_h|/q$. This finishes step $h + 1$.

In the second case, builder declares the turn is over after the h steps. There are two possibilities: $|S_h| = 1$ or $|S_h| > 1$. If $|S_h| = 1$, coordinator picks p_i to be the point in S_h . The number h of steps in this case satisfies $1 = |S_h| \geq |S_0|/q^h$, so $h \geq \log_q |S_0|$. Otherwise, $|S_h| > 1$, and we claim that there is still a point $z \in [n - 1]^q$ in a level less than r which can be chosen for p_i . Indeed, if $x = (x_1, \dots, x_q)$ and $y = (y_1, \dots, y_q)$ are distinct points in S_h , then the point $z = (z_1, \dots, z_q)$ with $z_j = \min(x_j, y_j)$ satisfies the desired properties. Indeed, as x and y are in level r , the level of z is $\sum_{i=1}^q \min(x_i, y_i) < \sum_{i=1}^q x_i = r$. Further, the set of possibilities for p_i is of the form that the k th coordinate is at least some m_k for each k , and, as x and y are in this set, then z is also in this set. The point z is already in the sequence, and coordinator lets p_i be this point. Since p_i is already in the sequence, after stage i builder is exactly in the same situation as before stage i in regards to winning the (q, n) -lattice game, and so the steps in this stage did not help get closer to ending the game.

In each stage for which a new point appears in the sequence at least $\log_q |S_0|$ steps were used, where $|S_0|$ is the number of points not yet in the sequence in the level currently being used to fill the new points. Let l_r denote the number of points in level r . We have $l_r = 1$ if and only if $r = q$

or $q(n-1)$. We thus have

$$\begin{aligned}
L(q, n) &\geq \sum_{r=q}^{q(n-1)} \sum_{m=1}^{l_r} \log_q m = \sum_{r=q+1}^{q(n-1)-1} \sum_{m=1}^{l_r} \log_q m \geq \sum_{r=q+1}^{q(n-1)-1} \int_1^{l_r-1} \log_q m \, dm \\
&= \sum_{r=q+1}^{q(n-1)-1} (m \log_q m - m / \ln q) \Big|_{m=1}^{l_r-1} = \sum_{r=q+1}^{q(n-1)-1} (l_r - 1) \log_q (l_r - 1) - (l_r - 2) / \ln q \\
&\geq ((q(n-2) - 1) \ell \log \ell - ((n-1)^q - 2 - 2(q(n-2) - 1) / \ln q)),
\end{aligned}$$

where in the second inequality we bounded a sum from below by an integral using the fact $f(x) = \log x$ is an increasing function, and the last inequality follows from Jensen's inequality and convexity of the function $f(x) = x \log x$, where we substitute $\sum_{r=q+1}^{q(n-1)-1} l_r = (n-1)^q - 2$, and $\ell = ((n-1)^q - 1 - q(n-2)) / (q(n-2) - 1)$ is the average of $l_r - 1$ with r ranging from $q+1$ to $q(n-1) - 1$. For q fixed, this bound can be simplified to

$$L(q, n) \geq (1 - o(1)) n^q \log_q n^{q-1} = (q - 1 - o(1)) n^q \log_q n,$$

which completes the proof. \square

5.2 Alternative upper bound on $N_k(q, n)$

In this subsection, we present the proof of Theorem 1.4, which gives an upper bound on $N_k(q, n)$ in terms of the online Ramsey number $V_{k-1}(q, n + k - 2)$.

Define $V'_k(q, n)$ exactly as $V_k(q, n)$, except in each stage $t \geq k$, builder must add at least one edge (whose largest vertex is v_t). We refer to the online Ramsey game in which builder must add at least one edge in each stage $t \geq k$ as the *modified online Ramsey game*. We have

$$V_k(q, n) \leq V'_k(q, n) \leq V_k(q, n + k - 1), \quad (3)$$

the lower bound being trivial. Note that in the usual online Ramsey graph, those vertices v_t which are not the largest vertex in an edge can only be among the first $k-1$ vertices of a monotone path. Hence, after a monochromatic monotone path of length $n + k - 1$ is created, the last n vertices of the path each are the largest vertex of some edge, which gives the upper bound. We will use this upper bound in the proof below.

Proof of Theorem 1.4. Let $\chi : \binom{[N]}{k} \rightarrow [q]$ be a q -coloring of the edges of K_N^k , where $N = q^{V_{k-1}(q, n+k-2)} + k - 2$. We have to show that K_N^k contains a monochromatic monotone path of length n .

We construct a set of vertices $\{v_1, v_2, \dots, v_t\}$ and a $(k-1)$ -uniform hypergraph H on these vertices with at most $V_{k-1}(q, n+k-2)$ edges such that for any $(k-1)$ -edge $e = \{v_{i_1}, \dots, v_{i_{k-1}}\}$, $i_1 < \dots < i_{k-1}$, the color of any k -edge $\{v_{i_1}, \dots, v_{i_{k-1}}, v_{i_k}\}$ in K_N^k with $i_k > i_{k-1}$ is the same, say $\chi'(e)$. Moreover, this $(k-1)$ -uniform hypergraph will contain a monochromatic path of length n , which one can easily see will define a monochromatic path of length n in K_N^k . As described below, the way the vertices of the set are constructed will be determined by playing the modified online Ramsey game.

We begin the construction of this set of vertices by setting $v_1 = 1$ and setting $S_1 = [N] \setminus \{v_1\}$. After stage t of the process, we have a set of vertices $\{v_1, \dots, v_t\}$ with $v_1 < \dots < v_t$ and a set S_t such

that $w > v_t$ for all $w \in S_t$ and for each $(k-1)$ -edge $\{v_{i_1}, \dots, v_{i_{k-1}}\}$ in H with $i_1 < \dots < i_{k-1} \leq t$, the color of the k -edge $\{v_{i_1}, \dots, v_{i_{k-1}}, w\}$ in K_N^k is the same for every w in S_t and for every $w = v_j$ with $i_{k-1} < j \leq t$.

In the beginning of stage $t+1$, we let v_{t+1} be the smallest element in S_t . We play the modified online Ramsey game, so that builder chooses the edges, which are $(k-1)$ -tuples $\{v_{i_1}, \dots, v_{i_{k-2}}, v_{t+1}\}$ with $i_1 < \dots < i_{k-2} < t+1$, to be drawn according to his strategy. Painter then colors these edges. For the first edge e_1 chosen, painter looks at all k -tuples containing this edge and a vertex from $S_t \setminus \{v_{t+1}\}$. The $(k-1)$ -edge is colored $r_1 \in [q]$ in χ' if there are at least $(|S_t| - 1)/q$ such k -tuples that have color r_1 , breaking ties between colors arbitrarily. Such a color r_1 exists by the pigeonhole principle. This defines a new subset $S_{t,1}$, which are all vertices in $S_t \setminus \{v_{t+1}\}$ such that together with edge e_1 form a k -tuple of color $r_1 = \chi'(e_1)$. After j edges have been drawn in this stage, we have a subset $S_{t,j}$, and we color the next drawn edge e_{j+1} color r_{j+1} if the number of k -tuples of color r_{j+1} containing it and a vertex from $S_{t,j}$ is at least $|S_{t,j}|/q$, breaking ties between colors arbitrarily. Then we define $S_{t,j+1}$ to be the set of at least $|S_{t,j}|/q$ vertices in $S_{t,j}$ which together with e_{j+1} form an edge of color r_{j+1} .

After builder has added all $(k-1)$ -edges with largest vertex v_{t+1} , the remaining set will be S_{t+1} . Let m_t be the number of $(k-1)$ -edges $e = \{v_{i_1}, \dots, v_{i_{k-2}}, v_t\}$ with $i_1 < \dots < i_{k-2} < t$ in H . Since we are playing the modified online Ramsey game, we have $m_t = 0$ for $1 \leq t \leq k-2$ and $m_t \geq 1$ for $t \geq k-1$.

We now show by induction that

$$|S_t| \geq \frac{N - (k-2)}{q^{\sum_{i=k-1}^t m_i}} - \sum_{i=k-1}^t \frac{1}{q^{\sum_{j=i}^t m_j}}.$$

For the base case $t = k-2$, we have

$$|S_{k-2}| = N - (k-2).$$

Suppose we have proved the desired inequality for t . When we draw a vertex v_{t+1} , the size of our set decreases by 1. Each time we draw an edge from v_{t+1} , the size of our set S goes down by a factor of q . Therefore

$$\begin{aligned} |S_{t+1}| &\geq \frac{|S_t| - 1}{q^{m_{t+1}}} \geq \frac{N - (k-2)}{q^{\sum_{i=k-1}^{t+1} m_i}} - \left(\sum_{i=k-1}^t \frac{1}{q^{\sum_{j=i}^{t+1} m_j}} \right) - \frac{1}{q^{m_{t+1}}} \\ &= \frac{N - (k-2)}{q^{\sum_{i=k-1}^{t+1} m_i}} - \sum_{i=k-1}^{t+1} \frac{1}{q^{\sum_{j=i}^{t+1} m_j}}. \end{aligned}$$

The number of edges drawn in the $(k-1)$ -uniform hypergraph after t stages is $\sum_{i=k-1}^t m_i$. If this is at most $V'_{k-1}(q, n) \leq V_{k-1}(q, n + k - 2)$, as $q \geq 2$ and $m_i \geq 1$ for all $i \geq k-1$, we have

$$\begin{aligned} |S_t| &\geq \frac{N - (k-2)}{q^{\sum_{i=k-1}^t m_i}} - \sum_{i=k-1}^t \frac{1}{q^{\sum_{j=i}^t m_j}} \geq \frac{N - (k-2)}{q^{V'_{k-1}(q, n)}} - \sum_{i=k-1}^t \frac{1}{q^{t-i+1}} \\ &> \frac{N - (k-2)}{q^{V'_{k-1}(q, n)}} - 1 = 0. \end{aligned}$$

Since $|S_t|$ is an integer, $|S_t| \geq 1$ and we can continue to the next stage. Thus, at the time we stop, $\sum_{i=1}^t m_i > V'_{k-1}(q, n)$ and we have constructed a $(k-1)$ -uniform hypergraph H that contains a monochromatic monotone path in coloring χ' on vertices v_{i_1}, \dots, v_{i_n} . On the other hand, it follows from the properties of our construction that the vertices v_{i_1}, \dots, v_{i_n} form a monochromatic monotone path of length n in K_N^k as required. \square

5.3 Size Ramsey numbers

We close this section by proving a lower bound on the size Ramsey number $S_2(q, n)$. Recall that $\chi(q, n)$ denotes the minimum chromatic number of an ordered graph G which is (q, n) -path Ramsey. A graph is t -degenerate if every subgraph of it has a vertex of degree at most t . Every t -degenerate graph has an ordering of its vertices such that each vertex is adjacent to at most t earlier vertices. Indeed, let a vertex of degree at most t be the last vertex in the ordering, and continue picking vertices out of degree at most t from the remaining induced subgraph, adding them to the end of the ordering, until all vertices are picked out. From this ordering, we get that every t -degenerate graph has chromatic number at most $t+1$ as we can color the vertices in order, picking the color of a vertex distinct the colors of its at most t neighbors that come before it in the ordering.

Theorem 5.6. *Let $n_1, n_2, q \geq 2$ be integers, and let $n = n_1 + n_2 - 1$. We have*

$$S_2(q, n) \geq (\chi(q, n_1) - 1)N_2(q, n_2)/2.$$

Proof. Let G be an ordered graph such that for every q -edge-coloring of G there is a monochromatic monotone path of length n . Suppose for contradiction that G has fewer than $(\chi(q, n_1) - 1)N_2(q, n_2)/2$ edges. Partition $V(G) = V_1 \cup V_2$, where V_2 is the t -core of G with $t = \chi(q, n_1) - 1$, which is formed by deleting, one-by-one, vertices from G of smallest degree until the remaining induced subgraph has minimum degree at least t . Since G has fewer than $(\chi(q, n_1) - 1)N_2(q, n_2)/2$ edges, and the minimum degree in the subgraph of G induced by the vertex set V_2 is at least t , we have the inequality $|V_2| < N_2(q, n_2)$, and hence there is a q -coloring with color set $[q]$ of the edges inside V_2 with no monochromatic monotone path of length n_2 . The subgraph of G induced by the vertices in V_1 is $(t-1)$ -degenerate, so it can be properly colored with $t = \chi(q, n_1) - 1$ colors. Hence, by definition of $\chi(q, n_1)$, there is a q -edge-coloring with color set $[q]$ of the edges inside V_1 without a monochromatic monotone path of length n_1 . We still have to color the edges between V_1 and V_2 . For any two vertices $i \in V_1$ and $j \in V_2$, if $i < j$, color the edge (i, j) color 1, and if $i > j$, color the edge (i, j) color 2. For each $j \in \{1, 2\}$, a monochromatic monotone path cannot start with a vertex in V_j , later have a vertex in V_{3-j} , and then, even later have a vertex in V_j , as otherwise it would contain an edge of color 1 and an edge of color 2. Therefore, the length of the longest monochromatic monotone path is at most the sum of the lengths of the longest monochromatic monotone paths within V_1 and within V_2 , which is at most $(n_1 - 1) + (n_2 - 1) = n - 1$, contradicting our assumption that every q -edge-coloring of G has a monochromatic monotone path of length n . The proof is complete. \square

Taking $n_1 = n_2 = n/2$ in the previous theorem, and using $N_2(q, n) = (n-1)^q + 1$ and Theorems 3.1 and 3.3, we have the following corollary.

Corollary 5.7. *For any integers $n, q \geq 2$, we have $S_2(q, n) \geq c_q n^{2q-1}$ for some constant $c_q > 0$ only depending on q .*

This should be compared with the trivial upper bound

$$S_2(q, n) \leq \binom{N_2(q, n)}{2} = \binom{(n-1)^q + 1}{2} \leq n^{2q}.$$

Thus, for fixed q , the lower and upper bound on $S_2(q, n)$ are roughly a factor n apart.

6 Transitive colorings

In this short section we prove a simple lemma which is used in the proof of Theorem 1.6. A *family* \mathcal{F} of k -element subsets of $[N]$ (i.e., a k -uniform hypergraph on the vertex set $[N]$) is said to be *transitive* if for any $i_1 < i_2 < \dots < i_{k+1}$ such that

$$\{i_1, i_2, \dots, i_k\}, \{i_2, i_3, \dots, i_{k+1}\} \in \mathcal{F},$$

we can conclude that *all* k -element subsets of $\{i_1, i_2, \dots, i_{k+1}\}$ belong to \mathcal{F} . A q -coloring of all k -element subsets of $[N]$ is called *transitive* if each of its color classes is transitive.

For the geometric application Theorem 1.6, we need the following lemma relating monochromatic cliques in transitive colorings to monochromatic monotone paths in general colorings.

Lemma 6.1. *Let $N = N_k(q, n)$ be the smallest integer such that for every q -coloring of all hyperedges of K_N^k , the complete k -uniform hypergraph on the vertex set $[N]$, there exists a monochromatic monotone path of length n . Then, for every transitive q -coloring of all hyperedges of K_N^k , there exists a monochromatic complete subhypergraph of size n .*

Lemma 6.1 directly follows from the definition of $N_k(q, n)$ and the following statement.

Lemma 6.2. *Let $n > k$, and let H be a k -uniform ordered hypergraph on the vertex set $[n]$, which contains a monotone path of length n , that is, $\{i, i+1, \dots, i+k-1\} \in E(H)$ for all $1 \leq i \leq n-k+1$.*

If $E(H)$ is transitive, then H is the complete k -uniform hypergraph on $[n]$.

Proof. We proceed by induction on n . The base case $n = k + 1$ follows from the definition of transitivity.

For the inductive step, suppose that $n > k + 1$ and that the statement has already been established for all ordered hypergraphs with fewer than n vertices. Let H be a hypergraph with n vertices meeting the conditions of the lemma. By the induction hypothesis, the induced subhypergraph $H[\{1, \dots, n-1\}] \subseteq H$, consisting of all edges of H contained in $\{1, \dots, n-1\}$ is complete, and so is the induced subhypergraph $H[\{2, \dots, n\}] \subseteq H$.

Observe that this implies that for any $1 \leq i \leq n$, the induced hypergraph $H[\{1, \dots, n\} \setminus \{i\}]$ contains a monotone path of length $n-1$. As the edge set of $H[\{1, \dots, n\} \setminus \{i\}]$ is clearly transitive, we can apply the induction hypothesis to conclude that $H[\{1, \dots, n\} \setminus \{i\}]$ is a complete k -uniform hypergraph, for all i . Therefore, H is also a complete k -uniform hypergraph. \square

7 Noncrossing convex bodies—Proof of Theorem 1.6

Let $\mathcal{C} = \{C_1, C_2, \dots, C_N\}$ be a family of $N = N_3(3, n)$ noncrossing convex bodies in general position in the plane, numbered from left to right, according to the x -coordinates of their left endpoints (leftmost points). We have to show that \mathcal{C} contains n members in convex position.

In order to complete the proof outlined at the end of Subsection 1.3 in the Introduction, we have to define a 3-coloring of all triples (C_i, C_j, C_k) with $i < j < k$, which satisfies some special properties. We need some definitions.

We say that the triple $(C_i, C_j, C_k), i < j < k$ has a *clockwise (counterclockwise) orientation* if there exist distinct points $q_i \in C_i, q_j \in C_j, q_k \in C_k$ such that they lie on the boundary of $\text{conv}(C_i \cup C_j \cup C_k)$ and appear there in clockwise (counterclockwise) order (cf. [29], [32]). Clearly, the orientation of a triple can be clockwise and counterclockwise at the same time. See Figure 1.




Figure 1: Orientations of convex bodies.

As we numbered the members of \mathcal{C} according to the order of their left endpoints, for any $i < j < k$, the left endpoint of C_i must lie on the boundary of $\text{conv}(C_i \cup C_j \cup C_k)$. The triple (C_i, C_j, C_k) is said to have a *strong-clockwise (strong-counterclockwise) orientation* if there exist points $q_j \in C_j, q_k \in C_k$ such that, starting at the left endpoint q_i^* of C_i , the triple (q_i^*, q_j, q_k) appears in clockwise (counterclockwise) order along the boundary of $\text{conv}(C_i \cup C_j \cup C_k)$. We say that (C_i, C_j, C_k) has *both strong orientations* if it has both a strong-clockwise and a strong-counterclockwise orientation. Notice that it is possible that (C_i, C_j, C_k) has both orientations, but only one *strong* orientation. On the other hand, every triple has at least one strong orientation. Obviously, if (C_i, C_j, C_k) has a strong-clockwise orientation, say, then it also has a clockwise orientation.

7.1 Definitions, notation, and observations

Let $\mathcal{C} = \{C_1, C_2, \dots, C_N\}$ be a family of noncrossing convex bodies in general position, numbered from left to right, according to the x -coordinates of their left endpoints (leftmost points). A subfamily of convex bodies \mathcal{S} is called *separable* if it has a member C such that $\text{conv}(\cup \mathcal{S}) \setminus C$ is disconnected. We say that C *separates* \mathcal{S} if $\text{conv}(\cup \mathcal{S}) \setminus C$ is disconnected. Clearly if a triple is not separable, then it only has one strong orientation. Since \mathcal{C} is a family of noncrossing convex sets, every triple has at most one member that separates it.

We will make several observations on the triples of \mathcal{C} using the following notation. For $i < j < k$, let β denote the boundary of $\text{conv}(C_i \cup C_j \cup C_k)$, q_i^* denote the left endpoint of C_i , and let $\gamma_i = C_i \cap \beta, \gamma_j = C_j \cap \beta, \gamma_k = C_k \cap \beta$.

Observation 7.1. For $i < j < k$, C_k cannot separate (C_i, C_j, C_k) .

Proof. For sake of contradiction, suppose that C_k separates the triple (C_i, C_j, C_k) . Then γ_i is a simple continuous curve (i.e., homeomorphic to the unit interval) that contains the left most point of β , γ_j is also a simple continuous curves, and γ_k consists of two disjoint simple continuous curves γ_{k_1} and γ_{k_2} . Since no two convex bodies have a common tangent that meets at their intersection, $\gamma_i, \gamma_j, \gamma_{k_1}, \gamma_{k_2}$ are pairwise disjoint. Moreover, $\gamma_i, \gamma_{k_1}, \gamma_j, \gamma_{k_2}$ appear in clockwise order along β .

However since $q_i^* \in \gamma_i$ and since the left endpoint of C_j lies inside $\text{conv}(C_i, C_j, C_k)$ to the left of the left endpoint of C_k , this implies that C_j and C_k cross, that is, they share more than two boundary points which is a contradiction. \square

Observation 7.2. *For $i < j < k$, the triple (C_i, C_j, C_k) has both strong orientations if and only if C_j separates it.*

Proof. If C_j separates the triple (C_i, C_j, C_k) , then γ_i and γ_k are simple continuous curves such that $q_i^* \in \gamma_i$, while γ_j consists of two disjoint simple continuous curves $\gamma_{j_1}, \gamma_{j_2}$. Moreover since \mathcal{C} is in general position, $\gamma_i, \gamma_{j_1}, \gamma_k, \gamma_{j_2}$ are pairwise disjoint and appear in clockwise order along β . Therefore we can find points $q_{j_1} \in \gamma_{j_1}, q_{j_2} \in \gamma_{j_2}$, and $q_k \in \gamma_k$ such that $q_i^*, q_{j_1}, q_k, q_{j_2}$ appear on β in clockwise order. Hence (C_i, C_j, C_k) has both strong orientations.

For the other direction, suppose that (C_i, C_j, C_k) has both strong orientations. Since the triple (C_i, C_j, C_k) is separable, and we know that C_k does not separate it, it suffices to show that C_i does not separate (C_i, C_j, C_k) .

Suppose C_i separates (C_i, C_j, C_k) . Then γ_j and γ_k are simple continuous curves, and γ_i consists of two disjoint simple continuous curves γ_{i_1} and γ_{i_2} . We can assume that $q_i^* \in \gamma_{i_1}$. Since C_i separates (C_i, C_j, C_k) , $\gamma_{i_1}, \gamma_j, \gamma_{i_2}, \gamma_k$ are pairwise disjoint and must appear in clockwise or counterclockwise order along β . If $\gamma_{i_1}, \gamma_j, \gamma_{i_2}, \gamma_k$ appear in clockwise order, then (C_i, C_j, C_k) has only a strong-clockwise orientation since there does not exist points $q_j \in \gamma_j$ and $q_k \in \gamma_k$ such that q_i^*, q_j, q_k appear in counterclockwise order along β . Likewise if $\gamma_{i_1}, \gamma_j, \gamma_{i_2}, \gamma_k$ appear in counterclockwise order along β , then (C_i, C_j, C_k) has only a strong-counterclockwise orientation. In either case, (C_i, C_j, C_k) does not have both strong orientations and we have a contradiction. \square

Suppose that C_i separates (C_i, C_j, C_k) . By the argument above (C_i, C_j, C_k) has only one strong orientation since $\gamma_{i_1}, \gamma_j, \gamma_{i_2}, \gamma_k$ must appear in clockwise or counterclockwise order along β . Let h be a fixed line that goes through q_i^* and intersects γ_{i_2} . We can assume that h is not vertical. Then by the noncrossing property and by convexity of $\text{conv}(C_i \cup C_j \cup C_k)$, regions $C_j \setminus C_i$ and $C_k \setminus C_i$ must be separated by h . Therefore we make the following observation.

Observation 7.3. *For $i < j < k$, suppose that C_i separates the triple (C_i, C_j, C_k) and let h be the line described as above. If $C_j \setminus C_i$ lies in the upper half-plane generated by h (and $C_k \setminus C_i$ lies in the lower half-plane), then (C_i, C_j, C_k) has only a strong-clockwise orientation. Likewise if $C_j \setminus C_i$ lies in the lower half-plane generated by h (and $C_k \setminus C_i$ lies in the upper half-plane), then (C_i, C_j, C_k) has only a strong-counterclockwise orientation.*

Proof. If $C_j \setminus C_i$ lies in the upper half-plane generated by h , then $\gamma_{i_1}, \gamma_j, \gamma_{i_2}, \gamma_k$ must appear in clockwise order along β . Hence there does not exist points $q_j \in \gamma_j$ and $q_k \in \gamma_k$ such that q_i^*, q_j, q_k appear in counterclockwise order along β . Therefore (C_i, C_j, C_k) does not have a strong-counterclockwise orientation, which implies that it only has a strong-clockwise orientation since every triple has at least one strong orientation.

By a similar argument, if $C_j \setminus C_i$ lies in the lower half-plane generated by h , this implies that (C_i, C_j, C_k) has only a strong-counterclockwise orientation. \square

For each pair of indices $i < j$, let \vec{l}_{ij} (and \vec{r}_{ij}) denote the *directed* common tangent line of C_i and C_j which meets C_i before C_j , such that $C_i \cup C_j$ lies completely to the right (left) of \vec{l}_{ij} (\vec{r}_{ij}).

Let \mathcal{C} be a family of noncrossing convex sets in general position, and let β denote the boundary of $\text{conv}(\cup \mathcal{C})$. We say that the pair (C_i, C_j) appears in *consecutive clockwise (counterclockwise) order* on β if there exists points $q_i \in C_i \cap \beta$ and $q_j \in C_j \cap \beta$ such that if γ is the arc generated by moving from q_i to q_j along β in the clockwise (counterclockwise) direction, then the interior of γ does not intersect any member of \mathcal{C} .

If (C_i, C_j) appears in consecutive clockwise (counterclockwise) order on β , then \vec{l}_{ij} (\vec{r}_{ij}) is the directed line going through points q_i and q_j (where q_i, q_j is defined as above) such that $\text{conv}(\cup \mathcal{C})$ lies to the right (left) of \vec{l}_{ij} (\vec{r}_{ij}). Therefore we have the following.

Observation 7.4. *For $i < j < k$, if (C_i, C_j, C_k) has both strong orientations, then $C_i \cup C_j \cup C_k$ lies to the right of \vec{l}_{ij} and \vec{l}_{jk} , and lies to the left of \vec{r}_{ij} and \vec{r}_{jk} . In other words, (C_i, C_j) (and (C_j, C_k)) appears in consecutive clockwise and counterclockwise order on the boundary of $\text{conv}(C_i, C_j, C_k)$.*

Proof. By Observation 7.2, C_j separates (C_i, C_j, C_k) . Therefore γ_i and γ_k are simple continuous curves while γ_j consists of two disjoint simple continuous curves γ_{j_1} and γ_{j_2} . Moreover, $\gamma_i, \gamma_{j_1}, \gamma_k, \gamma_{j_2}$ are pairwise disjoint and appear in clockwise order along the boundary of $\text{conv}(C_i \cup C_j \cup C_k)$. Hence (C_i, C_j) (and (C_j, C_k)) appears in consecutive clockwise and counterclockwise order on the boundary of $\text{conv}(C_i, C_j, C_k)$. \square

Observation 7.5. *For $i < j < k$, if (C_i, C_j, C_k) has only a strong-clockwise orientation, then $C_i \cup C_j \cup C_k$ lies to the right of \vec{l}_{ij} and to the left of \vec{r}_{ik} . In other words, (C_i, C_j) appears in consecutive clockwise order and (C_i, C_k) appears in consecutive counterclockwise order on the boundary of $\text{conv}(C_i \cup C_j \cup C_k)$.*

Proof. By Observation 7.1 and 7.2, either (C_i, C_j, C_k) is not separable or C_i separates it. If (C_i, C_j, C_k) is not separable, then $\gamma_i, \gamma_j, \gamma_k$ are pairwise disjoint simple continuous curves that appear on the boundary of $\text{conv}(C_i \cup C_j \cup C_k)$ in clockwise order. Hence (C_i, C_j) appears in consecutive clockwise order and (C_i, C_k) appears in consecutive counterclockwise order on the boundary of $\text{conv}(C_i \cup C_j \cup C_k)$ and we are done.

If C_i separates (C_i, C_j, C_k) , then γ_j and γ_k are simple continuous curves while γ_i consists of two disjoint simple continuous curves γ_{i_1} and γ_{i_2} . Moreover $\gamma_{i_1}, \gamma_j, \gamma_{i_2}, \gamma_k$ are pairwise disjoint and appear in clockwise order. Hence (C_i, C_j) appears in consecutive clockwise order and (C_i, C_k) appears in consecutive counterclockwise order on the boundary of $\text{conv}(C_i \cup C_j \cup C_k)$. \square

By a symmetric argument, we have the following.

Observation 7.6. *For $i < j < k$, if (C_i, C_j, C_k) has only a strong-counterclockwise orientation, then $C_i \cup C_j \cup C_k$ lies to the left of \vec{r}_{ij} and to the right of \vec{l}_{ik} .*

7.2 The transitive property and its application

Lemma 7.7. *Let $i < j < k < l$. If the triples (C_i, C_j, C_k) and (C_j, C_k, C_l) have both strong orientations, then the same is true for (C_i, C_j, C_l) and (C_i, C_k, C_l) .*

Proof. By Observations 7.2 and 7.4, both C_i and C_l must lie to the right of \vec{l}_{jk} and to the left of \vec{r}_{jk} as shown in Figure 2.

Hence C_j separates the triple (C_i, C_j, C_l) and C_k separates the triple (C_i, C_k, C_l) , which implies (C_i, C_j, C_l) and (C_i, C_k, C_l) have both strong orientations. \square




Figure 2: For Lemma 7.7.

Lemma 7.8. *Let $i < j < k < l$. If the triples (C_i, C_j, C_k) and (C_j, C_k, C_l) have only strong-clockwise orientations (or only strong-counterclockwise orientations), then the same is true for (C_i, C_j, C_l) and (C_i, C_k, C_l) .*

Proof. By symmetry, it is sufficient to verify the statement in the case when the triples (C_i, C_j, C_k) and (C_j, C_k, C_l) have only strong-clockwise orientations.

Claim 7.9. *(C_i, C_k, C_l) has only a strong-clockwise orientation.*

Notice that we can make the following assumption.

$$(*) \quad C_l \text{ lies to the left of } \vec{r}_{ik} \text{ and to the right of } \vec{l}_{jk}.$$

Indeed, by Observations 7.4 and 7.6, we can assume that C_l lies to the left of \vec{r}_{ik} since otherwise (C_i, C_k, C_l) would not have a strong-counterclockwise orientation and we would be done, that is, (C_i, C_k, C_l) has only a strong-clockwise orientation since every triple has at least one strong orientation. Moreover by Observation 7.5, C_l must lie to the right of \vec{l}_{jk} since (C_j, C_k, C_l) has only a strong-clockwise orientation.

We define L to be the vertical line through the left endpoint of C_k , p_1 to be the left endpoint of C_k , p_2 to be the first point on \vec{l}_{jk} from C_k in the direction of \vec{l}_{jk} , p_3 to be the first point on \vec{r}_{ik} from C_k in the direction of \vec{r}_{ik} , and p_4 to be the first point on \vec{r}_{jk} from C_k in the direction of \vec{r}_{jk} . For simplicity, we will assume that p_1, p_2, p_3, p_4 are distinct, since otherwise it would just make the remaining analysis easier.

Observation 7.10. p_1, p_2, p_3, p_4 appear in clockwise order along the boundary of C_k .

Proof. Let ∂ denote the boundary of $\text{conv}(C_j \cup C_k)$. Since C_j and C_k are noncrossing, $C_k \cap \partial$ is the arc generated by moving along the boundary of C_k from p_2 to p_4 in the clockwise direction. Since the left endpoint of C_j lies to the left of L and since p_1, p_2, p_4 are all distinct, p_1 must lie in the interior of $\text{conv}(C_j \cup C_k)$. Hence p_1, p_2, p_4 appear in clockwise order along the boundary of C_k . Therefore it suffices to show that points p_2, p_3, p_4 appear in clockwise order. By Observation 7.5, C_j must lie to the left of \vec{r}_{ik} and therefore $p_3 \in C_k \cap \partial$. Since p_2, p_3, p_4 are distinct, p_2, p_3, p_4 appear in clockwise order along the boundary of C_k and this completes the proof. \square

By the ordering on \mathcal{C} and by $(*)$, C_l must lie to the right of L , to the right of \vec{l}_{jk} , and to the left of \vec{r}_{ik} . Hence $C_l \setminus C_k$ must lie in one of the three regions defined as follows. Let *Region I* be the region enclosed by L , \vec{l}_{jk} , and the arc generated by moving from p_1 to p_2 along the boundary

of C_k in the clockwise direction, whose interior is disjoint from C_k . Let *Region II* be the region enclosed by \vec{l}_{jk} , \vec{r}_{ik} , and the arc generated by moving from p_2 to p_3 along the boundary of C_k in the clockwise direction, whose interior is disjoint from C_k . Let *Region III* be the region enclosed by L , \vec{r}_{ik} , and the arc generated by moving from p_3 to p_1 along the boundary of C_k in the clockwise direction, whose interior is disjoint from C_k . See Figure 3(a).

Observation 7.11. $C_l \setminus C_k$ does not lie in *Region I*.

Proof. For sake of contradiction, suppose $C_l \setminus C_k$ lies in *Region I*. Then the proof falls into two cases.

Case 1. Suppose that C_j does not intersect *Region I*. This implies that the tangent point of C_j on \vec{l}_{jk} lies to the left of L . Therefore, the triangle T_1 whose vertices is this tangent point, p_1 , and p_2 contains *Region I*. Since the vertices of T_1 are in $C_j \cup C_k$, the triangle T_1 is a subset of the convex hull $\text{conv}(C_j \cup C_k)$. Hence,

$$C_l \subset (\text{Region I} \cup C_k) \subset (T_1 \cup C_k) \subset \text{conv}(C_j \cup C_k),$$

which is a contradiction.

Case 2. Suppose C_j does intersect *Region I*. Let h be the line that goes through the left endpoint of C_j and a point on $C_j \cap \vec{l}_{jk}$. If $C_l \setminus C_j$ lies below h , then $C_l \subset \text{conv}(C_j \cup C_k)$ which is a contradiction. If $C_l \setminus C_k$ lies above h , then C_j separates (C_j, C_k, C_l) and by Observation 7.3, (C_j, C_k, C_l) has a strong counter-clockwise orientation which is also a contradiction. See Figure 3(b).




Figure 3: *Regions I, II, and III.*

□

Observation 7.12. $C_l \setminus C_k$ does not lie in *Region II*.

Proof. For sake of contradiction, suppose $C_l \setminus C_k$ lies in *Region II*. By Observation 7.10, p_2, p_3, p_4 appear in clockwise order. Hence C_l lies to the left of \vec{r}_{jk} . This implies that C_k separates the triple

(C_j, C_k, C_l) . By Observation 7.2, (C_j, C_k, C_l) has both strong orientations which is a contradiction. See Figure 4(a). □

Therefore $C_l \setminus C_k$ must lie in Region III. Suppose for contradiction that C_i does not intersect Region III. Then the tangent point of C_i on \vec{r}_{ik} lies to the left of L . Therefore, the triangle T_2 whose vertices is this tangent point, p_1 , and p_3 contains Region III. Since the vertices of T_2 are in $C_i \cup C_k$, the triangle T_2 is a subset of the convex hull $\text{conv}(C_i \cup C_k)$. Hence,

$$C_l \subset (\text{Region III} \cup C_k) \subset (T_2 \cup C_k) \subset \text{conv}(C_i \cup C_k),$$

which is a contradiction. Hence, C_i must intersect Region III. Moreover, $C_l \setminus C_i$ must lie in the lower half-plane generated by the line that goes through the left endpoint of C_i and a point on $C_i \cap \vec{r}_{ik}$ as otherwise again $C_l \subset \text{conv}(C_i \cup C_k)$. Since C_i separates (C_i, C_k, C_l) and by Observation 7.3, (C_i, C_k, C_l) has only a strong-clockwise orientation. See Figure 4(b). This completes the proof of Claim 7.9.




Figure 4: Regions II and III.

Now we will show that (C_i, C_j, C_l) has only a strong-clockwise orientation by a very similar argument. □

Claim 7.13. (C_i, C_j, C_l) has only a strong-clockwise orientation.

Just as before, we can assume the following.

$$(**) \quad C_l \text{ lies to the left of } \vec{r}_{ij} \text{ and to the right of } \vec{l}_{jk}.$$

Indeed, by Observation 7.6, we can assume that C_l lies to the left of \vec{r}_{ij} since otherwise (C_i, C_j, C_l) would not have a strong-counterclockwise orientation and we would be done, that is, (C_i, C_j, C_l) has only a strong-clockwise orientation since every triple has at least one strong orientation. Moreover

by Observation 7.5, C_l must lie to the right of \vec{l}_{jk} since (C_j, C_k, C_l) has only a strong-clockwise orientation.

Since (C_i, C_j, C_k) has only a strong-clockwise orientation, then either (C_i, C_j, C_k) is not separable, or C_i separates it. Now the proof falls into two cases.

CASE 1: Assume that (C_i, C_j, C_k) is not separable. Then we define L' to be the vertical line through the left endpoint of C_j , p_5 to be the left endpoint of C_j , p_6 to be the first point on \vec{l}_{jk} from C_j in the direction of \vec{l}_{jk} , p_7 to be the first point on \vec{r}_{ij} from C_j in the direction of \vec{r}_{ij} , and p_8 to be the first point on \vec{r}_{jk} from C_j in the direction of \vec{r}_{jk} . For simplicity, we will assume that p_5, p_6, p_7, p_8 are distinct, since otherwise it would just make the remaining analysis easier. Now we make the following observation.

Observation 7.14. p_5, p_6, p_7, p_8 appear in clockwise order along the boundary of C_j .

Proof. Let ∂ denote the boundary of $\text{conv}(C_j \cup C_k)$. Then $C_j \cap \partial$ is the arc generated by moving along the boundary of C_j from p_8 to p_6 in the clockwise direction. Since p_5 is the leftmost point of $\text{conv}(C_j \cup C_k)$, $p_5 \in C_j \cap \partial$. Hence p_5, p_6, p_8 appear in clockwise order along the boundary of C_j . Therefore it suffices to show that p_6, p_7, p_8 appear in clockwise order. Since (C_i, C_j, C_k) is not separable and has a strong-clockwise orientation, C_k does not lie completely to the left of \vec{r}_{ij} . Since p_6, p_7, p_8 are all distinct, p_7 lies in the interior of $\text{conv}(C_j \cup C_k)$. Thus p_6, p_7, p_8 appear in clockwise order along the boundary of C_j and this completes the proof. \square

By (**) and the ordering on \mathcal{C} , C_l must lie to the right of L' , to the left of \vec{r}_{ij} , and to the right of \vec{l}_{jk} . Therefore $C_l \setminus C_j$ must lie in one of the three regions defined as follows. Let *Region IV* be the region enclosed by \vec{l}_{jk} , L' , and the arc generated by moving from p_5 to p_6 along the boundary of C_j in the clockwise direction, whose interior is disjoint from C_j . Let *Region V* be the region enclosed by \vec{l}_{jk} , \vec{r}_{ij} , and the arc generated by moving from p_6 to p_7 along the boundary of C_j in the clockwise direction, whose interior is disjoint from C_j . Let *Region VI* be the region enclosed by L' , \vec{r}_{ij} , and the arc generated by moving from p_7 to p_5 along the boundary of C_j in the clockwise direction, whose interior is disjoint from C_j . See Figures 5(a) and 5(b).




Figure 5: Regions IV, V, and VI.

Observation 7.15. $C_l \setminus C_j$ cannot lie in Region IV

Proof. For sake of contradiction, assume that $C_l \setminus C_j$ does lie in Region IV. Since p_5, p_6, p_8 appear in clockwise order along the boundary of C_j , C_j separates the triple (C_j, C_k, C_l) . Moreover, $C_l \setminus C_j$ lies in the upper half-plane generated by the line that goes through p_5 and p_6 . By Observation 7.3, (C_j, C_k, C_l) has only a strong-counterclockwise orientation and we have a contradiction. See Figure 6(a). □

Observation 7.16. $C_l \setminus C_j$ cannot lie in Region V.

Proof. For sake of contradiction, assume that $C_l \setminus C_j$ does lie in Region V. Since C_l lies to the left of \vec{r}_{ij} and since p_7, p_8 appear in clockwise order along the boundary of C_j , we can conclude that C_l lies to the left of \vec{r}_{jk} . Since $C_l \not\subset \text{conv}(C_j \cup C_k)$, C_k separates the triple (C_j, C_k, C_l) . By Observation 7.2, (C_j, C_k, C_l) has both strong orientations and we have a contradiction. See Figure 6(b). □

Therefore $C_l \setminus C_j$ must lie in Region VI. Suppose for contradiction that C_i does not intersect Region VI. Then the tangent point of C_i on \vec{r}_{ij} lies to the left of L' . Therefore, the triangle T_3 whose vertices is this tangent point, p_5 , and p_7 contains Region VI. Since the vertices of T_3 are in $C_i \cup C_j$, the triangle T_3 is a subset of the convex hull $\text{conv}(C_i \cup C_j)$. Hence,

$$C_l \subset (\text{Region VI} \cup C_j) \subset (T_3 \cup C_j) \subset \text{conv}(C_i \cup C_j),$$

which is a contradiction. Therefore C_i must intersect Region VI and $C_l \setminus C_i$ must lie in the lower half-plane generated by the line that goes through the left endpoint of C_i and a point on $C_i \cap \vec{r}_{ij}$. Therefore C_i separates (C_i, C_j, C_l) and by Observation 7.3, (C_i, C_j, C_l) has only a strong-clockwise orientation. See Figure 6(c).

CASE 2: Assume that C_i separates (C_i, C_j, C_k) .

Observation 7.17. p_5, p_7, p_6 appear in clockwise order along the boundary of C_j .

Proof. Let β denote the boundary of $\text{conv}(C_i \cup C_j)$. Since C_i separates (C_i, C_j, C_k) , the arc generated by moving along the boundary of C_j from p_5 to p_6 in the clockwise direction must contain $C_j \cap \beta$. Since $p_7 \in C_j \cap \beta$, p_5, p_7, p_6 appear in clockwise order along the boundary of C_j . □

By (***) and the ordering on \mathcal{C} , C_l must lie to the right of L' and to the right of \vec{l}_{jk} . Hence $C_l \setminus C_j$ must lie in one of the following two regions. Let *Region VII* be the region enclosed by \vec{l}_{jk} , L' , and the arc generated by moving from p_5 to p_6 along the boundary of C_j in the clockwise direction, whose interior is disjoint from C_j . Finally let *Region VIII* be the region enclosed by L' , \vec{l}_{jk} , and the arc generated by moving from p_6 to p_5 along the boundary of C_j in the clockwise direction, whose interior is disjoint from C_j . See Figures 7(a) and 7(b).

Observation 7.18. $C_l \setminus C_j$ cannot lie in Regions VII.

Proof. For sake of contradiction, suppose $C_l \setminus C_j$ lies in Region VII. Then C_j separates the triple (C_j, C_k, C_l) , and $C_l \setminus C_j$ lies in the upper half-plane generated by the line that goes through the




Figure 6: Regions IV, V, and VI.

points p_5 and p_6 . By Observation 7.3, (C_j, C_k, C_l) has only a strong-counterclockwise orientation and we have a contradiction. See Figure 8(a). □

Therefore $C_l \setminus C_j$ must lie in Region VIII. Since C_i separates the triple (C_i, C_j, C_k) , which has only a strong-clockwise orientation, C_i must intersect Region VIII and \vec{l}_{jk} . Since $C_l \notin \text{conv}(C_i \cup C_j)$, $C_l \setminus C_i$ must lie in the lower half plane generated by the line that goes through the left endpoint of C_i and a point on $C_i \cap \vec{r}_{ij}$. Thus C_i separates (C_i, C_j, C_l) and by Observation 7.3, (C_i, C_j, C_l) has only a strong-clockwise orientation. See Figure 8(b). This completes the proof of Claim 7.13 and Lemma 7.8. □

Now we are ready to define a 3-coloring of the complete 3-uniform ordered hypergraph $K_{\mathcal{C}}^3$ formed by all triples in $\mathcal{C} = \{C_1, \dots, C_N\}$. A triple (C_i, C_j, C_k) , $i < j < k$ is colored

1. *red* if it has only a strong-clockwise orientation;
2. *blue* if it has only a strong-counterclockwise orientation;
3. *green* if it has both strong orientations.

Lemmas 7.7 and 7.8 imply that the above coloring is transitive.

Since $N = N_3(3, n)$, the edges of $K_{\mathcal{C}}^3$, i.e., the triples of \mathcal{C} , determine a monochromatic monotone path of length n . As this coloring is transitive, Lemma 6.2 implies that the members of this path induce a monochromatic *complete* 3-uniform subhypergraph $K \subseteq K_{\mathcal{C}}^3$. It follows from the definition of the coloring that all hyperedges of K are triples which have the same strong orientations. The proof of Theorem 1.6 is now a direct consequence of the following statement from [23] and Theorem 1.2, which gives an upper bound on $N = N_3(3, n)$.




Figure 7: C_i separates (C_i, C_j, C_k) .




Figure 8: Regions VII and VIII.

Lemma 7.19. *If a family of noncrossing convex bodies in general position in the plane can be ordered in such a way that every triple has a clockwise orientation (or every triple has a counterclockwise orientation), then all members of the family are in convex position.*

□

8 Concluding Remarks

1. We proved estimates on the minimum number of vertices $N_k(q, n)$ of a complete ordered k -uniform hypergraph for which every q -coloring of its edges has a monochromatic monotone path of length n . In particular, for $q \geq 3$ and $n \geq q + 2$, we have

$$2^{(n/q)^{q-1}} \leq N_3(q, n) \leq n^{n^{q-1}}.$$

It would be interesting to close the gap between the lower and upper bound.

2. We proved that the smallest integer $M(n)$ such that every family of $M(n)$ noncrossing convex bodies in general position in the plane has n members in convex position satisfies $M(n) \leq N_3(3, n) \leq$

n^2 . It would be interesting to improve the bound on $M(n)$ further and, in particular, decide whether $M(n)$ grow exponential in n .

3. We proved that triples of noncrossing convex bodies with *only* a strong-clockwise orientation have the transitive property. However it is easy to see that triples with a strong-clockwise orientation do not. In particular, it is possible to have four convex bodies ordered from left to right according to their left endpoints, such that (C_1, C_2, C_3) and (C_2, C_3, C_4) have a strong-clockwise orientation, but (C_1, C_3, C_4) does not as shown in Figure 9.




Figure 9: (C_1, C_3, C_4) does not have a strong-clockwise orientation.

4. We proved that the vertex online Ramsey number $V_2(q, n)$ and the lattice game number $L(q, n)$ are equal, asymptotically determined these numbers for $q = 2$, and determined the order of growth of these numbers for fixed q . We leave as an open problem to asymptotically determine these numbers for fixed $q > 2$.

5. We proved that the size Ramsey number $S_2(q, n)$, which is the minimum number of edges of a graph which is (q, n) -path Ramsey, satisfies $c_q n^{2q-1} \leq S_2(q, n) \leq c'_q n^{2q}$ for some constants c_q, c'_q depending on n . It would be interesting to determine the growth of $S_2(q, n)$ for fixed q .

References

- [1] Balogh, J., Butterfield, J.: Online Ramsey games for triangles in random graphs, *Discrete Math.* **310** (2010), 3653–3657.
- [2] Beck, J.: On size Ramsey number of paths, trees and circuits. I, *J. Graph Theory* **7** (1983), 115–129.
- [3] Beck, J.: Achievement games and the probabilistic method, *Combinatorics, Paul Erdős is Eighty*, Bolyai Soc. Math. Stud. **1** (1993), 51–78.
- [4] Belfrage, M., Mütze, T., Spohel, R.: Probabilistic one-player Ramsey games via deterministic two-player games, submitted.
- [5] Bisztriczky, T., Fejes Tóth, G.: A generalization of the Erdős-Szekeres convex n -gon theorem, *Journal für die reine und angewandte Mathematik* **395** (1989), 167–170.
- [6] Bisztriczky, T., Fejes Tóth, G.: Convexly independent sets, *Combinatorica* **10** (1990), 195–202.
- [7] de Bruijn, N. G., van Ebbenhorst Tengbergen, C., Kruyswijk, D.: On the set of divisors of a number, *Nieuw Arch. Wiskunde* **23** (1951), 191–193.

- [8] Conlon, D.: On-line Ramsey numbers, *SIAM J. Discrete Math.* **23** (2009), 1954–1963.
- [9] Conlon, D., Fox, J., Sudakov, B.: Ramsey numbers of sparse hypergraphs, *Random Struct. Algorithms* **35** (2009), 1–14.
- [10] Conlon, D., Fox, J., Sudakov, B.: Hypergraph Ramsey numbers, *Journal of the American Mathematical Society* **23** (2010), 247–266.
- [11] Conlon, D., Fox, J., Sudakov, B.: An improved bound for the stepping-up lemma, *Discrete Applied Mathematics*, to appear.
- [12] Cooley, O., Fountoulakis, N., Kühn, D., Osthus, D.: Embeddings and Ramsey numbers of sparse k -uniform hypergraphs, *Combinatorica* **29** (2009), 263–297.
- [13] Dilworth, R. P.: A decomposition theorem for partially ordered sets, *Annals of Mathematics* **51** (1950), 161–166.
- [14] Erdős, P., Faudree, R. J., Rousseau, C. C., Schelp, R. C.: The size Ramsey number, *Period. Math. Hungar.* **9** (1978), 145–161.
- [15] Erdős, P., Gallai, T.: On maximal paths and circuits of graphs, *Acta Math. Acad. Sci. Hungar* **10** (1959), 337–356.
- [16] Erdős, P., Szekeres, G.: A combinatorial problem in geometry, *Compositio Math.* **2** (1935), 463–470.
- [17] Figaj, A., Łuczak, T.: The Ramsey number for a triple of long even cycles, *J. Combin. Theory Ser. B* **97** (2007), 584–596.
- [18] Friedgut, E., Kohayakawa, Y., Rödl, V., Ruciński, A.: and P. Tetali, Ramsey games against a one-armed bandit, *Combin. Prob. Comp.* **12** (2003), 515–545.
- [19] Gerencsér, L., Gyárfás, A.: On Ramsey-type problems, *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.* **10** (1967), 167–170.
- [20] Graham, R. L., Rothschild, B. L., Spencer, J. H.: *Ramsey Theory, 2nd ed.*, Wiley, New York, 1990.
- [21] Gyárfás, A., Ruszinkó, M., Sárközy, G. N., Szemerédi, E.: Three-color Ramsey numbers for paths, *Combinatorica* **27** (2007), 35–69.
- [22] Haxell, P. E., Łuczak, T., Peng, Y., Rödl, V., Rućinski, A., Skokan, J.: The Ramsey number for 3-uniform tight hypergraph cycles, *Combin. Probab. Comput.* **18** (2009), 165–203.
- [23] Hubard, A., Montejano, L., Mora, E., Suk, A.: Order types of convex bodies, *Order*, DOI: 10.1007/s11083-010-9156-2.
- [24] Kohayakawa, Y., Rödl, V., Schacht, M., Szemerédi, E.: Sparse partition universal graphs for graphs of bounded degree, *Adv. Math.* **226** (2011), 5041–5065.
- [25] Kurek, A., Ruciński, A.: Two variants of the size Ramsey number, *Discuss. Math. Graph Theory* **25** (2005), 141–149.

- [26] Nagle, B., Olsen, S., Rödl, V., Schacht, M.: On the Ramsey number of sparse 3-graphs, *Graphs Combin.* **24** (2008), 205–228.
- [27] Pach, J., Tóth, G.: A generalization of the Erdős-Szekeres theorem to disjoint convex sets, *Discrete and Computational Geometry* **19** (1998), 437–445.
- [28] Pach, J., Tóth, G.: Erdős-Szekeres-type theorems for segments and noncrossing convex sets, *Geometriae Dedicata* **81** (2000), 1–12.
- [29] Pach, J., Tóth, G.: Families of convex sets not representable by points, in: *Indian Statistical Institute Platinum Jubilee Commemorative Volume—Architecture and Algorithms*, World Scientific, Singapore, 2009, 43–53.
- [30] Ramsey, F. P.: On a problem in formal logic, *Proc. London Math. Soc.* **30** (1930), 264–286.
- [31] Rödl, V., Szemerédi, E.: On size Ramsey numbers of graphs with bounded degree, *Combinatorica* **20** (2000), 257–262.
- [32] Suk, A.: On the order types of systems of segments in the plane, *Order* **27** (2010), 63–68.