A structure theorem for pseudo-segments and its applications

Jacob Fox \square

Department of Mathematics, Stanford University, Stanford, CA, USA

János Pach ⊠

HUN-REN Alfréd Rényi Institute of Mathematics, Budapest, Hungary

And rew Suk \square

Department of Mathematics, University of California San Diego, La Jolla, CA, USA

– Abstract -

- We prove a far-reaching strengthening of Szemerédi's regularity lemma for intersection graphs of 2
- pseudo-segments. It shows that the vertex set of such graphs can be partitioned into a bounded
- number of parts of roughly the same size such that almost all of the bipartite graphs between pairs
- of parts are complete or empty. We use this to get an improved bound on disjoint edges in simple
- topological graphs, showing that every n-vertex simple topological graph with no k pairwise disjoint
- edges has at most $n(\log n)^{O(\log k)}$ edges.

2012 ACM Subject Classification Mathematics of computing \rightarrow Combinatorics

Keywords and phrases Regularity lemma, pseudo-segments, intersection graphs

Related Version : \@@@par

Funding Jacob Fox: supported by a Packard Fellowship and by NSF Awards DMS-1800053 and DMS-2154169.

János Pach: Rényi Institute, Budapest. Supported by NKFIH grant K-176529, Austrian Science Fund Z 342-N31 and ERC Advanced Grant "GeoScape."

Andrew Suk: Supported by NSF CAREER award DMS-1800746, and NSF awards DMS-1952786 and DMS-2246847.

1 Introduction

Given a set of curves \mathcal{C} in the plane, we say that \mathcal{C} is a collection of *pseudo-sequents* if any 9 two members in $\mathcal C$ have at most one point in common, and no three members in $\mathcal C$ have a 10 point in common. The intersection graph of a collection \mathcal{C} of sets has vertex set \mathcal{C} and two 11 sets in C are adjacent if and only if they a have nonempty intersection. 12

A partition of a set is an *equipartition* if each pair of parts in the partition differ in size 13 by at most one. Szemerédi's celebrated regularity lemma roughly says that the vertex set 14 of any graph has an equipartition such that the bipartite graph between almost all pairs of 15 parts is random-like. Our main result is a strengthening of Szemerédi's regularity lemma for 16 intersection graphs of pseudo-segments. It replaces the condition that the bipartite graphs 17 between almost all pairs of parts is random-like to being complete or empty. 18

▶ **Theorem 1.** For each $\varepsilon > 0$ there is $K = K(\varepsilon)$ such that for every finite collection C of 19 pseudo-segments in the plane, there is an equipartition of C into K parts C_1, \ldots, C_K such 20 that for all but at most εK^2 pairs $\mathcal{C}_i, \mathcal{C}_j$ of parts, either every curve in \mathcal{C}_i crosses every curve 21 in C_i , or every curve in C_i is disjoint from every curve in C_i . 22

Pach and Solymosi [17] proved the special case of Theorem 1 where \mathcal{C} is a collection of 23 segments in the plane, and this result was later extended to semi-algebraic graphs [2] and 24 hypergraphs [5] of bounded description complexity. However, the techniques used to prove 25

© J. Fox. J. Pach. A. Suk: \odot





Editors: Wolfgang Mulzer and Jeff M. Phillips; Article No. XX; pp. XX:1–XX:14

Leibniz International Proceedings in Informatics

LIPICS Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

XX:2 A structure theorem for pseudo-segments and its applications

these results heavily rely on the algebraic structure. In fact, while it follows from the Milnor-Thom theorem that there are only $2^{O(n \log n)}$ graphs on n vertices which are semialgebraic

- of bounded description complexity (see [18, 2, 20]) there are many more (namely $2^{\Omega(n^{4/3})}$)
- $_{29}$ graphs on *n* vertices which are intersection graphs of pseudo-segments [6].
- Next, we discuss an application of Theorem 1 in graph drawing.

Disjoint edges in simple topological graphs. A topological graph is a graph drawn 31 in the plane such that its vertices are represented by points and its edges are represented 32 by nonself-intersecting arcs connecting the corresponding points. The edges are allowed to 33 intersect, but they may not intersect vertices apart from their endpoints. Furthermore, no 34 two edges are tangent, i.e., if two edges share an interior point, then they must properly cross 35 at that point in common. A topological graph is *simple* if every pair of its edges intersect at 36 most once. Two edges of a topological graph cross if their interiors share a point, and are 37 disjoint if they neither share a common vertex nor cross. 38

³⁹ Determining the maximum number of edges in a simple topological graph with no k⁴⁰ pairwise disjoint edges seems to be a difficult task. When k = 2, a linear upper bound is ⁴¹ known [16, 3, 10, 11]. When $k \ge 3$, Pach and Tóth [19] showed that every *n*-vertex simple ⁴² topological graph with no k pairwise disjoint edges has at most $O(n \log^{4k-8} n)$ edges. They ⁴³ conjectured that for every fixed k, the number of edges in such graphs is at most $O_k(n)$. Our ⁴⁴ next result substantially improves the upper bound for large k.

⁴⁵ ► **Theorem 2.** If G = (V, E) is an n-vertex simple topological graph with no k pairwise ⁴⁶ disjoint edges, then $|E(G)| \le n(\log n)^{O(\log k)}$.

⁴⁷ The proof of Theorem 2 follows the arguments in [19, 21], and is by double induction on ⁴⁸ *n* and *k*. We consider the cases when there are many or few disjoint pairs of edges in *G*. In ⁴⁹ the former case, we apply a variant of Theorem 1 and induction on *k*. In the latter case, we ⁵⁰ apply a bisection width result due to Pach and Tóth [19] and induction on *n*. See [7] for more ⁵¹ details. In [9], Fox and Sudakov showed that every dense *n*-vertex simple topological graph ⁵² contains $\Omega(\log^{1+\delta} n)$ pairwise disjoint edges, where $\delta \approx 1/40$. As an immediate Corollary to ⁵³ Theorem 2, we improve this bound to nearly polynomial under a much weaker assumption.

Corollary 3. Let $\varepsilon > 0$, and let G = (V, E) be an n-vertex simple topological graph with at least $2n^{1+\varepsilon}$ edges. Then G has $n^{\Omega(\varepsilon/\log \log n)}$ pairwise disjoint edges.

⁵⁶ For complete *n*-vertex simple topological graphs, Aichholzer et al. [1] showed that one can ⁵⁷ always find $\Omega(n^{1/2})$ pairwise disjoint edges.

The proofs of the above theorems heavily rely on the following bipartite Ramsey-type result for intersection graphs of pseudo-segments. As shown in [7], the main result in this paper, Theorem 1, is equivalent to the following.

⁶¹ ► **Theorem 4.** Let \mathcal{R} be a set of n red curves, and \mathcal{B} be a set of n blue curves in the plane ⁶² such that $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-segments. Then there are subsets $\mathcal{R}' \subset \mathcal{R}$ and ⁶³ $\mathcal{B}' \subset \mathcal{B}$, where $|\mathcal{R}'|, |\mathcal{B}'| \ge \Omega(n)$, such that either every curve in \mathcal{R}' crosses every curves in ⁶⁴ \mathcal{B}' , or every curve in \mathcal{R}' is disjoint from every curve in \mathcal{B}' .

The rest of this paper is devoted to proving Theorem 4. In the next section, we recall that any finite collection of pseudo-segments in the plane contains a linear-sized subset with the property that only a small fraction of pairs in the subset are crossing, or nearly all of them cross. In Section 3, we prove Theorem 4 in the special case where one of the families is double grounded. Building on these results, in Section 4, we establish our bipartite Ramsey-type theorem (Theorem 4) for any two families of pseudo-segments with the property that for
each family, only a small fraction of pairs are crossing, or nearly all of them cross. Finally, in
Section 5, we prove Theorem 4 in its full generality.

73 2 Tools

⁷⁴ We say that a graph G is ε -homogeneous if the edge density in G is less than ε or greater ⁷⁵ than $1 - \varepsilon$. For the proof of Theorem 4, we need the following result from [4].

Theorem 5 ([4]). There is an constant c' > 0 such that the following holds. Let C be a collection of n pseudo-segments in the plane with at least εn^2 crossing pairs. Then there are subsets $C_1, C_2 \subset C$, each of size $c'\varepsilon n$, such that every curve in C_1 crosses every curve in C_2 .

⁷⁹ Given a collection C of curves in the plane, let G(C) denote the intersection graph of ⁸⁰ C. In [8], Fox, Pach, and Tóth showed that pseudo-segments has the strong Erdős-Hajnal ⁸¹ property, which implies the following.

Corollary 6 ([8]). The family of intersection graphs of pseudo-segments has the polynomial Rödl property. That is, there is an absolute constant $c_1 > 0$ such that the following holds. Let $\varepsilon > 0$ and C be a collection of n pseudo-segments in the plane. Then there is a subset $C' \subset C$ of size $\varepsilon^{c_1} n$ whose intersection graph G(C') is ε -homogeneous.

We will frequently use the following simple lemma in this paper. See [7] for the proof.

▶ Lemma 7. Let G = (V, E) be a graph on n vertices. If the edge density of G is at most ε , then any induced subgraph on δn vertices has edge density at most $2\varepsilon/\delta^2$. Likewise, if the edge density of G is at least $1 - \varepsilon$, then any induced subgraph on δn vertices has edge density at least $1 - 2\varepsilon/\delta^2$.

⁹¹ **3** Proof of Theorem 4 – for double grounded red curves

Given a collection of curves C in the plane, we say that C is *double grounded* if there are two distinct curves γ_1 and γ_2 such that for each curve $\alpha \in C$, α has one endpoint on γ_1 and the other on γ_2 , and the interior of α is disjoint from γ_1 and γ_2 . Throughout this paper, for simplicity, we will always assume that both endpoints of each of our curves have distinct *x*-coordinates. We refer to the endpoint of a curve with the smaller (larger) *x*-coordinate as its *left (right) endpoint*. The aim of this section is to prove Theorem 4 in the special case where one of the color classes (the red one, say) consists of double grounded curves.

A curve in the plane is called *x*-monotone if every vertical line intersects it in at most one point. We start by considering double grounded *x*-monotone curves, and at the end of this section, we will remove the *x*-monotone condition. We will need the following result, known as the cutting-lemma for *x*-monotone curves. See, for example, Proposition 2.11 in [14].

▶ Lemma 8 (The Cutting Lemma). Let C be a collection of n double grounded x-monotone curves, whose grounds are disjoint vertical segments γ_1 and γ_2 , and let r > 1 be a parameter. Then $\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2)$ can be subdivided into t connected regions $\Delta_1, \ldots, \Delta_t$, such that the interior of each Δ_i is intersected by at most n/r curves from C, and we have $t = O(r^2)$.

¹⁰⁷ Throughout the paper, we will implicitly use the Jordan curve theorem.

Lemma 9. Let \mathcal{R} be a set of n red double grounded x-monotone curves, whose grounds are disjoint vertical segments γ_1 and γ_2 . Let \mathcal{B} be a set of n blue curves (not necessarily

XX:4 A structure theorem for pseudo-segments and its applications



Figure 1 Case 1, α_0 and α are disjoint.

¹¹⁰ x-monotone) such that every blue curve in \mathcal{B} is disjoint from grounds γ_1 and γ_2 , and suppose ¹¹¹ that $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-segments. Then there are subsets $\mathcal{R}' \subset \mathcal{R}$ and $\mathcal{B}' \subset \mathcal{B}$ ¹¹² such that $|\mathcal{R}'|, |\mathcal{B}'| \geq \Omega(n)$, and either every curve in \mathcal{R}' crosses every curve in \mathcal{B}' , or every ¹¹³ curve in \mathcal{R}' is disjoint from every curve in \mathcal{B}' .

Proof. Let P be the set of left-endpoints of the curves in \mathcal{B} . We apply Lemma 8 to \mathcal{R} with parameter r = 4 to obtain a subdivision $\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2) = \Delta_1 \cup \cdots \cup \Delta_t$, such that for each Δ_i , the interior of Δ_i intersects at most n/4 members in \mathcal{R} , and $t \leq c_0 4^2$ where c_0 is an absolute constant from Lemma 8. By the pigeonhole principle, there is a region Δ_i such that Δ_i contains at least $n/c_0 4^2$ points from P. Let $\mathcal{B}_0 \subset \mathcal{B}$ be the set of blue curves whose left endpoints are in Δ_i . Hence $|\mathcal{B}_0| = \Omega(n)$.

Let Q be the right endpoints of the curves in \mathcal{B}_0 . Using the same subdivision described above, there is a region Δ_j such that Δ_j contains at least $|Q|/(c_0 4^2) \ge n/(c_0 4^2)^2$ points from Q. Let $\mathcal{B}_1 \subset \mathcal{B}_0$ be the set of blue curves with their left endpoint in Δ_i and right endpoint in Δ_j . Let $\mathcal{R}_1 \subset \mathcal{R}$ consists of all red curves that do not intersect the interior of Δ_i and Δ_j . Lemma 8 implies that $|\mathcal{R}_1| \ge n - \frac{2n}{4} = \frac{n}{2}$, and $|\mathcal{B}_1| = \Omega(n)$. Recall that each blue curve in \mathcal{B}_1 does not intersect the grounds γ_1 nor γ_2 . Fix an arbitrary curve $\alpha_0 \in \mathcal{R}_1$. The proof now falls into the following cases.

¹²⁷ Case 1. Suppose at least $|\mathcal{R}_1|/2$ curves in \mathcal{R}_1 are disjoint from α_0 . Let $\mathcal{R}_2 \subset \mathcal{R}_1$ be the set ¹²⁸ of red curves disjoint from α_0 . For each $\alpha \in \mathcal{R}_2$, $\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2 \cup \alpha_0 \cup \alpha)$, consists of two ¹²⁹ connected components, one bounded and the other unbounded.

¹³³ Case 1.a. Suppose for at least $|\mathcal{R}_2|/2$ red curves $\alpha \in \mathcal{R}_2$, both Δ_i and Δ_j lie in the same ¹³⁴ connected component of $\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2 \cup \alpha_0 \cup \alpha)$. See Figure 1a. Let $\mathcal{R}_3 \subset \mathcal{R}_2$ be the collection ¹³⁵ of such red curves. Then for each $\alpha \in \mathcal{R}_3$, each blue curve $\beta \in \mathcal{B}_1$ crosses α if and only if ¹³⁶ β crosses α_0 . Hence, there is a subset $\mathcal{B}_2 \subset \mathcal{B}_1$ of size at least $\Omega(n)$, such that either every ¹³⁷ blue curve in \mathcal{B}_2 crosses every red curve in \mathcal{R}_3 , or every blue curve in \mathcal{B}_2 is disjoint from ¹³⁸ every red curve in \mathcal{R}_3 . Moreover, $|\mathcal{R}_3| = \Omega(n)$ and we are done.

¹³⁹ Case 1.b. Suppose for at least $|\mathcal{R}_2|/2$ red curves $\alpha \in \mathcal{R}_2$, regions Δ_i and Δ_j lie in different ¹⁴⁰ connected component of $\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2 \cup \alpha_0 \cup \alpha)$. See Figure 1b. Similar to above, let $\mathcal{R}_3 \subset \mathcal{R}_2$ ¹⁴¹ be the collection of such red curves. By the pseudo-segment condition, for each $\alpha \in \mathcal{R}_3$, each ¹⁴² blue curve $\beta \in \mathcal{B}_1$ crosses α if and only if β is disjoint from α_0 . Hence, there is a subset ¹⁴³ $\mathcal{B}_2 \subset \mathcal{B}_1$ of size $\Omega_r(n)$, such that either every blue curve in \mathcal{B}_2 crosses every red curve in \mathcal{R}_3 , ¹⁴⁴ or every blue curve in \mathcal{B}_2 is disjoint from every red curve in \mathcal{R}_3 . Moreover, $|\mathcal{R}_3| = \Omega(n)$ and ¹⁴⁵ we are done.

¹⁴⁶ Case 2. Suppose at least $|\mathcal{R}_1|/2$ curves in \mathcal{R}_1 cross α_0 . Let $\mathcal{R}_2 \subset \mathcal{R}_1$ be the set of red curves ¹⁴⁷ that crosses α_0 . For each $\alpha \in \mathcal{R}_2 \setminus \{\alpha_0\}, \mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2 \cup \alpha_0 \cup \alpha)$ consists of three connected ¹⁴⁸ components, two of which are bounded and the other unbounded.

J. Fox, J. Pach, A. Suk



Figure 2 Case 2, α_0 and α cross.

¹⁴⁹ Case 2.a. Suppose for at least $|\mathcal{R}_2|/3$ red curves $\alpha \in \mathcal{R}_2$, Both Δ_i and Δ_j lie in the same ¹⁵⁰ connected component of $\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2 \cup \alpha_0 \cup \alpha)$. See Figure 2a. Let $\mathcal{R}_3 \subset \mathcal{R}_2$ be the collection ¹⁵¹ of such red curves. By the pseudo-segment condition, for each $\alpha \in \mathcal{R}_3$, each blue curve ¹⁵² $\beta \in \mathcal{B}_1$ crosses α if and only if β crosses α_0 . Hence, there is a subset $\mathcal{B}_2 \subset \mathcal{B}_1$ of size at least ¹⁵³ $\Omega(n)$, such that either every blue curve in \mathcal{B}_2 crosses every red curve in \mathcal{R}_3 , or every blue ¹⁵⁴ curve in \mathcal{B}_2 is disjoint from every red curve in \mathcal{R}_3 . Moreover, $|\mathcal{R}_3| = \Omega(n)$.

¹⁵⁵ Case 2.b. Suppose for at least $|\mathcal{R}_2|/3$ red curves $\alpha \in \mathcal{R}_2$, regions Δ_i and Δ_j lie in different ¹⁵⁶ bounded connected components of $\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2 \cup \alpha_0 \cup \alpha)$. Let $\mathcal{R}_3 \subset \mathcal{R}_2$ be the collection of ¹⁵⁷ such red curves. Then for each $\alpha \in \mathcal{R}_3$, every blue curve $\beta \in \mathcal{B}_1$ crosses α . Since $|\mathcal{R}_3| = \Omega(n)$, ¹⁵⁸ we have $|\mathcal{B}_1| = \Omega(n)$.

¹⁵⁹ Case 2.c. Suppose for at least $|\mathcal{R}_2|/3$ red curves $\alpha \in \mathcal{R}_2$, regions Δ_i and Δ_j lie in different ¹⁶⁰ connected components of $\mathbb{R}^2 \setminus (\gamma_1 \cup \gamma_2 \cup \alpha_0 \cup \alpha)$, one of which is bounded and the other ¹⁶¹ unbounded. See Figure 2b. Let $\mathcal{R}_3 \subset \mathcal{R}_2$ be the collection of such red curves. By the ¹⁶² pseudo-segment condition, for each $\alpha \in \mathcal{R}_3$, each blue curve $\beta \in \mathcal{B}_1$ crosses α if and only if ¹⁶³ β is disjoint from α_0 . Hence, there is a subset $\mathcal{B}_2 \subset \mathcal{B}_1$ of size $\Omega(n)$, such that either every ¹⁶⁴ blue curve in \mathcal{B}_2 crosses every red curve in \mathcal{R}_3 , or every blue curve in \mathcal{B}_2 is disjoint from ¹⁶⁵ every red curve in \mathcal{R}_3 . Moreover, $|\mathcal{R}_3| = \Omega(n)$, and we are done.

Recall that a *pseudoline* is an unbounded arc in \mathbb{R}^2 , whose complement is disconnected. An *arrangement of pseudolines* is a set of pseudolines such that every pair meets exactly once, and no three members have a point in common. A classic result of Goodman [12] states that every arrangement of pseudolines is isomorphic to an arrangement of *wiring diagram* (bi-infinite *x*-monotone curves). Moreover, Goodman and Pollack showed the following.

Theorem 10 ([13]). Every arrangement of pseudolines can be continuously deformed
 (through isomorphic arrangements) to a wiring diagram.

¹⁷⁶ We also need the following simple lemma.

Lemma 11. Given a finite linearly ordered set whose elements are colored red or blue, we
 can select half of the red elements and half of the blue elements such that all of the selected
 elements of one color come before all of the selected elements of the other color.

180 We are now ready to establish the main result of this section:

Theorem 12. Let \mathcal{R} be a set of n red double grounded curves with grounds γ_1 and γ_2 , where γ_1 and γ_2 cross each other. Let \mathcal{B} be a set of n blue curves such that $\mathcal{R} \cup \mathcal{B} \cup \{\gamma_1, \gamma_2\}$ is a collection of pseudo-segments. Then there are subsets $\mathcal{R}' \subset \mathcal{R}$ and $\mathcal{B}' \subset \mathcal{B}$ such that $|\mathcal{R}'|, |\mathcal{B}'| \ge \Omega(n)$, and either every curve in \mathcal{R}' crosses every curve in \mathcal{B}' , or every curve in \mathcal{R}' is disjoint from every curve in \mathcal{B}' .

XX:6 A structure theorem for pseudo-segments and its applications

Proof. By passing to linear-sized subsets of \mathcal{R} and \mathcal{B} and subcurves of γ_1 and γ_2 , we will reduce the problem to the setting of Lemma 9. Let us assume that γ_1 and γ_2 cross at point p. Hence, $(\gamma_1 \setminus \gamma_2) \cup (\gamma_2 \setminus \gamma_1)$ consists of four connected components. By the pigeonhole principle, there is a subset $\mathcal{R}_1 \subset \mathcal{R}$ of size n/4 such that every curve in \mathcal{R}_1 has an endpoint on one of the connected components of $\gamma_1 \setminus \gamma_2$, and all of the other endpoints lie on one of the connected components of $\gamma_2 \setminus \gamma_1$. Let $\gamma'_i \subset \gamma_i$, for i = 1, 2, be these connected components so that they have a common endpoint at p and their interiors are disjoint.

For each $\alpha \in \mathcal{R}_1$, the sequence of curves $(\gamma'_1, \gamma'_2, \alpha)$ appear either in *clockwise* or *counterclockwise* order along the unique simple closed curve that lies in $\gamma'_1 \cup \gamma'_2 \cup \alpha$. Without loss of generality, we can assume that there is a subset $\mathcal{R}_2 \subset \mathcal{R}_1$, where $|\mathcal{R}_2| = \Omega(n)$, such that for every curve $\alpha \in \mathcal{R}_2$, the sequence $(\gamma'_1, \gamma'_2, \alpha)$ appears in clockwise order, since a symmetric argument would follow otherwise.

We define the *orientation* of each curve $\alpha \in \mathcal{R}_2$ as the sequence of turns, either *left*-198 *left, left-right, right-left,* or *right-right,* made by starting at p and moving along γ'_1 in the 199 arrangement $\gamma'_1 \cup \gamma'_2 \cup \alpha$, until we return back to p. More precisely, starting at p we move 200 along γ'_1 until we reach the endpoint of α . We then turn either left or right to move along α 201 towards γ'_2 . Once we've reached γ'_2 , we either turn left or right in order to move along γ'_2 202 and reach p again. By the pigeonhole principle, there is a subset $\mathcal{R}_3 \subset \mathcal{R}_2$ of size at least 203 $\Omega(n)$ such that all curves in \mathcal{R}_3 have the same orientation. Without loss of generality, we can 204 assume that the orientation is *left-left*, since a symmetric argument would follow otherwise. 205

Starting at p and moving along γ'_1 towards its other endpoint, let us consider the sequence 206 of curves from $\mathcal{R}_3 \cup \mathcal{B}$ intersecting γ'_1 . Then, by Lemma 11, there are subsets $\mathcal{R}_4 \subset \mathcal{R}_3$ 207 and $\mathcal{B}_1 \subset \mathcal{B}$, where $|\mathcal{R}_4| \geq |\mathcal{R}_3|/2$ and $|\mathcal{B}_1| \geq |\mathcal{B}|/2$, such that either all of the curves in \mathcal{R}_4 208 appear before all of the curves in \mathcal{B}_1 that intersect γ'_1 in this sequence, or all of the curves in 209 \mathcal{R}_4 appear after all of the curves in \mathcal{B}_1 in this sequence. Note that \mathcal{B}_1 consists of the blue 210 curves in \mathcal{B} that are disjoint to γ'_1 and at least half of the curves in \mathcal{B} that intersect γ'_1 found 211 by the application of Lemma 11. Hence, there is a subcurve $\gamma_1'' \subset \gamma_1'$ such that γ_1'' is one of 212 the grounds for \mathcal{R}_4 , and is disjoint from every curve in \mathcal{B}_1 . We apply the same argument to 213 $\mathcal{R}_4 \cup \mathcal{B}_1$ and γ'_2 , and obtain subsets $\mathcal{R}_5 \subset \mathcal{R}_4$, $\mathcal{B}_2 \subset \mathcal{B}_1$, and a subcurve $\gamma''_2 \subset \gamma'_2$, such that 214 $|\mathcal{R}_5|, |\mathcal{B}_2| = \Omega(n)$, and \mathcal{R}_5 is double grounded with disjoint grounds γ_1'' and γ_2'' , and every 215 curve in \mathcal{B}_2 is disjoint from γ_1'' and γ_2'' . 216

For $i \in \{1, 2\}$, let p_i be the endpoint of γ''_i that lies closest to p along γ'_i . Starting at p_i 217 and moving along γ''_i , let π_i be the sequence of curves in \mathcal{R}_5 that appear on γ''_i . Since every 218 curve in \mathcal{R}_5 has the same left-left orientation, and appears clockwise order with respect to 219 γ'_1 and γ'_2 , two curves $\alpha, \alpha' \in \mathcal{R}_5$ cross if and only if the order in which they appear in π_1 220 and π_2 changes. Let γ_3'' be a curve very close to γ_2'' such that γ_3'' has the same endpoints as 221 γ_2'' , and is disjoint from all curves in $\mathcal{R}_5 \cup \mathcal{B}_2$. Hence, $\gamma_2'' \cup \gamma_3''$ makes an empty lens in the 222 arrangement $\mathcal{R}_5 \cup \mathcal{B}_2$. We slightly extend each curve $\alpha \in \mathcal{R}_5$ through this lens to γ''_3 so that 223 the resulting curve, α' properly crosses γ''_2 and has its new endpoint on γ''_3 . Moreover, the 224 extension will be made in such a way that the sequence π_3 of curves in \mathcal{R}_5 appearing along 225 γ_3'' starting from p_2 will appear in the opposite order of π_1 . Let $\mathcal{R}'_5 = \{\alpha' : \alpha \in \mathcal{R}_5\}$. Thus, 226 every pair of curves in \mathcal{R}'_5 will cross exactly once. 227

For each curve $\alpha' \in \mathcal{R}'_5$, we further extend α' by moving both endpoints towards p along γ_1 and γ_2 , so that we do not create any additional crossings within \mathcal{R}'_5 . Let $\hat{\alpha}$ be the resulting extension, where both endpoints of $\hat{\alpha}$ lie arbitrarily close to p. Set $\hat{\mathcal{R}}_5 = \{\hat{\alpha} : \alpha' \in \mathcal{R}'_5\}$. See Figure 3. Furthermore, we can assume that p lies in the unbounded face of the arrangement $\hat{\mathcal{R}}_5$, since otherwise we could project the arrangement $\hat{\mathcal{R}}_5$ onto a sphere, and then project it back to the plane so that p lies in the unbounded face, without creating or removing any



Figure 3 The resulting extension $\hat{\mathcal{R}}_5$.

crossing. Therefore, $\hat{\mathcal{R}}_5$ can be extended to a family of pseudolines. By Theorem 10, we can 234 apply a continuous deformation of the plane so that $\hat{\mathcal{R}}_5$ becomes a collection of unbounded 235 x-monotone curves. Hence, after the deformation, the original set \mathcal{R}_5 becomes a collection 236 of double grounded x-monotone curves, with grounds γ_1'', γ_2'' , such that every curve in \mathcal{B}_2 is 237 disjoint from the grounds γ_1'' and γ_2'' , the crossing pattern in the arrangement $\mathcal{R}_5 \cup \mathcal{B}_2$ is the 238 same as before. Moreover, γ_1'' and γ_2'' will be disjoint vertical segments. We apply Lemma 9 239 to \mathcal{R}_5 and \mathcal{B}_2 and obtain subsets $\mathcal{R}_6 \subset \mathcal{R}_5$ and $\mathcal{B}_3 \subset \mathcal{B}_2$, each of size $\Omega(n)$, such that either 240 every curve in \mathcal{R}_6 crosses every curve in \mathcal{B}_3 , or every curve in \mathcal{R}_6 is disjoint from every curve 241 in \mathcal{B}_3 . This completes the proof. 242

By combining Theorem 12 with a variant of Szemerédi's regularity lemma due to Kómlos [15], we have the following (see [7] for more details).

▶ Theorem 13. There is a constant c' > 0 such that the following holds. Let \mathcal{R} be a collection of n red double grounded curves with grounds γ_1 and γ_2 , such that γ_1 and γ_2 cross. Let \mathcal{B} be a collection of n blue curves such that $\mathcal{R} \cup \mathcal{B} \cup \{\gamma_1, \gamma_2\}$ is a collection of pseudo-segments. If there are at least εn^2 crossing pairs in $\mathcal{R} \times \mathcal{B}$, then there are subsets $\mathcal{R}' \subset \mathcal{R}, \mathcal{B}' \subset \mathcal{B}$, where $|\mathcal{R}'|, |\mathcal{B}'| \ge \varepsilon^{c'} n$, such that every curve in \mathcal{R}' crosses every curve in \mathcal{B}' .

▶ **Theorem 14.** There is an absolute constant c' > 0 such that the following holds. Let \mathcal{R} be a collection of n red double grounded curves with grounds γ_1 and γ_2 , such that γ_1 and γ_2 cross. Let \mathcal{B} be a collection of n blue curves such that $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-segments. If there are at least εn^2 disjoint pairs in $\mathcal{R} \times \mathcal{B}$, then there are subsets $\mathcal{R}' \subset \mathcal{R}, \mathcal{B}' \subset \mathcal{B}$, where $|\mathcal{R}'|, |\mathcal{B}'| \ge \varepsilon^{c'} n$, such that every curve in \mathcal{R}' is disjoint from every curve in \mathcal{B}' .

²⁵⁶ We will apply Theorems 12 and 13 in the next section.

4 Proof of Theorem 4 – for ε -homogeneous families

The aim of this section is to prove Theorem 4, the main result of this paper, in the special case where the edge density of the intersection graph of the red curves is nearly 0 or nearly 1, and the same is true for the intersection graph of the blue curves. This will easily imply Theorem 4 in its full generality, as shown in the next section.

²⁶² 4.1 Low versus low density

By Corollary 6, we can reduce to the case that the intersection graphs $G(\mathcal{R})$ and $G(\mathcal{B})$ are both ε -homogeneous. Below, we first consider the cases when both $G(\mathcal{R}), G(\mathcal{B})$ has edge density less than ε .

XX:8 A structure theorem for pseudo-segments and its applications



Figure 4 Partitioning of the red curve $\alpha = \alpha_u \cup \alpha_\ell$.

▶ **Theorem 15.** There is an absolute constant $\varepsilon_1 > 0$ such that the following holds. Let \mathcal{R} be a set of n red curves and \mathcal{B} be a set of n blue curves in the plane such that $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-segments. If the edge densities of the intersection graphs $G(\mathcal{R})$ and $G(\mathcal{B})$ are both less than ε_1 , then there are subsets $\mathcal{R}' \subset \mathcal{R}$ and $\mathcal{B}' \subset \mathcal{B}$, each of size $\Omega(n)$, such that every red curve in \mathcal{R}' crosses every blue curve in \mathcal{B}' , or every red curve in \mathcal{R}' is disjoint from every blue curve in \mathcal{B}' .

The proof of Theorem 15 is a simple application of a separator theorem from [4] (see [7]).

²⁷³ 4.2 High versus low edge density

In this subsection, we consider the case when the intersection graph $G(\mathcal{R})$ has edge density at least $1 - \varepsilon$, and $G(\mathcal{B})$ has edge density less than ε . Since the edge density in the intersection graph $G(\mathcal{R})$ is at least $1 - \varepsilon$, we can further reduce to the case when there is a red curve γ_1 that crosses every member in \mathcal{R} exactly once.

▶ Lemma 16. For each integer $t \ge 1$, there is a constant $\varepsilon'_t > 0$ such that the following holds. 278 Let \mathcal{R} be a set of n red curves in the plane, all crossed by a curve γ_1 exactly once, and \mathcal{B} be 279 a set of n blue curves in the plane such that $\mathcal{R} \cup \mathcal{B} \cup \{\gamma_1\}$ is a collection of pseudo-segments. 280 Suppose that the intersection graph $G(\mathcal{B})$ has edge density less than ε'_t , and $G(\mathcal{R})$ has edge 281 density at least $1 - \varepsilon'_t$. Then there are subsets $\hat{\mathcal{R}} \subset \mathcal{R}$, $\hat{\mathcal{B}} \subset \mathcal{B}$, each of size $\Omega_{\varepsilon'_t}(n)$, such that 282 either every red curve in $\hat{\mathcal{R}}$ crosses every blue curve in $\hat{\mathcal{B}}$, or every red curve in $\hat{\mathcal{R}}$ is disjoint 283 from every blue curve in $\hat{\mathcal{B}}$, or each curve $\alpha \in \hat{\mathcal{R}}$ has a partition into two connected parts 284 $\alpha = \hat{\alpha}_u \cup \hat{\alpha}_\ell$, such that for 285

$$\hat{\mathcal{U}} = \{ \hat{\alpha}_u : \alpha \in \hat{\mathcal{R}}, \alpha = \hat{\alpha}_u \cup \hat{\alpha}_\ell \} \quad \text{and} \quad \hat{\mathcal{L}} = \{ \hat{\alpha}_\ell : \alpha \in \hat{\mathcal{R}}, \alpha = \hat{\alpha}_u \cup \hat{\alpha}_\ell \},$$

every curve in $\hat{\mathcal{L}}$ is disjoint to every curve in $\hat{\mathcal{B}}$, and the edge density of $G(\hat{\mathcal{U}})$ is less than 2^{-t} .

Proof. Each curve $\alpha \in \mathcal{R}$ is partitioned into two connected parts by γ_1 , say an upper and lower part. More precisely, we have the partition $\alpha = \alpha_u \cup \alpha_\ell$, where the parts α_u and α_ℓ are defined, as follows. We start at the left endpoint of γ_1 and move along γ_1 until we reach $\alpha \cap \gamma_1$. At this point, we turn left along α to obtain α_u and right to obtain α_ℓ . See Figure 4. Let $\mathcal{U}(\mathcal{L})$ be the upper (lower) part of each curve in \mathcal{R} , that is,

²⁹³
$$\mathcal{U} = \{ \alpha_u : \alpha \in \mathcal{R}, \alpha = \alpha_\ell \cup \alpha_u \}$$
 and $\mathcal{L} = \{ \alpha_\ell : \alpha \in \mathcal{R}, \alpha = \alpha_\ell \cup \alpha_u \}.$

In what follows, for every integer $t \geq 1$, we will obtain subsets $\mathcal{R}^{(t)} \subset \mathcal{R}$, $\mathcal{B}^{(t)} \subset \mathcal{B}$, each of size $\Omega_{\varepsilon'_t}(n)$, such that either every red curve in $\mathcal{R}^{(t)}$ crosses every blue curve in $\mathcal{B}^{(t)}$, or every red curve in $\mathcal{R}^{(t)}$ is disjoint from every blue curve in $\mathcal{B}^{(t)}$, or each curve $\alpha \in \mathcal{R}^{(t)}$ has a new partition into upper and lower parts $\alpha = \alpha'_u \cup \alpha'_\ell$, such that the following holds.

J. Fox, J. Pach, A. Suk

- 1. We have $\alpha'_u \subset \alpha_u$, that is, the upper part α'_u is a subcurve of the previous upper part α_u .
- 2. The lower part α'_{ℓ} of each curve in $\mathcal{R}^{(t)}$ is disjoint from each blue curve in $\mathcal{B}^{(t)}$.
- 301 3. There is an equipartition $\mathcal{R}^{(t)} = \mathcal{R}_1^{(t)} \cup \cdots \cup \mathcal{R}_{2^t}^{(t)}$ into 2^t parts such that for $1 \le i < j \le 2^{t-1}$, 302 the upper part α'_u of each curve $\alpha \in \mathcal{R}_i^{(t)}$ is disjoint from the upper part β'_u of each curve 303 $\beta \in \mathcal{R}_i^{(t)}$.

Hence, the lemma follows from the statement above by setting $\hat{\mathcal{B}} = \mathcal{B}^{(t)}, \hat{\mathcal{R}} = \mathcal{R}^{(t)}$.

We proceed by induction on t. The bulk of the argument below is actually for the base case t = 1, since we will just repeat the entire argument for the inductive step with parameter ε'_t . Let ε'_1 be a small positive constant that will be determined later such that $\varepsilon'_1 < \varepsilon_1$, where ε_1 is from Theorem 15. Thus, $G(\mathcal{R})$ has edge density at least $1 - \varepsilon'_1$ and $G(\mathcal{B})$ has edge density less than ε'_1 .

Let $\delta > 0$ also be a sufficiently small constant determined later, such that $\varepsilon'_1 < \delta < \varepsilon_1$. We apply Corollary 6 to \mathcal{L} with parameter δ and obtain a subset $\mathcal{L}_1 \subset \mathcal{L}$ such that \mathcal{L}_1 is δ -homogeneous and $|\mathcal{L}_1| = \Omega_{\delta}(n)$. Let $\mathcal{R}_1 \subset \mathcal{R}$ be the red curves in \mathcal{R} corresponding to the curves in \mathcal{L}_1 , and let $\mathcal{U}_1 \subset \mathcal{U}$ be the curves in \mathcal{U} that corresponds to the red curves in \mathcal{R}_1 .

Without loss of generality, we can assume that the intersection graph $G(\mathcal{L}_1)$ has edge 314 density less than δ . Indeed, otherwise if $G(\mathcal{L}_1)$ has edge density greater than $1-\delta$, by the 315 pseudo-segment condition, the intersection graph $G(\mathcal{U}_1)$ must have edge density less than δ 316 and a symmetric argument would follow. In order to apply Theorem 15, we need two subsets 317 of equal size. By averaging, there is a subset $\mathcal{B}' \subset \mathcal{B}$ with $|\mathcal{B}'| = |\mathcal{L}_1|$ such that the edge 318 density of $G(\mathcal{B}')$ is at most that of $G(\mathcal{B})$. Since $G(\mathcal{L}_1)$ has edge density less than δ and $G(\mathcal{B}')$ 319 has edge density less than ε'_1 , by setting $\varepsilon'_1 < \delta < \varepsilon_1$, we can apply Theorem 15 to \mathcal{L}_1 and 320 \mathcal{B}' and obtain subsets $\mathcal{L}_2 \subset \mathcal{L}_1$ and $\mathcal{B}_1 \subset \mathcal{B}'$, each of size $\Omega_{\delta}(n)$, such that every curve in 321 \mathcal{L}_2 crosses every blue curve in \mathcal{B}_1 , or every curve in \mathcal{L}_2 is disjoint from every blue curve in 322 \mathcal{B}_1 . If we are in the former case, then we are done. Hence, we can assume that we are in 323 the latter case. Let $\mathcal{R}_2 \subset \mathcal{R}_1$ be the red curves that corresponds to \mathcal{L}_2 , and let $\mathcal{U}_2 \subset \mathcal{U}_1$ be 324 the curves in \mathcal{U}_1 that corresponds to \mathcal{R}_2 . We apply Corollary 6 to \mathcal{U}_2 with parameter δ and 325 obtain a subset $\mathcal{U}_3 \subset \mathcal{U}_2$ such that \mathcal{U}_3 is δ -homogeneous and $|\mathcal{U}_3| = \Omega_{\delta}(n)$. Let \mathcal{R}_3 be the 326 red curves in \mathcal{R} corresponding to \mathcal{U}_3 , and let \mathcal{L}_3 be the curves in \mathcal{L}_2 that corresponds to \mathcal{R}_3 . 327 Suppose that the intersection graph $G(\mathcal{U}_3)$ has edge density less than δ . Since $|\mathcal{B}_1| = \delta_0 n$, 328 where $\delta_0 = \delta_0(\delta, \varepsilon_1)$, by Lemma 7, the intersection graph $G(\mathcal{B}_1)$ has edge density at most 329

where $\sigma_0 = \sigma_0(\sigma, \varepsilon_1)$, by hermitar, the intersection graph $\mathcal{C}(\mathcal{L}_1)$ has edge density at most 2 $\varepsilon_1'/\delta_0^2$. Thus, we set δ and ε_1' sufficiently small so that $\delta < \varepsilon_1$ and $2\varepsilon_1'/\delta_0^2 < \varepsilon_1$. By averaging, we can find subsets of \mathcal{U}_3 and \mathcal{B}_1 , each of size $\min(|\mathcal{U}_3|, |\mathcal{B}_1|)$ and with densities less than ε_1 , and apply Theorem 15 to these subsets and obtain subsets $\mathcal{U}_4 \subset \mathcal{U}_3$ and $\mathcal{B}_2 \subset \mathcal{B}_1$, each of size $\Omega_{\delta}(n)$, such that every curve in \mathcal{U}_4 crosses every blue curve in \mathcal{B}_2 , or every curve in \mathcal{U}_4 is disjoint from every blue curve in \mathcal{B}_2 . In both cases, we are done since every curve in \mathcal{L}_3 is disjoint from every curve in \mathcal{B}_2 . Therefore, we can assume that $G(\mathcal{U}_3)$ has edge density greater than $1 - \delta$.

For each curve $\alpha \in \mathcal{U}_3$, let $N(\alpha)$ denote the set of curves in \mathcal{U}_3 that intersects α , and let $d(\alpha) = |N(\alpha)|$. We label the curves $\beta \in N(\alpha)$ with integers 0 to $d(\alpha) - 1$ according to their closest intersection point to the ground γ_1 along α , that is, the label $f_{\alpha}(\beta)$ of $\beta \in N(\alpha)$ is the number of curves in \mathcal{U}_3 that intersects the portion of α strictly between γ_1 and $\alpha \cap \beta$. Since $\sum_{\alpha \in \mathcal{U}_3} d(\alpha) - 1 \ge 2(1 - \delta) {|\mathcal{U}_3| \choose 2} - |\mathcal{U}_3|$, by Jensen's inequality, we have

$$\sum_{\alpha \in \mathcal{U}_3} \sum_{\beta \in N(\alpha)} f_{\alpha}(\beta) = \sum_{\alpha \in \mathcal{U}_3} \binom{d(\alpha)}{2} \ge |\mathcal{U}_3| \binom{\frac{\sum_{\alpha \in \mathcal{U}_3} d(\alpha)}{|\mathcal{U}_3|}}{2} \ge \frac{|\mathcal{U}_3|^3}{4}.$$

XX:10 A structure theorem for pseudo-segments and its applications

Let the weight $w(\beta)$ of a curve $\beta \in \mathcal{U}_3$ be the sum of its labels, that is, $w(\beta) = \sum_{\alpha:\beta \in N(\alpha)} f_{\alpha}(\beta)$.

Hence, the weight $w(\beta)$ is the total number of crossing points along curves α strictly between γ_1 and β , where α crosses both γ_1 and β . By averaging, there is a curve $\gamma_2 \in \mathcal{U}_3$ whose weight is at least $|\mathcal{U}_3|^2/4$.

Using γ_2 , we partition each curve $\alpha \in \mathcal{U}_3 \setminus \{\gamma_2\}$ that crosses γ_2 into two connected parts, $\alpha = \alpha_w \cup \alpha_m$, where α_m is the connected subcurve with endpoints on γ_1 and γ_2 , and α_w is the other connected part. Set

$$\mathcal{W}_3 = \{ \alpha_w : \alpha \in \mathcal{U}_3 \setminus \{\gamma_2\}, \alpha \cap \gamma_2 \neq \emptyset \} \quad \text{and} \quad \mathcal{M}_3 = \{ \alpha_m : \alpha \in \mathcal{U}_3 \setminus \{\gamma_2\}, \alpha \cap \gamma_2 \neq \emptyset \}.$$

Since γ_2 has weight at least $|\mathcal{U}_3|^2/4$, by the pigeonhole principle, there are at least $|\mathcal{U}_3|^2/8$ intersecting pairs in $\mathcal{M}_3 \times \mathcal{M}_3$, or at least $|\mathcal{U}_3|^2/8$ intersecting pairs in $\mathcal{M}_3 \times \mathcal{W}_3$.

Case 1. Suppose there are at least $|\mathcal{U}_3|^2/8$ pairs in $\mathcal{M}_3 \times \mathcal{W}_3$ that cross. The set \mathcal{M}_3 is double 353 grounded with grounds γ_1 and γ_2 that cross exactly once, and every curve in \mathcal{W}_3 is disjoint 354 from γ_1 and γ_2 . As $|\mathcal{M}_3|, |\mathcal{W}_3| \leq |\mathcal{U}_3|$, the density of edges in the bipartite intersection graph 355 of \mathcal{M}_3 and \mathcal{W}_3 is at least 1/8. By averaging, we can find subsets of \mathcal{M}_3 and \mathcal{W}_3 each of size 356 $\min(|\mathcal{M}_3|, |\mathcal{W}_3|)$ such that the density of edges in the bipartite intersection graph of these 357 subsets is at least 1/8. By setting $\delta > 0$ sufficiently small, we can apply Theorem 13 to these 358 subsets of \mathcal{M}_3 and \mathcal{W}_3 and obtain subsets $\mathcal{M}_4 \subset \mathcal{M}_3$ and $\mathcal{W}'_4 \subset \mathcal{W}_3$, each of size $\Omega_{\delta}(n)$, such 359 that each curve in \mathcal{M}_4 crosses each curve in \mathcal{W}'_4 . Moreover, by the pseudo-segment condition, 360 each curve in $\mathcal{M}_4 \cup \mathcal{W}'_4$ corresponds to a unique curve in \mathcal{U}_3 . Let $\mathcal{U}_4 \subset \mathcal{U}_3$ be the curves that 361 corresponds to \mathcal{M}_4 and let $\mathcal{U}'_4 \subset \mathcal{U}_3$ be the curves that corresponds to \mathcal{W}'_4 . Hence, we set 362

363
$$\mathcal{W}_4 = \{\alpha_m : \alpha \in \mathcal{U}_4, \alpha = \alpha_w \cup \alpha_m\}$$
 and $\mathcal{M}'_4 = \{\alpha_m : \alpha \in \mathcal{U}'_4, \alpha = \alpha_w \cup \alpha_m\}$

See Figure 5a. We apply Theorem 12 to arbitrary subsets of \mathcal{M}_4 and \mathcal{B}_2 , each of size min $(|\mathcal{M}_4|, |\mathcal{B}_2|)$, and obtain subsets $\mathcal{M}_5 \subset \mathcal{M}_4$ and $\mathcal{B}_3 \subset \mathcal{B}_2$, each of size $\Omega_{\delta}(n)$, such that either every red curve in \mathcal{M}_5 crosses every blue curve in \mathcal{B}_3 , or every red curve in \mathcal{M}_5 is disjoint from every blue curve in \mathcal{B}_3 . In the former case, we are done. Hence, we can assume that we are in the latter case.

We again apply Theorem 12 to arbitrary subsets of \mathcal{M}'_4 and \mathcal{B}_3 , each of size $\min(|\mathcal{M}'_4|, |\mathcal{B}_3|)$, to obtain subsets $\mathcal{M}'_5 \subset \mathcal{M}'_4$ and $\mathcal{B}_4 \subset \mathcal{B}_3$, each of size $\Omega_{\delta}(n)$, such that either every red curve in \mathcal{M}'_5 crosses every blue curve in \mathcal{B}_4 , or every red curve in \mathcal{M}'_5 is disjoint from every blue curve in \mathcal{B}_4 . Again, if we are in the former case, we are done. Hence, we can assume that we are in the latter case. Let

$$\mathcal{W}_5 = \{ \alpha_w : \alpha = \alpha_w \cup \alpha_m, \alpha_m \in \mathcal{M}_5 \} \quad \text{and} \quad \mathcal{W}_5' = \{ \alpha_w : \alpha = \alpha_w \cup \alpha_m, \alpha_m \in \mathcal{M}_5' \},$$

and recall that every element in \mathcal{M}_5 crosses every element in \mathcal{W}'_5 . By the pseudo-segment condition, every element in \mathcal{W}_5 is disjoint from every element in \mathcal{W}'_5 .

Let \mathcal{R}_5 be the red curves in \mathcal{R} that corresponds to \mathcal{W}_5 , and let \mathcal{R}'_5 be the red curves in \mathcal{R} that corresponds to \mathcal{W}'_5 . We have $|\mathcal{R}_5|, |\mathcal{R}'_5| = \Omega_\delta(n)$, and moreover, we can assume that $|\mathcal{R}_5| = |\mathcal{R}'_5|$. For each curve $\alpha \in \mathcal{R}_5 \cup \mathcal{R}'_5$, and its original partition $\alpha = \alpha_u \cup \alpha_\ell$ defined by γ_1 , we have a new partition $\alpha = \alpha'_u \cup \alpha'_\ell$ defined by γ_2 , where $\alpha'_u = \alpha_w$ and $\alpha'_\ell = \alpha_m \cup \alpha_\ell$. By setting $\mathcal{R}^{(1)} = \mathcal{R}_5 \cup \mathcal{R}'_5$, and $\mathcal{B}^{(1)} = \mathcal{B}_4$, where each curve $\alpha \in \mathcal{R}^{(1)}$ is equipped with the partition $\alpha = \alpha'_u \cup \alpha'_\ell$, we satisfy the base case of the statement.



Figure 5 In both cases, \mathcal{W}_4 is disjoint to \mathcal{W}'_4 .

³⁸³ Case 2. The argument is essentially the same as Case 1. Suppose we have at least $|\mathcal{U}_3|^2/8$ ³⁸⁴ crossing pairs in $\mathcal{M}_3 \times \mathcal{M}_3$. Then by Theorem 5, there are subsets $\mathcal{M}_4, \mathcal{M}'_4 \subset \mathcal{M}_3$, each ³⁸⁵ of size $\Omega_{\delta}(n)$, such that every curve in \mathcal{M}_4 crosses every curve in \mathcal{M}'_4 . Let $\mathcal{U}_4 \subset \mathcal{U}$ be the ³⁸⁶ curves that corresponds to \mathcal{M}_4 and let $\mathcal{U}'_4 \subset \mathcal{U}$ be the curves that corresponds to \mathcal{M}'_4 . Set

387 $\mathcal{W}_4 = \{ \alpha_w : \alpha \in \mathcal{U}_4, \alpha = \alpha_w \cup \alpha_m \}$ and $\mathcal{W}'_4 = \{ \alpha_w : \alpha \in \mathcal{U}'_4, \alpha = \alpha_w \cup \alpha_m \}.$

See Figure 5b. Hence, by the pseudo-segment condition, every curve in W_4 is disjoint from every curve in W'_4 . By taking arbitrary subsets of \mathcal{M}_4 and \mathcal{B}_2 of size min $(|\mathcal{M}_4|, |\mathcal{B}_2|)$, we can apply Theorem 12 to these subsets and obtain subsets $\mathcal{M}_5 \subset \mathcal{M}_4$ and $\mathcal{B}_3 \subset \mathcal{B}_2$, each of size $\Omega_{\delta}(n)$, such that either every red curve in \mathcal{M}_5 crosses every blue curve in \mathcal{B}_3 , or every red curve in \mathcal{M}_5 is disjoint from every blue curve in \mathcal{B}_3 . In the former case, we are done. Hence, we can assume that we are in the latter case.

Again, we take an arbitrary subset of \mathcal{M}'_4 and \mathcal{B}_3 of size $\min(|\mathcal{M}'_4|, |\mathcal{B}_3|)$ and apply Theorem 12 to \mathcal{M}'_4 and \mathcal{B}_3 , to obtain subsets $\mathcal{M}'_5 \subset \mathcal{M}'_4$ and $\mathcal{B}_4 \subset \mathcal{B}_3$, each of size $\Omega_{\delta}(n)$, such that either every red curve in \mathcal{M}'_5 crosses every blue curve in \mathcal{B}_4 , or every red curve in \mathcal{M}'_5 is disjoint from every blue curve in \mathcal{B}_4 . Again, if we are in the former case, we are done. Hence, we can assume that we are in the latter case. Set \mathcal{R}_5 be the red curves in \mathcal{R} that corresponds to \mathcal{M}_5 , and let \mathcal{R}'_5 be the red curves in \mathcal{R} that corresponds to \mathcal{M}'_5 .

We have $|\mathcal{R}_5|, |\mathcal{R}'_5| = \Omega_{\delta}(n)$, and moreover, we can assume that $|\mathcal{R}_5| = |\mathcal{R}'_5|$. For each curve $\alpha \in \mathcal{R}_5 \cup \mathcal{R}'_5$, and its original partition $\alpha = \alpha_u \cup \alpha_\ell$ defined by γ_1 , we have a new partition $\alpha = \alpha'_u \cup \alpha'_\ell$ defined by γ_2 , where $\alpha'_u = \alpha_w$ and $\alpha'_\ell = \alpha_m \cup \alpha_\ell$. By setting $\mathcal{R}^{(1)} = \mathcal{R}_5 \cup \mathcal{R}'_5$, and $\mathcal{B}^{(1)} = \mathcal{B}_4$, where each curve $\alpha \in \mathcal{R}^{(1)}$ is equipped with the partition $\alpha = \alpha'_u \cup \alpha'_\ell$, we satisfy the base case of the statement.

For the inductive step, suppose we have obtained constants $\varepsilon'_{t-1} < \cdots < \varepsilon'_1$ such that 408 the statement follows. Let ε'_t be a small constant that will be determined later such that 409 $\varepsilon'_t < \varepsilon'_{t-1}$. Let \mathcal{R} be a set of n red curves in the plane, all crossed by a curve γ_1 exactly 410 once, and \mathcal{B} be a set of n blue curves in the plane such that $\mathcal{R} \cup \mathcal{B} \cup \{\gamma_1\}$ is a collection of 411 pseudo-segments. Moreover, $G(\mathcal{R})$ has edge density at least $1 - \varepsilon'_t$ and $G(\mathcal{B})$ has edge density 412 less than ε'_t . We set $\delta' < 0$ to be a small constant such that $\varepsilon'_t < \delta' < \varepsilon_{t-1}$. We repeat the 413 entire argument above, replacing ε'_1 with ε'_t and δ with δ' , to obtain subsets $\mathcal{R}_5, \mathcal{R}'_5 \subset \mathcal{R}$ 414 and $\mathcal{B}_4 \subset \mathcal{B}$, each of size $\Omega_{\delta'}(n)$, such that each $\alpha \in \mathcal{R}_5 \cup \mathcal{R}'_5$ is equipped with the partition 415 $\alpha = \alpha'_u \cup \alpha'_\ell$, and α'_ℓ is disjoint to every blue curve in \mathcal{B}_4 . Moreover, for $\alpha \in \mathcal{R}_5$ and $\beta \in \mathcal{R}'_5$, 416 where $\alpha = \alpha'_u \cup \alpha'_\ell$ and $\beta = \beta'_u \cup \beta'_\ell$, α'_u is disjoint to β'_u . 417

Since $|\mathcal{R}_5|, |\mathcal{B}_4| \geq \delta_1 n$, where δ_1 depends only on δ' , by Theorem 7, $G(\mathcal{R}_5)$ has edge density at least $1 - 2\varepsilon'_t/\delta_1^2$ and $G(\mathcal{B}_4)$ has edge density less than $2\varepsilon'_t/\delta_1^2$. By setting ε'_t sufficiently small, $G(\mathcal{R}_5)$ has edge density at least $1 - \varepsilon'_{t-1}$, and $G(\mathcal{B}_4)$ has edge density less

XX:12 A structure theorem for pseudo-segments and its applications

than ε'_{t-1} . By averaging, we can find subsets of \mathcal{R}_5 and \mathcal{B}_4 , each of size min $(|\mathcal{R}_5|, |\mathcal{B}_4|)$ and 421 with densities at least $1 - \varepsilon'_{t-1}$ and less than ε'_{t-1} respectively, and apply induction to these 422 subsets parameter t' = t - 1, and obtain subsets $\mathcal{R}^{(t-1)} \subset \mathcal{R}_5$, $\mathcal{B}^{(t-1)} \subset \mathcal{B}_4$, each of size 423 $\Omega_{\varepsilon'_{+-1}}(n)$, with the desired properties. If every red curve in $\mathcal{R}^{(t-1)}$ is disjoint from every blue 424 curve in $\mathcal{B}^{(t-1)}$, or if every red curve in $\mathcal{R}^{(t-1)}$ crosses every blue curve in $\mathcal{B}^{(t-1)}$, then we 425 are done. Hence, we can assume that each curve $\alpha \in \mathcal{R}^{(t-1)}$ has a partition $\alpha = \alpha''_{\mu} \cup \alpha''_{\ell}$ 426 such that α''_u is a subcurve of α'_u , α''_ℓ is disjoint from every blue curve in $\mathcal{B}^{(t-1)}$, and there is an equipartition $\mathcal{R}^{(t-1)} = \mathcal{R}^{(t-1)}_1 \cup \cdots \cup \mathcal{R}^{(t-1)}_{2^{t-1}}$, such that for $1 \leq i < j \leq 2^{t-1}$, the upper 427 428 part α''_u of each curve $\alpha \in \mathcal{R}_i^{(t-1)}$ is disjoint the upper part β''_u of each curve $\beta \in \mathcal{R}_j^{(t-1)}$. 429

Finally, since $|\mathcal{R}'_5|, |\mathcal{B}^{(t-1)}| \geq \delta_2 n$, where δ_2 depends only on δ' , by Theorem 7, $G(\mathcal{R}'_5)$ 430 has edge density at least $1 - 2\varepsilon'_t/\delta_2^2$ and $G(\mathcal{B}^{(t-1)})$ has edge density less than $2\varepsilon'_t/\delta_2^2$. By 431 setting ε'_t sufficiently small, $G(\mathcal{R}'_5)$ has edge density at least $1 - \varepsilon'_{t-1}$, and $G(\mathcal{B}^{(t-1)})$ has 432 edge density less than ε'_{t-1} . By averaging, we can find subsets of \mathcal{R}'_5 and $\mathcal{B}^{(t-1)}$, each of 433 size $\min(|\mathcal{R}'_5|, |\mathcal{B}^{(t-1)}|)$ and with densities at least $1 - \varepsilon'_{t-1}$ and less than ε'_{t-1} respectively, 434 and apply induction to these subsets parameter t' = t - 1, and obtain subsets $\mathcal{S}^{(t-1)} \subset \mathcal{R}'_5$, 435 $\mathcal{B}^{(t)} \subset \mathcal{B}^{(t-1)}$, each of size $\Omega_{\varepsilon'_{t-1}}(n)$, with the desired properties. If every red curve in $\mathcal{S}^{(t-1)}$ 436 is disjoint from every blue curve in $\mathcal{B}^{(t)}$, or if every red curve in $\mathcal{S}^{(t-1)}$ crosses every blue 437 curve in $\mathcal{B}^{(t)}$, then we are done. Hence, we can assume that each curve $\alpha \in \mathcal{S}^{(t-1)}$ has 438 a partition $\alpha = \alpha''_u \cup \alpha''_\ell$ such that α''_u is a subcurve of α'_u , α''_ℓ is disjoint from every blue curve in $\mathcal{B}^{(t-1)}$, and there is an equipartition $\mathcal{S}^{(t-1)} = \mathcal{S}^{(t-1)}_1 \cup \cdots \cup \mathcal{S}^{(t-1)}_{2^{t-1}}$, such that for 439 440 $1 \leq i < j \leq 2^{t-1}$, the upper part α''_u of each curve $\alpha \in \mathcal{S}_i^{(t-1)}$ is disjoint the upper part β''_u of 441 each curve $\beta \in \mathcal{S}_i^{(t-1)}$. We then (arbitrarily) remove curves from each part in $\mathcal{R}_i^{(t-1)}$ and 442 $\mathcal{S}_{i}^{(t-1)}$ such that the resulting parts all have the same size and for

444
$$\mathcal{R}^{(t)} = \mathcal{R}_1^{(t-1)} \cup \dots \cup \mathcal{R}_{2^{t-1}}^{(t-1)} \cup \mathcal{S}_1^{(t-1)} \cup \dots \cup \mathcal{S}_{2^{t-1}}^{(t-1)},$$

we have $|\mathcal{R}^{(t)}| = \Omega_{\varepsilon'_{4-1}}(n)$. Then $\mathcal{R}^{(t)}$ and $\mathcal{B}^{(t)}$ has the desired properties.

◄

446 We now prove the following.

⁴⁴⁷ ► **Theorem 17.** There is an absolute constant $\varepsilon_3 > 0$ such that the following holds. Let \mathcal{R} ⁴⁴⁸ be a set of n red curves in the plane and \mathcal{B} be a set of n blue curves in the plane such that ⁴⁴⁹ $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-segments, and the intersection graph $G(\mathcal{B})$ has edge density ⁴⁵⁰ less than ε_3 , and $G(\mathcal{R})$ has edge density at least $1 - \varepsilon_3$. Then there are subsets $\mathcal{R}' \subset \mathcal{R}$, ⁴⁵¹ $\mathcal{B}' \subset \mathcal{B}$, each of size $\Omega(n)$, such that either every red curve in \mathcal{R} crosses every blue curve in ⁴⁵² \mathcal{B} , or every red curve in \mathcal{R} is disjoint from every blue curve in \mathcal{B} .

Proof. Let t be a fixed large integer such that $2^{-t} < \varepsilon_1$, where ε_1 is defined in Theorem 453 15. Let ε_3 be a small constant determined later such that $\varepsilon_3 < \varepsilon'_t$, where ε'_t is defined in 454 Lemma 16. Recall that $\varepsilon'_t < \varepsilon_1$. Since $G(\mathcal{R})$ has edge density at least $1 - \varepsilon_3$, there is a curve 455 $\gamma_1 \in \mathcal{R}$ such that γ_1 crosses at least n/2 red curves in \mathcal{R} . Let $\mathcal{R}_0 \subset \mathcal{R}$ be the red curves that 456 crosses γ_1 . By Lemma 7, $G(\mathcal{R}_0)$ has edge density at least $1 - 8\varepsilon_3$. By averaging, we can find 457 a subset $\mathcal{B}' \subset \mathcal{B}$ of size $|\mathcal{R}_0|$ whose edge density is less than ε_3 . By setting ε_3 sufficiently 458 small so that $8\varepsilon_3 < \varepsilon'_t$, we can apply Lemma 16 to \mathcal{R}_0 and \mathcal{B}' with parameter t, and obtain 459 subsets $\hat{\mathcal{R}} \subset \mathcal{R}_0, \hat{\mathcal{B}} \subset \mathcal{B}$, each of size $\Omega_{\varepsilon'}(n)$, with the desired properties. If every red curve 460 in $\hat{\mathcal{R}}$ crosses every blue curve in $\hat{\mathcal{B}}$, or every red curve in $\hat{\mathcal{R}}$ is disjoint from every blue curve 461 in $\hat{\mathcal{B}}$, then we are done. Therefore, we can assume that each curve $\alpha \in \hat{\mathcal{R}}$ has a partition 462 into two parts $\alpha = \alpha'_{\mu} \cup \alpha'_{\ell}$ with the properties described in Lemma 16. Set 463

464 $\mathcal{U} = \{ \alpha'_u : \alpha \in \hat{\mathcal{R}}, \alpha = \alpha'_u \cup \alpha'_\ell \}$ and $\mathcal{L} = \{ \alpha'_\ell : \alpha \in \hat{\mathcal{R}}, \alpha = \alpha'_u \cup \alpha'_\ell \}.$

Hence, every curve in \mathcal{L} is disjoint from every curve in $\hat{\mathcal{B}}$, and $G(\mathcal{U})$ has edge density at 465 most $2^{-t} < \varepsilon_1$. Since $|\hat{\mathcal{B}}| \ge \delta n$, where δ depends only on ε'_t , by Lemma 7, $G(\hat{\mathcal{B}})$ has edge 466 density at most $2\varepsilon_3/\delta^2$. By setting ε_3 sufficiently small so that $2\varepsilon_3/\delta_0^2 < \varepsilon_1$, $G(\hat{\mathcal{B}})$ has edge 467 density at most ε_1 . By averaging, we can find subsets of \mathcal{U} and $\hat{\mathcal{B}}$, each of size min $(|\mathcal{U}|, |\hat{\mathcal{B}}|)$ 468 and with densities at most ε_1 , and apply Theorem 15 to these subsets to obtain subsets 469 $\mathcal{U}' \subset \mathcal{U}$ and $\mathcal{B}' \subset \hat{\mathcal{B}}$, each of size $\Omega_{\varepsilon_3}(n)$, such that every curve in \mathcal{U}' is disjoint from every 470 curve in \mathcal{B}' , or every curve in \mathcal{U}' crosses every curve in \mathcal{B}' . By setting \mathcal{R}' to be the red curves 471 in \mathcal{R} corresponding to \mathcal{U}' , every red curve in \mathcal{R}' is disjoint from every blue curve in \mathcal{B}' , or 472 every red curve in \mathcal{R}' crosses every blue curve in \mathcal{B}' , and each subset has size $\Omega_{\varepsilon_3}(n)$. 473

474 4.3 High versus high edge density

Finally, we consider the case when the intersection graphs $G(\mathcal{R})$ and $G(\mathcal{B})$ both have edge densities at least $1 - \varepsilon$. By copying the proof of Theorem 17, except using Theorem 17 (high versus low density) instead of Theorem 15 (low versus low density), we obtain the following.

▶ Theorem 18. There is an absolute constant $ε_4 > 0$ such that the following holds. Let \mathcal{R} be a set of n red curves in the plane and \mathcal{B} be a set of n blue curves in the plane such that $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-segments, and the intersection graphs $G(\mathcal{B})$ and $G(\mathcal{R})$ both have edge density at least $1 - ε_4$. Then there are subsets $\mathcal{R}' \subset \mathcal{R}$, $\mathcal{B}' \subset \mathcal{B}$, each of size $\Omega(n)$, such that either every red curve in \mathcal{R} crosses every blue curve in \mathcal{B} , or every red curve in \mathcal{R} is disjoint from every blue curve in \mathcal{B} .

⁴⁸⁴ **5** Proof of Theorem 4

Let \mathcal{R} be a set of n red curves in the plane, and \mathcal{B} be a set of n blue curves in the plane 485 such that $\mathcal{R} \cup \mathcal{B}$ is a collection of pseudo-segments. Let ε be a sufficiently small constant 486 such that $\varepsilon < \varepsilon_4 < \varepsilon_3 < \varepsilon_1$, where ε_1 is from Theorem 15, ε_3 is from Theorem 17, and ε_4 487 is from Theorem 18. We apply Corollary 6 to both \mathcal{R} and \mathcal{B} and obtain subsets $\mathcal{R}_1 \subset \mathcal{R}$ 488 and $\mathcal{B}_1 \subset \mathcal{B}$ such that both $G(\mathcal{R}_1)$ and $G(\mathcal{B}_1)$ are ε -homogeneous. Moreover, we can assume 489 that $|\mathcal{R}_1| = |\mathcal{B}_1|$. If both $G(\mathcal{R}_1)$ and $G(\mathcal{B}_1)$ have edge densities less than ε , then, since ε is 490 sufficiently small, we can apply Theorem 15 to obtain subsets $\mathcal{R}_2 \subset \mathcal{R}_1$ and $\mathcal{B}_2 \subset \mathcal{B}_1$, each 491 of size $\Omega_{\varepsilon}(n)$, such that either every red curve in \mathcal{R}_2 is disjoint from every blue curve in \mathcal{B}_2 . 492 or every red curve in \mathcal{R}_2 crosses every blue curve in \mathcal{B}_2 . If one of the graphs of $G(\mathcal{R}_1)$ and 493 $G(\mathcal{B}_1)$ has edge density less than ε , and the other has edge density greater than $1-\varepsilon$, then 494 we apply Theorem 17 to \mathcal{R}_1 and \mathcal{B}_1 to obtain subsets $\mathcal{R}_2 \subset \mathcal{R}_1$ and $\mathcal{B}_2 \subset \mathcal{B}_1$, each of size 495 $\Omega_{\varepsilon}(n)$, such that either every red curve in \mathcal{R}_2 is disjoint from every blue curve in \mathcal{B}_2 , or every 496 red curve in \mathcal{R}_2 crosses every blue curve in \mathcal{B}_2 . Finally, if both $G(\mathcal{R}_1)$ and $G(\mathcal{B}_1)$ have edge 497 densities at least $1 - \varepsilon$, then, since ε is sufficiently small, we can apply Theorem 18 to obtain 498 subsets $\mathcal{R}_2 \subset \mathcal{R}_1$ and $\mathcal{B}_2 \subset \mathcal{B}_1$, each of size $\Omega_{\varepsilon}(n)$, such that either every red curve in \mathcal{R}_2 is 499 disjoint from every blue curve in \mathcal{B}_2 , or every red curve in \mathcal{R}_2 crosses every blue curve in \mathcal{B}_2 . 500

501 — References

Oswin Aichholzer, Alfredo García, Javier Tejel, Birgit Vogtenhuber, and Alexandra Weinberger.
 Twisted ways to find plane structures in simple drawings of complete graphs. In Xavier Goaoc
 and Michael Kerber, editors, 38th International Symposium on Computational Geometry,

XX:14 A structure theorem for pseudo-segments and its applications

- 505 SoCG 2022, June 7-10, 2022, Berlin, Germany, volume 224 of LIPIcs, pages 5:1–5:18. Schloss
- 506 Dagstuhl Leibniz-Zentrum für Informatik, 2022.
- Noga Alon, János Pach, Rom Pinchasi, Rados Radoicic, and Micha Sharir. Crossing patterns of semi-algebraic sets. J. Comb. Theory, Ser. A, 111(2):310–326, 2005.
- Grant Cairns and Yury Nikolayevsky. Bounds for generalized thrackles. Discret. Comput.
 Geom., 23(2):191–206, 2000.
- Jacob Fox and János Pach. A separator theorem for string graphs and its applications. Comb.
 Probab. Comput., 19(3):371–390, 2010.
- Jacob Fox, János Pach, and Andrew Suk. A polynomial regularity lemma for semialgebraic
 hypergraphs and its applications in geometry and property testing. SIAM J. Comput.,
 45(6):2199–2223, 2016.
- Jacob Fox, János Pach, and Andrew Suk. Enumeration of intersection graphs of x-monotone curves. In preparation, 2024+.
- Jacob Fox, János Pach, and Andrew Suk. A structure theorem for pseudo-segments and its applications. arxiv:2312.01028, 2024.
- Jacob Fox, János Pach, and Csaba D. Tóth. Intersection patterns of curves. J. Lond. Math.
 Soc., 83(2):389-406, 2011.
- Jacob Fox and Benny Sudakov. Density theorems for bipartite graphs and related fRamsey-type
 results. *Combinatorica*, 29(2):153–196, 2009.
- Radoslav Fulek and János Pach. A computational approach to Conway's thrackle conjecture.
 Comput. Geom., 44(6-7):345–355, 2011.
- Radoslav Fulek and János Pach. Thrackles: An improved upper bound. Discret. Appl. Math.,
 259:226-231, 2019.
- Jacob E. Goodman. Proof of a conjecture of Burr, Grünbaum, and Sloane. Discret. Math.,
 32(1):27–35, 1980.
- Jacob E. Goodman and Richard Pollack. Semispaces of configurations, cell complexes of arrangements. J. Comb. Theory, Ser. A, 37(3):257–293, 1984.
- Sariel Har-Peled. Constructing planar cuttings in theory and practice. SIAM J. Comput.,
 29(6):2016–2039, 2000.
- János Komlós, Ali Shokoufandeh, Miklós Simonovits, and Endre Szemerédi. The regularity
 lemma and its applications in graph theory. In Gholamreza B. Khosrovshahi, Ali Shokoufandeh,
 and Mohammad Amin Shokrollahi, editors, *Theoretical Aspects of Computer Science, Advanced Lectures (First Summer School on Theoretical Aspects of Computer Science, Tehran, Iran, July 2000)*, volume 2292 of *Lecture Notes in Computer Science*, pages 84–112. Springer, 2000.
- László Lovász, János Pach, and Mario Szegedy. On Conway's thrackle conjecture. Discret.
 Comput. Geom., 18(4):369-376, 1997.
- János Pach and József Solymosi. Structure theorems for systems of segments. In Jin Akiyama, Mikio Kano, and Masatsugu Urabe, editors, *Discrete and Computational Geometry, Japanese Conference, JCDCG 2000, Tokyo, Japan, November, 22-25, 2000, Revised Papers*, volume
 2098 of *Lecture Notes in Computer Science*, pages 308–317. Springer, 2000.
- János Pach and József Solymosi. Crossing patterns of segments. J. Comb. Theory, Ser. A, 96(2):316-325, 2001.

János Pach and Géza Tóth. Disjoint edges in topological graphs. In Jin Akiyama, Edy Tri
 Baskoro, and Mikio Kano, editors, Combinatorial Geometry and Graph Theory, Indonesia Japan Joint Conference, IJCCGGT 2003, Bandung, Indonesia, September 13-16, 2003, Revised
 Selected Papers, volume 3330 of Lecture Notes in Computer Science, pages 133–140. Springer,

- 2003.
 200 Lisa Sauermann. On the speed of algebraically defined graph classes. Adv. Math., 380:107593, 2021.
 - Andrew Suk. Density theorems for intersection graphs of t-monotone curves. SIAM J. Discret.
 Math., 27(3):1323-1334, 2013.