## A geometric approach to Wigner-type theorems

Mark Pankov

University of Warmia & Mazury, Poland and JKU Linz, Austria

Let H be a complex Hilbert space and let  $\mathcal{P}(H)$  be the associated projective space (the set of rank-one projections). Suppose that dim  $H \geq 3$ . We prove the following Wigner-type theorem: if H is finite-dimensional, then an arbitrary orthogonality preserving transformation of  $\mathcal{P}(H)$  (i.e. sending orthogonal projections to orthogonal projections without the assumption that the orthogonality relation is preserved in both directions) is induced by a unitary or anti-unitary operator. In the case when H is infinite-dimensional, this fails.

The problem is reduced to a description of orthogonality preserving lineations. Lineations are maps between projective spaces which send lines to subsets of lines. In general, the behavior of lineations is complicated; they are not injective and can send lines to parts of lines only. Our version of Wigner's theorem is a consequence of the following result: every orthogonality preserving lineation of  $\mathcal{P}(V)$  to itself is induced by a linear or conjugate-linear isometry (now, we do not require that H is finite-dimensional).

As an application, we describe (not necessarily injective) transformations of Grassmannians preserving some types of principal angles.

This is a joint work with Thomas Vetterlein.

16 Sep 2020, 12:00 (BST)

You can join the event via this link: https://istaustria.zoom.us/j/97142391866?pwd=U0VKemxUdGhyV04rY3NqL0VnNHp4dz09 Meeting ID: 971 4239 1866 Passcode: 823120

More information about the webinar series can be found at the following homepages https://www.renyi.hu/~titkos/preserverwebinar.html https://researchseminars.org/seminar/PreserverWebinar

The slides and video recordings of all previous talks are available in this google drive folder: drive.google.com/drive/folders/1FZJ6KP1b6dpLrbY2wJMoqj3hYK-IYomx?usp=sharing